The Journal of Open Source Software

CTSEG: A segment picture quantum impurity solver based on TRIQS

Nikita Kavokine [1,2], **Hao Lu** [2], **Thomas Ayral** [3], **Michel Ferrero** [4,5], **Nils Wentzell** [1], and **Olivier Parcollet** [1]

1 Center for Computational Quantum Physics, Flatiron Institute, 162 5th Avenue, NY 10010, New York, USA **2** Max Planck Institute for Polymer Research, Ackermannweg 10, 55128 Mainz, Germany **3** Eviden Quantum Laboratory, 78340 Les Clayes-sous-Bois, France **4** CPHT, CNRS, Ecole Polytechnique, Institut Polytechnique de Paris, 91128 Palaiseau, France **5** Collège de France, 11 place Marcelin Berthelot, 75005 Paris, France

DOI: 10.xxxxxx/draft

Software

- Review
- Repository
- Archive

Editor: Lucy Whalley

Reviewers:

- @HugoStrand
- @egcpvanloon

Submitted: 18 September 2024
Published: unpublished

License
Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License (CC BY 4.0).

DRAFT

Summary

Electron-electron interactions are a key determinant of the electrical and optical response properties in solid-state materials. Yet, the many-electron problem is outstandingly difficult, and can be tackled analytically only when interactions are weak, or in rare exactly solvable cases. One of the popular numerical schemes for addressing strongly interacting systems is dynamical mean-field theory (DMFT) (Georges et al., 1996). In DMFT, the many-body problem formulated on a crystal lattice is self-consistently mapped onto a single atom (or impurity) immersed in an effective environment (or bath). The remaining computational task is then the solution of the impurity problem, which can be carried out through various quantum Monte Carlo algorithms (Gull et al., 2011). Here, we present an implementation of the continous time hybridization expansion algorithm in the segment picture (`CTSEG`) based on `TRIQS`, a comprehensive library for the numerical investigation of interacting quantum systems (Parcollet et al., 2015).

Statement of need

The Monte Carlo algorithms for quantum impurity problems are based on stochastically exploring the terms in the perturbative expansion of the solution around an exactly solvable limit. Continuous time hybridization expansion algorithms – chief of which `CTHYB` – involve expanding around the limit of an isolated atom (Gull et al., 2011). Currently, there exist implementations of `CTHYB` within three different libraries: `ALPS` (Shinaoka et al., 2017), `w2dynamics` (Wallerberger et al., 2019) and `TRIQS` (Seth et al., 2016).

However, a simpler and potentially faster version of the `CTHYB` algorithm, called `CTSEG`, can be used under the restriction of density-density interactions on the impurity (Haule, 2007; Werner et al., 2006; Werner & Millis, 2006). `CTSEG` can be further generalized to allow for time-dependent(Werner & Millis, 2007, 2010) and spin-spin interactions (Otsuki, 2013; Steiner et al., 2015): see `@werner2016` for a review. To our knowledge, there exists so far one published implementation of `CTSEG` based on ALPS (Hafermann et al., 2013), but it does not allow for spin-spin interactions.

Our `CTSEG` solver is about twice as fast as `TRIQS-CTHYB` for a single orbital problem, and has better scaling with the number of orbitals (40 times faster in our 5 orbital test case, see Fig. 1a). `CTSEG` has already allowed us to obtain the first numerically-exact solution of the quantum Heisenberg spin glass (Kavokine et al., 2024).

The Journal of Open Source Software



Figure 1: **a**. Running time comparison between the TRIQS implementations of CTSEG and CTHYB. The test system is a multi-orbital impurity at half-filling and inverse temperature $\beta = 20$. The Coulomb repulsion is $U = 2$ for two electrons on the same orbital and $U' = 1$ for two electrons on different orbitals. The hybridization is diagonal and identical for all orbitals: $\Delta(\omega) = 1/(\omega - 0.3)$. **b**. Spin-spin correlation function $\chi(\tau) = \langle \mathbf{S}(\tau) \cdot \mathbf{S}(0) \rangle$ of the $t - J - U$ model studied by Dumitrescu et al., obtained using CTSEG at inverse temperature $\beta = 300$ and different values of doping p. At long times $\chi(\tau) \sim 1/\tau^{\theta}$, with $\theta = 1$ at the QCP. Inset: exponent θ as a function of doping p. The QCP is located at $p \approx 0.16$.

Example of use

As a further illustration of our solver's performance, we apply it to the fully connected $t-J-U$ model studied by Dumitrescu et al. (2022). At half-filling, the model forms a spin glass phase, which melts into a metal at a doping-induced quantum critical point (QCP). Dumitrescu et al. obtained solutions at inverse temperatures up to $\beta = 65$, limited by the fermionic sign problem of their interaction expansion solver. The hybridization expansion carried out by CTSEG is sign-problem-free for the $t - J - U$ model, allowing us to reach $\beta = 300$ and to obtain a

more accurate localization of the QCP at doping $p \approx 0.16$ (Fig. 1b).

Acknowledgements

We thank Alexander Hampel for testing the code and providing valuable feedback. The Flatiron Institute is a division of the Simons Foundation.

References

Dumitrescu, P. T., Wentzell, N., Georges, A., & Parcollet, O. (2022). Planckian metal at a doping-induced quantum critical point. *Phys. Rev. B*, *105*, L180404. https://doi.org/10.1103/PhysRevB.105.L180404

Georges, A., Kotliar, G., Krauth, W., & Rozenberg, M. J. (1996). Dynamical mean-field theory of strongly correlated fermion systems and the limit of infinite dimensions. *Rev. Mod. Phys.*, *68*(1), 13–125. https://doi.org/10.1103/RevModPhys.68.13

Gull, E., Millis, A. J., Lichtenstein, A. I., Rubtsov, A. N., Troyer, M., & Werner, P. (2011). Continuous-time Monte Carlo methods for quantum impurity models. *Rev. Mod. Phys.*, *83*(2), 349–404. https://doi.org/10.1103/RevModPhys.83.349

Hafermann, H., Werner, P., & Gull, E. (2013). Efficient implementation of the continuous-time hybridization expansion quantum impurity solver. *Computer Physics Communications*, *184*(4), 1280–1286. https://doi.org/10.1016/j.cpc.2012.12.013

Haule, K. (2007). Quantum monte carlo impurity solver for cluster dynamical mean-field theory and electronic structure calculations with adjustable cluster base. *Phys. Rev. B*, *75*, 155113. https://doi.org/10.1103/PhysRevB.75.155113

Kavokine, N., Müller, M., Geroges, A., & Parcollet, O. (2024). Exact numerical solution of the fully connected classical and quantum Heisenberg spin glass. *Phys. Rev. Lett.*, *133*, 016501. https://doi.org/10.1103/PhysRevLett.133.016501

Otsuki, J. (2013). Spin-boson coupling in continuous-time quantum Monte Carlo. *Phys. Rev. B*, *87*(12), 1–7. https://doi.org/10.1103/PhysRevB.87.125102

Parcollet, O., Ferrero, M., Ayral, T., Hafermann, H., Krivenko, I., Messio, L., & Seth, P. (2015). TRIQS: A toolbox for research on interacting quantum systems. *Computer Physics Communications*, *196*, 398–415. https://doi.org/10.1016/j.cpc.2015.04.023

Seth, P., Krivenko, I., Ferrero, M., & Parcollet, O. (2016). TRIQS/CTHYB: A continuous-time quantum Monte Carlo hybridisation expansion solver for quantum impurity problems. *Computer Physics Communications*, *200*, 274–284. https://doi.org/10.1016/j.cpc.2015.10.023

Shinaoka, H., Gull, E., & Werner, P. (2017). Continuous-time hybridization expansion quantum impurity solver for multi-orbital systems with complex hybridizations. *Computer Physics Communications*, *215*, 128–136. https://doi.org/10.1016/j.cpc.2017.01.003

Steiner, K., Nomura, Y., & Werner, P. (2015). Double-expansion impurity solver for multiorbital models with dynamically screened U and J. *Phys. Rev. B*, *92*, 115123. https://doi.org/10.1103/PhysRevB.92.115123

Wallerberger, M., Hausoel, A., Gunacker, P., Kowalski, A., Parragh, N., Goth, F., Held, K., & Sangiovanni, G. (2019). w2dynamics: Local one- and two-particle quantities from dynamical mean field theory. *Computer Physics Communications*, *235*, 388–399. https://doi.org/10.1016/j.cpc.2018.09.007

Werner, P., Comanac, A., Medici, L. de', Troyer, M., & Millis, A. J. (2006). Continuous-time

DRAFT

solver for quantum impurity models. *Phys. Rev. Lett.*, *97*, 076405. https://doi.org/10.1103/PhysRevLett.97.076405

Werner, P., & Millis, A. J. (2006). Hybridization expansion impurity solver: General formulation and application to kondo lattice and two-orbital models. *Phys. Rev. B*, *74*, 155107. https://doi.org/10.1103/PhysRevB.74.155107

Werner, P., & Millis, A. J. (2007). Efficient dynamical mean field simulation of the holstein-hubbard model. *Phys. Rev. Lett.*, *99*, 146404. https://doi.org/10.1103/PhysRevLett.99.146404

Werner, P., & Millis, A. J. (2010). Dynamical screening in correlated electron materials. *Phys. Rev. Lett.*, *104*, 146401. https://doi.org/10.1103/PhysRevLett.104.146401

DRAFT