The Journal of Open Source Software

pDEMtools: conveniently search, download, and process ArcticDEM and REMA products

Thomas R. Chudley [1¶] **and Ian M. Howat** [2,3]

1 Department of Geography, Durham University, Durham, UK **2** Byrd Polar and Climate Research Center, Ohio State University, Columbus, OH, USA **3** School of Earth Sciences, Ohio State University, Columbus, OH, USA ¶ Corresponding author

DOI: 10.xxxxxx/draft

Software

- Review
- Repository
- Archive

Editor:

Submitted: 14 May 2024
Published: unpublished

License
Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License (CC BY 4.0).

DRAFT

Summary

pdemtools is a Python package designed for accessing, processing, and handling high-resolution Digital Elevation Models (DEMs) of the polar regions from the ArcticDEM and Reference Elevation Model of Antarctica (REMA) projects. Tools are provided to search, filter, and download ArcticDEM and REMA data, as well as to fulfill common preprocessing requirements such as geoid correction, coregistration, and calculating terrain attributes. The aim is to allow users to quickly move beyond basic DEM data management into their own analyses.

Statement of need

ArcticDEM and REMA are high-resolution, time-stamped 2-metre-resolution DEMs of the polar regions provided by the Polar Geospatial Center (PGC). They are extracted by applying stereo auto-correlation techniques (Noh & Howat, 2017) to pairs of submetre Maxar satellite imagery, including Worldview-1, Worldview-2, Worldview-3, and GeoEye-1, beginning in 2007 (ArcticDEM) or 2009 (REMA) and ongoing to the present day. Products are available as tens of thousands of time-stamped 'strips' (Howat et al., 2022b; Porter et al., 2022) constructed from individual scene-pairs, or as a single mosaic (Howat et al., 2022a; Porter et al., 2023) compiled from the combined stack of strips. Strips allow users to perform change detection by comparing data from different seasons or years, whilst mosaics provide a consistent and comprehensive product over the entire polar regions.

As Earth Science has moved into the 'big data' era, increasing amounts of Arctic- and Antarctic-focused resources are available as public, cloud-optimised datasets. New approaches are providing Python tools to act as combined API and processing tools, such as `icepyx` (Scheick et al., 2023) or `pypromice` (How et al., 2023). From 2022 (ArcticDEM v4.1 and REMA v2), the PGC DEM products are hosted as Cloud Optimised GeoTIFFs (CoGs) in a SpatioTemporal Asset Catalog (STAC), a standardised structure for cataloguing spatiotemporal data. However, the PGC STAC is not currently a dynamic STAC, and as such lacks a RESTful API interface for searching and downloading datasets in response to user queries. This limits the ability of users to programmatically interact with ArcticDEM and REMA data in a quick and efficient manner. The `pdemtools` package has two aims: the first is to provide a Python-focussed alternative for searching and downloading ArcticDEM and REMA data, emulating dynamic STAC query tools such as `pystac` (Radiant Earth, 2017 - 2024); whilst the second is to provide commonly used processing functions specific to the needs of ArcticDEM and REMA users (a focus on ice sheet and cryosphere work), as well as the particular strengths of the ArcticDEM and REMA datasets (high-resolution and multitemporal).

The `pdemtools search()` tool and `load` module allow for convenient access to the ArcticDEM and REMA datasets. Mosaics can be downloaded from a one-line `tools.mosaic()` function,

whilst the `search()` function allows for convenient filtering of a locally downloading ArcticDEM/REMA strip index according to variables such as date, region of interest, spatial coverage, temporal baseline, source sensors, accuracy, and cross-track data. The results of searches are returned as a `geopandas` (Van den Bossche et al., 2016 - 2024) dataframe, and can be downloaded using the `load.from_search()` function. Elevation models are returned as `xarray` DataArrays (Hoyer & Hamman, 2017) with geospatial metadata via the `rioxarray` extension (Corteva, 2019 - 2024) - a standard format for storing and processing n-dimensional geospatial data within the Python community. By utilising standardised formats, the aim is to allow the user to quickly move beyond `pdemtools` into their own analysis in whatever format they desire, be that `xarray`, `numpy` or `dask` datasets, DEM analysis Python packages such as `xdem` (GlacioHack Community, 2020 - 2023) for advanced coregistration or `richdem` (Barnes, 2016 - 2022) for flow analysis, or exporting to geospatial file formats for analysis beyond Python.

After download, there exist a number of (pre-)processing steps that are near universally common in topographic analyses. These include geoid-correction, co-registration of time-series data, and/or the construction of terrain parameters such as hillshade, slope, aspect, and curvature. `pdemtools` contains pre-built functions to perform these processing steps, as well as further functionality specific to ArcticDEM and REMA use cases. For instance, we include functions to quickly extract the EIGEN-6C4 geoid (Foerste et al., 2014) and Greenland/Antartctic bedrock masks directly from local versions of the Greenland and Antarctic BedMachine datasets (Morlighem, 2020a, 2020b), reprojecting and resampling the data to match the target DEM. Options for ingesting user-provided data are also provided. Additionally, partial deriviatives of the surface used to calculate terrain parameters ($\frac{\partial z}{\partial x}$, $\frac{\partial z}{\partial y}$, $\frac{\partial^2 z}{\partial x^2}$, $\frac{\partial^2 z}{\partial y^2}$, $\frac{\partial^2 z}{\partial x \partial y}$) are calculated following Florinsky (2009), as opposed to more common methods such as Zevenbergen & Thorne (1987). The newer approach computes partial derivatives of elevation based on fitting a third-order polynomial, by the least-squares approach, to a 5×5 window as opposed to the more common 3×3 window. This is more appropriate for high-resolution DEMs: curvature over a 10 m window for the 2 m resolution ArcticDEM/REMA strips will lead to a local denoising effect that limits the impact of noise common in high-resolution photogrammetric products. These methods are also adapted into our co-registration routine, which otherwise follows the commonly used approach of Nuth & Kääb (2011).

We aim to grow `pdemtools` by implementing new methods developed by the ArcticDEM and REMA research community. For instance, we currently include sea-level-filtering and iceberg detection routines outlined by Shiggins et al. (2023), and invite community contributions or requests of other routines that will be of use to users of `pdemtools`. Ongoing research projects making use of `pdemtools` are applying ArcticDEM and REMA data to the mapping of crevasses, ice cliff heights, and subglacial lakes, as well as the initiation of ice sheet models. It has also been used within training exercises at the 2024 Polar Geospatial Center Data Workshop, contributing to a growing international network of `pdemtools` users.

Documentation

Package documentation is available at the `pdemtools` readthedocs.

Acknowledgements

Chudley is supported by a Leverhulme Early Career Fellowship (ECF-2022-589). ArcticDEM data are provided by the Polar Geospatial Center under NSF-OPP awards 1043681, 1559691, and 1542736. REMA data are provided by the Byrd Polar and Climate Research Center and the Polar Geospatial Center under NSF-OPP awards 1543501, 1810976, 1542736, 1559691, 1043681, 1541332, 0753663, 1548562, 1238993 and NASA award NNX10AN61G. Computer time provided through a Blue Waters Innovation Initiative. DEMs produced using data from Maxar. We are grateful to the PGC for making index strips in the parquet format available

DRAFT

from their website; to Connor Shiggins for his input on the iceberg detection function; to Andrew Sole and Adrien Wehrle for identifying and helping to resolve bugs in early versions of `pdemtools`; and to the colleagues who helped to test this package in the early phases of development.

References

Barnes, R. (2016 - 2022). *RichDEM: Terrain analysis software* (Version 2.3.1). http://github.com/r-barnes/richdem

Corteva. (2019 - 2024). *Rioxarray: Geospatial xarray extension powered by rasterio* (Version 0.15.5). https://github.com/corteva/rioxarray

Florinsky, I. V. (2009). Computation of the third-order partial derivatives from a digital elevation model. *International Journal of Geographical Information Science*, *23*(2), 213–231. https://doi.org/10.1080/13658810802527499

Foerste, C., Bruinsma, Sean. L., Abrykosov, O., Lemoine, J.-M., Marty, J. C., Flechtner, F., Balmino, G., Barthelmes, F., & Biancale, R. (2014). *EIGEN-6C4 the latest combined global gravity field model including GOCE data up to degree and order 2190 of GFZ potsdam and GRGS toulouse*. GFZ Data Services. https://doi.org/10.5880/icgem.2015.1

GlacioHack Community. (2020 - 2023). *xDEM: Analysis of digital elevation models* (Version 0.0.18). https://github.com/GlacioHack/xdem

How, P. R., Wright, P. J., Mankoff, K. D., Vandecrux, B., Fausto, R. S., & Ahlstrøm, A. P. (2023). Pypromice: A Python package for processing automated weather station data. *Journal of Open Source Software*, *8*(86), 5298. https://doi.org/10.21105/joss.05298

Howat, I., Porter, C., Noh, M.-J., Husby, E., Khuvis, S., Danish, E., Tomko, K., Gardiner, J., Negrete, A., Yadav, B., Klassen, J., Kelleher, C., Cloutier, M., Bakker, J., Enos, J., Arnold, G., Bauer, G., & Morin, P. (2022a). *The Reference Elevation Model of Antarctica - Mosaics, Version 2*. Harvard Dataverse. https://doi.org/10.7910/DVN/EBW8UC

Howat, I., Porter, C., Noh, M.-J., Husby, E., Khuvis, S., Danish, E., Tomko, K., Gardiner, J., Negrete, A., Yadav, B., Klassen, J., Kelleher, C., Cloutier, M., Bakker, J., Enos, J., Arnold, G., Bauer, G., & Morin, P. (2022b). *The Reference Elevation Model of Antarctica - Strips, Version 4.1*. Harvard Dataverse. https://doi.org/10.7910/DVN/X7NDNY

Hoyer, S., & Hamman, J. (2017). xarray: N-d labeled Arrays and Datasets in Python. *Journal of Open Research Software*, *5*(1), 10. https://doi.org/10.5334/jors.148

Morlighem, M. (2020a). *IceBridge BedMachine Greenland, Version 5*. NASA National Snow; Ice Data Center Distributed Active Archive Center. https://doi.org/10.5067/GMEVBWFLWA7X

Morlighem, M. (2020b). *MEaSUREs BedMachine Antarctica, Version 3*. NASA National Snow; Ice Data Center Distributed Active Archive Center. https://doi.org/10.5067/FPSU0V1MWUB6

Noh, M.-J., & Howat, I. M. (2017). The Surface Extraction from TIN based Search-space Minimization (SETSM) algorithm. *ISPRS Journal of Photogrammetry and Remote Sensing*, *129*, 55–76. https://doi.org/10.1016/j.isprsjprs.2017.04.019

Nuth, C., & Kääb, A. (2011). Co-registration and bias corrections of satellite elevation data sets for quantifying glacier thickness change. *The Cryosphere*, *5*(1), 271–290. https://doi.org/10.5194/tc-5-271-2011

Porter, C., Howat, I., Noh, M.-J., Husby, E., Khuvis, S., Danish, E., Tomko, K., Gardiner, J., Negrete, A., Yadav, B., Klassen, J., Kelleher, C., Cloutier, M., Bakker, J., Enos, J., Arnold,

DRAFT

G., Bauer, G., & Morin, P. (2022). *ArcticDEM - Strips, Version 4.1*. Harvard Dataverse. https://doi.org/10.7910/DVN/C98DVS

Porter, C., Howat, I., Noh, M.-J., Husby, E., Khuvis, S., Danish, E., Tomko, K., Gardiner, J., Negrete, A., Yadav, B., Klassen, J., Kelleher, C., Cloutier, M., Bakker, J., Enos, J., Arnold, G., Bauer, G., & Morin, P. (2023). *ArcticDEM - Mosaics, Version 4.1*. Harvard Dataverse. https://doi.org/10.7910/DVN/3VDC4W

Radiant Earth. (2017 - 2024). *PySTAC* (Version 1.10.1). https://github.com/stac-utils/pystac/

Scheick, J., Leong, W. J., Bisson, K., Arendt, A., Bhushan, S., Fair, Z., Hagen, N. R., Henderson, S., Knuth, F., Li, T., Liu, Z., Piunno, R., Ravinder, N., Landung, Sutterley, T., Swinski, J., & Anubhav. (2023). Icepyx: Querying, obtaining, analyzing, and manipulating ICESat-2 datasets. *Journal of Open Source Software*, *8*(84), 4912. https://doi.org/10.21105/joss.04912

Shiggins, C. J., Lea, J. M., & Brough, S. (2023). Automated ArcticDEM iceberg detection tool: Insights into area and volume distributions, and their potential application to satellite imagery and modelling of glacier–iceberg–ocean systems. *The Cryosphere*, *17*(1), 15–32. https://doi.org/10.5194/tc-17-15-2023

Van den Bossche, J., Jordahl, K., Fleischmann, M., Richards, M., McBride, J., Wasserman, J., Badaracco, A. G., Snow, A. D., Ward, B., Tratner, J., Gerard, J., Perry, M., Farmer, C., Hjelle, G. A., Taves, M., Hoeven, E. ter, Cochran, M., Bell, R., rraymondgh, … Kaushik. (2016 - 2024). *Geopandas* (Version 0.14.4). https://doi.org/10.5281/zenodo.11080352

Zevenbergen, L. W., & Thorne, C. R. (1987). Quantitative analysis of land surface topography. *Earth Surface Processes and Landforms*, *12*(1), 47–56. https://doi.org/10.1002/esp.3290120107

DRAFT