

WhereTheWaterFlows.jl: Hydrological flow routing on digital elevation models in Julia

Mauro A. Werder ^{1,2}  and Christophe Ogier ^{1,2}

¹ Laboratory of Hydraulics, Hydrology and Glaciology (VAW), ETH Zurich, Zurich, Switzerland  ² Swiss Federal Institute for Forest, Snow and Landscape Research (WSL), bâtiment ALPOLE, Sion, Switzerland   Corresponding author

DOI: [10.21105/joss.10834](https://doi.org/10.21105/joss.10834)

Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Editor: [Jayaram Hariharan](#)  

Reviewers:

- [@evetion](#)
- [@xkykai](#)

Submitted: 28 May 2026

Published: 10 October 2026

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

Summary

WhereTheWaterFlows.jl (WWF) is a Julia package for computing water flow paths and drainage basins on digital elevation models (DEMs) or other hydropotential surfaces. Given a DEM, the package determines *where the water flows* across the landscape, accumulating upslope area and delineating catchment basins. Besides this classic, deterministic routing on surface topographies, WWF also includes routines to route water in subglacial settings as well as uncertainty quantification tools based on a Monte Carlo approach.

Key outputs include the flow-direction field, the upslope contributing area or discharge, the depression-filled DEM, a labelled catchment-basin map, and the location of all sinks. Plotting functions for result visualisation are included. The package is registered in the Julia General registry, has a full test suite and complete [online documentation](#).

WWF implements the widely-used D8 single-flow-direction algorithm, which routes water from each grid cell to its lowest-elevation neighbour among the eight surrounding cells, together with a breach-type depression-filling algorithm to handle closed depressions that would otherwise trap flow (O'Callaghan & Mark, 1984). The underlying graph traversal follows the O(n) recursive strategy of Braun & Willett (2013), giving linear scaling with the number of grid cells. WWF performs comparably to or better than existing routing software, as shown in our [benchmarks](#).

Statement of need

Hydrological flow routing on DEMs is a fundamental operation in geosciences: it underpins catchment delineation, runoff modelling, subglacial hydrology, and landscape-evolution studies. Software implementations of flow routing date back to O'Callaghan & Mark (1984) and, nowadays, many implementations exist. WWF is the first native Julia package for this task.

WWF is suitable for any domain where DEM-based flow analysis is required. However, the package has been designed with glaciological applications in mind where it has been used in six publications to date, see section [Research impact](#). The glaciological focus required the implementation of glacier specific aspects, such as Shreve potential routing (Shreve, 1972) in the Subglacially submodule, as well as the development of uncertainty quantification via Monte Carlo methods (the Randomly submodule). The latter aspect is needed in glaciology as uncertainties in glacier bed DEMs and Shreve-potential are large and thus warrant quantification.

State of the field

Several mature tools exist for hydrological flow routing on DEMs, examples include:

- **Geomorphometry.jl** is a Julia package providing many geomorphic analysis tools, among them flow routing. (Pronk & Schouten, 2026)
- **GRASS GIS** (command line/C/Python/QGIS), the component `r.watershed` implements several water routing and related algorithms. (GRASS-Team et al., 2026)
- **TauDEM** (command line/ArcGIS) provides command-line tools designed for large-scale catchment analysis. (Tesfa et al., 2011)
- **TopoToolbox** (MATLAB/Python) provides widely used functionalities for topographic analysis including flow routing. (Schwanghart & Scherler, 2014)
- **Whitebox Tools** (Rust/Python) provides an extensive geomorphometry toolkit as a Rust library with Python bindings and unofficial [Julia bindings](#). (Lindsay, 2016)
- **RichDEM** (Python/C++) offers a broad set of flow-routing methods. (Barnes et al., 2014)

To our knowledge, none of the existing routing software packages implement specialised subglacial routing functionality or have uncertainty quantification capabilities.

Software design

The central function is `waterflows(dem)`, which accepts a two-dimensional array of elevation values and returns a named tuple of outputs such as upstream area and catchments. Boundary conditions (open or closed edges, fixed sinks) and other options can be specified via keyword arguments. A novel feature of WWF is the feedback-function (via the `feedback_fn` argument of `waterflows`) which is called on each cell before routing its water downstream, enabling feedbacks to occur. This allows for extensive customisations, as demonstrated in the provided examples (see `examples/` directory) on sediment transport and subglacial melt calculations.

For subglacial routing, the main function is `waterflows_subglacial(surfdem, beddem, dx, f)` with surface and bed DEMs, grid spacing and water pressure as flotation fraction as inputs. In the subglacial setting, the `beddem` and `f` input fields have large uncertainties associated with them. This spurred the development of the Monte Carlo based uncertainty quantification in WWF (which can be used independently of subglacial routing). The main function there is `map_mc(model, sample, reduce!, n)` which runs the `model` function (i.e. the water routing) on `n` samples of surface, bed and flotation fractions fields and reduces the results with `reduce!` (to avoid huge data volumes). The uncertainty in the input fields is modelled using Gaussian Random Fields, which provide spatially correlated random noise. The random fields are generated using a FFT-based method (Räss et al., 2019). However, the software design allows to couple more sophisticated, external geostatistical models to generate the uncertainty fields.

From the outset, WWF has been developed with subglacial applications in mind where uncertainties in inputs are large and would likely dominate over routing algorithm choices. Therefore, the simplest routing algorithm, D8, was deemed sufficient and implemented. Incidentally, the ensemble of flow paths resulting from stochastic routing exhibits an effectively diffusive flow (which is what more sophisticated routing schemes, such as D-inf provide). Although subglacial routing was the motivation to develop WWF, the main WWF module and the `Randomly` submodule were designed, again from the outset, to be usable and useful for general settings. Making WWF useful for general settings also motivated our choice to keep the subglacial and stochastic functionality tucked away in submodules, which makes it clear that these are optional features and that WWF can be used without them. The API of WWF is intentionally kept simple and is based around using generic vectors and arrays as inputs and no custom datatypes are defined (with the exception of the `Uncertainty` struct). This makes WWF interoperable with many grid based Julia packages, in particular georeferenced raster data of the standard packages [Rasters.jl](#) and [GeoArrays.jl](#) can be used and leads to georeferenced WWF outputs (except for `Randomly` workflows which currently return normal arrays).

Example

The code and figure below show an example of a drainage network calculation on a small synthetic DEM:

```
using WhereTheWaterFlows, CairoMakie
n = 200
x = y = range(-π, π, length=n)
dem = sin.(x) .* cos.(y') .+ 0.05 * rand(n, n)

out = waterflows(dem)
plt_area(x, y, out.area)
```




Figure 1: Upslope contributing area on a synthetic DEM from running the provided code example. Brighter colours indicate larger contributing areas, tracing the drainage network emerging from the synthetic topography.

Research impact

WhereTheWaterFlows.jl has been used in published glaciological research: Malczyk et al. (2023) used it to route subglacial melt water and constrain melt fluxes beneath an ice sheet from observations of active subglacial lake recharge. The study employed WWF's uncertainty quantification features. Delaney et al. (2023) built a 2D subglacial sediment transport model on top of WWF. Ogier et al. (2025) employed flow routing to analyse the likelihood of subglacial water pockets in Alpine glaciers, contributing to a new inventory for the Swiss Alps. Horgan et al. (2025) used flow routing to delineate the subglacial catchment of the Kamb Ice Stream. Washam et al. (2026) demonstrated that volcanic heat sources within the Kamb catchment likely contribute to enhanced subglacial melt. Ogier et al. (2026) used the package to assess the potential contribution of subglacial water reservoirs to the catastrophic 2024 La

Bérarde flood.

WWF is used for teaching in the lecture courses “Physics of Glaciers” at ETH Zurich (Switzerland) and “Introduction to geoscientific programming” at Uni Mainz (Germany). The software is archived on Zenodo, has accumulated 20+ GitHub stars and 5+ forks since its public release in 2019. It has four contributors and 200+ commits across 20+ tagged releases.

AI usage disclosure

AI tools from Anthropic and OpenAI are used in the development of WWF. Most of the code predates 2022, the advent of capable LLMs, and was thus written by hand. This also applies to the Randomly and Subglacially submodules which were previously developed in a separate, private repository and only now merged into WWF. Recently, code refactoring was helped by LLMs. Much of the documentation, examples and benchmarks were generated with the help of LLMs. The tools employed to date are Claude Sonnet 4.6 and GPT-5.3-Codex. All AI generated content was reviewed by a human.

The first draft of this manuscript was created by Claude Sonnet 4.6. The tool was used for initial text generation and structural organisation of the paper sections. The text was then reviewed, heavily edited and rewritten by the two authors.

Acknowledgements

Development of WWF was partly funded by the European Space Agency’s project 4DAntarctica (ESA: Grant 4000128611/19/I-DT) and by the Swiss National Science Foundation’s project DIWING (grant nr. 212061).

We thank the reviewers, Xin Kai Lee @xkykai and Maarten Pronk @evetion, and the editor Jayaram Hariharan @elbeejay for their suggestions, inputs and for their time.

References

- Barnes, R., Lehman, C., & Mulla, D. (2014). Priority-flood: An optimal depression-filling and watershed-labeling algorithm for digital elevation models. *Computers & Geosciences*, 62, 117–127. <https://doi.org/10.1016/j.cageo.2013.04.024>
- Braun, J., & Willett, S. D. (2013). A very efficient $O(n)$, implicit and parallel method to solve the stream power equation governing fluvial incision and landscape evolution. *Geomorphology*, 180–181, 170–179. <https://doi.org/10.1016/j.geomorph.2012.10.008>
- Delaney, I., Anderson, L., & Herman, F. (2023). Modeling the spatially distributed nature of subglacial sediment transport and erosion. *Earth Surface Dynamics*, 11(4), 663–680. <https://doi.org/10.5194/esurf-11-663-2023>
- GRASS-Team, Landa, M., Neteler, M., Metz, M., Petrášová, A., Petráš, V., Clements, G., Zigo, T., Larsson, N., Kládíková, L., Haedrich, C., Blumentrath, S., Andreo, V., Cho, H., Gebbert, S., Nartišs, M., Kudrnovsky, H., Delucchi, L., Zambelli, P., ... Bowman, H. (2026). GRASS (Version 8.5.0). Zenodo. <https://doi.org/10.5281/ZENODO.5176030>
- Horgan, H. J., Stewart, C., Stevens, C., Dunbar, G., Balfoort, L., Schmidt, B. E., Washam, P., Werder, M. A., Mandeno, D., Marschalek, J., Hulbe, C., Holschuh, N., Levy, R., Hurwitz, B., Jendersie, S., Johnson, K., Lawrence, J., Morgenstern, R., Mullen, A. D., ... Whiteford, A. (2025). A West Antarctic grounding-zone environment shaped by episodic water flow. *Nature Geoscience*, 1–7. <https://doi.org/10.1038/s41561-025-01687-3>
- Lindsay, J. B. (2016). Whitebox GAT: A case study in geomorphometric analysis. *Computers & Geosciences*, 95, 75–84. <https://doi.org/10.1016/j.cageo.2016.07.003>

- Malczyk, G., Gourmelen, N., Werder, M., Wearing, M., & Goldberg, D. (2023). Constraints on subglacial melt fluxes from observations of active subglacial lake recharge. *Journal of Glaciology*, 1–15. <https://doi.org/10.1017/jog.2023.70>
- O'Callaghan, J. F., & Mark, D. M. (1984). The extraction of drainage networks from digital elevation data. *Computer Vision, Graphics, and Image Processing*, 28(3), 323–344. [https://doi.org/10.1016/S0734-189X\(84\)80011-0](https://doi.org/10.1016/S0734-189X(84)80011-0)
- Ogier, C., Fischer, M., Werder, M. A., Huss, M., Hupfer, M., Jacquemart, M., Gagliardini, O., Gilbert, A., Hösli, L., Thibert, E., Vincent, C., & Farinotti, D. (2025). Definition, formation and rupture mechanisms of water pockets in alpine glaciers: Insights from an updated inventory for the Swiss Alps. *Journal of Glaciology*, 71, e82. <https://doi.org/10.1017/jog.2025.43>
- Ogier, C., Werder, M. A., Gagliardini, O., Santin, I., Moser, R., Hugonnet, R., Blanc, A., & Farinotti, D. (2026). Potential glacier contributions to the 2024 La Bérarde flood. *Natural Hazards and Earth System Sciences*, 26(8), 3901–3917. <https://doi.org/10.5194/nhess-26-3901-2026>
- Pronk, M., & Schouten, R. (2026). Geomorphometry.jl: Analyzing and visualizing the shape of the Earth in Julia. *Journal of Open Source Software*, 11(122), 10395. <https://doi.org/10.21105/joss.10395>
- Räss, L., Kolyukhin, D., & Minakov, A. (2019). Efficient parallel random field generator for large 3-D geophysical problems. *Computers & Geosciences*, 131, 158–169. <https://doi.org/10.1016/j.cageo.2019.06.007>
- Schwanghart, W., & Scherler, D. (2014). TopoToolbox 2 – MATLAB-based software for topographic analysis and modeling in Earth surface sciences. *Earth Surface Dynamics*, 2(1), 1–7. <https://doi.org/10.5194/esurf-2-1-2014>
- Shreve, R. L. (1972). Movement of water in glaciers. *Journal of Glaciology*, 11(62), 205–214. <https://doi.org/10.3189/S002214300002219X>
- Tesfa, T. K., Tarboton, D. G., Watson, D. W., Schreuders, K. A. T., Baker, M. E., & Wallace, R. M. (2011). Extraction of hydrological proximity measures from DEMs using parallel processing. *Environmental Modelling & Software*, 26(12), 1696–1709. <https://doi.org/10.1016/j.envsoft.2011.07.018>
- Washam, P., Schmidt, B. E., Loose, B., Horgan, H. J., Stewart, C. L., Stevens, C. L., Lawrence, J. D., Hulbe, C. L., Mullen, A. D., Quartini, E. S., & Hurwitz, B. (2026). Oceanic and volcanic heat converge in a subglacial channel of the Kamb Ice Stream in West Antarctica. *Communications Earth & Environment*. <https://doi.org/10.1038/s43247-026-03508-w>