The Journal of Open Source Software

divent: An R Package for Diversity Measures Based on Entropy

Eric Marcon [1,2¶] **and Florence Puech** [3]

1 AgroParisTech, Montpellier, France **2** AMAP, CIRAD, CNRS, INRAE, IRD, Univ Montpellier, Montpellier, France **3** Université Paris-Saclay, INRAE, AgroParisTech, Paris-Saclay Applied Economics, F-91120, Palaiseau, France ¶ Corresponding author

DOI: 10.21105/joss.10860

Software

- Review
- Repository
- Archive

Editor: Matthias Grenié

Reviewers:

- @Bisaloo
- @ahasverus

Submitted: 04 May 2026
Published: 09 November 2026

License
Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License (CC BY 4.0).

Summary

`divent` is an R package dedicated to the measurement of diversity in general, and biodiversity in particular. It provides functions to estimate α, β and γ diversity of communities, including phylogenetic and functional diversity.

It is a reboot of the `entropart` R package to make it easier to use and computationally more efficient. It also extends its functionality.

Data may be abundance vectors or matrices (the number of individual per species, possibly per site), lists of individuals, or point patterns to address spatially-explicit diversity. Random communities can be simulated (function `rcommunity()`) and spatialized (function `rspcommunity()`). The main functions allow computing:

- Taxonomic entropy: HCDT entropy (`ent_tsallis()`) including Shannon and Simpson's entropies.
- Phylogenetic (`ent_phylo()`) or functional (`ent_similarity()`) entropy when dissimilarities between species are taken into account in a dendrogram or a similarity matrix.
- Effective numbers of species, aka Hill numbers, for all entropies (`div_hill()`, `div_phylo()`, `div_similarity()`).
- Diversity partitioning (`div_part()`), i.e. α, β and γ diversity of a set of communities.
- Diversity profiles (`profile_hill()`, `profile_phylo()`, `profile_similarity()`), with respect to the order or diversity.
- Diversity accumulation curves (`accum_hill()`) with respect to the sample size or coverage.
- Spatial accumulation of diversity (`accum_sp_hill()`) with respect to the distance from individuals, with maps (`plot_map()`).

Many estimators are proposed, with sensible automatic choice, to deal with incomplete sampling.

Statement of need

Measuring biological diversity (Magurran, 2004) has been a theoretical and empirical challenge for decades. A consensus emerged (Jost, 2006) to derive it from the theory of information (Shannon, 1948). Individuals are sampled and their species is recorded.

Entropy is the expected amount of information brought by an individual. Generalized entropy (Tsallis, 1988) allows choosing the balance between rare and abundant species. Classical indices of biodiversity such as Shannon's and Simpson's indices are just special cases of generalized entropy. Phylogenetic and functional entropy extend the definition taking into account the dissimilarity between species summarized respectively in a dendrogram (Marcon & Hérault,

2015a) or a distance matrix (Leinster & Cobbold, 2012). Finally, Hill numbers (Hill, 1973) are the number of species with equal abundance that would have the same entropy as the data, allowing measuring diversity as a effective number.

Data are generally a sample of a community, requiring estimators of entropy (Marcon, 2015). Estimators of richness (i.e., the number of species), Shannon's entropy or generalized entropy rely on elaborate computing implemented in `divent` (converting estimated entropy into diversity is straightforward). When the sampling effort is not sufficient to allow estimating asymptotic diversity, standardized diversity for a chosen sampling level is necessary and diversity accumulation with respect to the sample size can be estimated (Chao et al., 2014).

Spatially-explicit diversity raises an increasing interest (Wiegand et al., 2017) to address the diversity of the neighborhood of individuals rather than the diversity of an arbitrary plot. The spatial accumulation of diversity with respect to the size of the neighborhood is an efficient tool to understand community assembly rules (Shen et al., 2013; Wiegand et al., 2017). When individual location is not available, the partitioning of the diversity of a set of communities (Whittaker, 1960), called γ diversity, into the average diversity of communities (α diversity) and their divergence (β diversity) is necessary, whatever the diversity considered, for the same purposes.

The `divent` R package enables all these diversity features to be computed.

State of the field

The `entropart` R package (Marcon & Hérault, 2015b) was first published in 2015 and augmented following the literature and the research programs of the authors. It suffers from its incremental development and some obsolete choices such as the organisation of the data, camel-case names of objects, and the insufficient consistence between the behavior of different functions. These issues justified a complete refactoring. Spatially-explicit diversity (Marcon & Puech, 2023; Shimatani, 2001) is completely new in `divent`.

The other major R package with the same purposes is `iNEXT` (Hsieh et al., 2016) and its update `iNEXT.3D` (Chao et al., 2021). The most important features are identical. Additionally, `iNEXT.3D` addresses temporal changes in diversity (`divent` does not yet), while `divent` addresses spatially-explicit diversity. `divent` proposes more atomic functions to calculate various aspects of diversity separately (thus faster, allowing including the results in larger analyses) while `iNEXT.3D` relies on the single, end-user oriented `estimate3D()` function for all purposes.

`divent` is built on some explicit theoretical choices that differ from those of `iNEXT.3D`:

- Diversity is explicitly derived from entropy (Jost, 2006), following the information theory.
- Richness estimation is not limited to Chao's estimators (Marcon, 2015). The jackknife estimator (Burnham & Overton, 1979) is often employed to reduce diversity underestimation owing to undersampling. Far more estimators of entropy in general are available.
- Phylogenies must be ultrametric (Marcon & Hérault, 2015a) and no functional distance cutoff is proposed following Pavoine et al. (2016) rather than Chao et al. (2019) for theoretical arguments.

Software design

`divent` is a written in S3, the object-oriented programming framework provided directly within base R. It follows the state-of-the-art design of R packages (Wickham & Bryan, 2023), namely function naming with prefixes (e.g., `ent_*` functions compute entropy), extensive documentation thanks to packages `roxygen2` (Wickham, Danenberg, et al., 2025) for R objects, `Rdpack` (Boshnakov, 2026) for references and `pkgdown` (Wickham, Hesselberth, et al.,

2025) for online manual. Continuous integration including unit tests (Wickham, 2011) and coverage (Hester, 2025) is hosted by GitHub.

Generic functions (e.g., `div_hill()`) are used to allow two data formats, addressed by their respective method: numeric vectors containing the number of individual per species (function `div_hill.numeric()`) or objects of class `species_distributions` (function `div_hill.species_distribution()`), which are `data.frames` whose rows are communities, columns are species and values are abundances (objects of class `abundances`, that inherits from `species_distributions`) or probabilities (class `probabilities`). Functions to convert a classical abundance matrix, a list of individuals or a point pattern to an `abundances` object are provided.

Spatially-explicit diversity measures require counting the number of neighbors around each individual. This task is run by a short C++ code integrated and parallelized by the `RcppParallel` package (Allaire et al., 2023).

Research impact statement

`divent` is the successor of the `entropart` R package (Marcon & Hérault, 2015b) that is widely used (around 40 citations every year according to Google Scholar). It is often employed in tropical forest ecology (Poorter et al., 2021; Réjou-Méchain et al., 2021) where correctly estimating diversity from incomplete sampling is critical, but also in other disciplines such as evolutionary (Hafer-Hahmann & Vorburger, 2020), microbial (Kouakou et al., 2025) or marine ecology (Huang et al., 2024), and even agronomy (Mandal et al., 2018) or environmental sociology (Dago et al., 2025).

`divent` was first adopted by Prunot et al. (2026) for estimating the diversity of Carabidae. Furthermore, the `MiscMetabar` R package (Taudière, 2023) has relied on `divent` for its diversity calculations since version 0.15.1.

AI usage disclosure

No AI tools were used at any stage of the package development nor in this article.

Acknowledgements

This work benefited from the support of "Investissements d'avenir" of the French National Agency for Research (Labex CEBA, ref. ANR-10-LABX-25-01). The authors thank Bruno Hérault for his contributions to the entropart package.

References

Allaire, J., Francois, R., Ushey, K., Vandenbrouck, G., Geelnard, M., & Intel. (2023). *RcppParallel: Parallel programming tools for 'Rcpp'* [Manual]. https://doi.org/10.32614/CRAN.package.RcppParallel

Boshnakov, G. N. (2026). *Rdpack: Update and manipulate rd documentation objects* [Manual]. https://doi.org/10.32614/CRAN.package.Rdpack

Burnham, K. P., & Overton, W. S. (1979). Robust estimation of population size when capture probabilities vary among animals. *Ecology*, *60*(5), 927–936. https://doi.org/10.2307/1936861

Chao, A., Chiu, C.-H., Villéger, S., Sun, I.-F., Thorn, S., Lin, Y.-C., Chiang, J.-M., & Sherwin, W. B. (2019). An attribute-diversity approach to functional diversity, functional beta

diversity, and related (dis)similarity measures. *Ecological Monographs*, *89*(2). https://doi.org/10.1002/ecm.1343

Chao, A., Gotelli, N. J., Hsieh, T. C., Sander, E. L., Ma, K. H., Colwell, R. K., & Ellison, A. M. (2014). Rarefaction and extrapolation with Hill numbers: A framework for sampling and estimation in species diversity studies. *Ecological Monographs*, *84*(1), 45–67. https://doi.org/10.1890/13-0133.1

Chao, A., Henderson, P. A., Chiu, C.-H., Moyes, F., Hu, K.-H., Dornelas, M., & Magurran, A. E. (2021). Measuring temporal change in alpha diversity: A framework integrating taxonomic, phylogenetic and functional diversity and the iNEXT.3D standardization. *Methods in Ecology and Evolution*, *12*(10), 1926–1940. https://doi.org/10.1111/2041-210X.13682

Dago, M. R., Zo-Bi, I. C., Konan, I. K., Kouassi, A. K., Guei, S., Jagoret, P., N'Guessan, A. E., Sanial, E., Tankam, C., Traoré, S., & Hérault, B. (2025). What motivates West African cocoa farmers to value trees? Taking the 4 W approach to the heart of the field. *People and Nature*, *7*(1), 215–230. https://doi.org/10.1002/pan3.10754

Hafer-Hahmann, N., & Vorburger, C. (2020). Parasitoids as drivers of symbiont diversity in an insect host. *Ecology Letters*, *23*(8), 1232–1241. https://doi.org/10.1111/ele.13526

Hester, J. (2025). *Covr: Test coverage for packages* [Manual]. https://doi.org/10.32614/CRAN.package.covr

Hill, M. O. (1973). Diversity and evenness: A unifying notation and its consequences. *Ecology*, *54*(2), 427–432. https://doi.org/10.2307/1934352

Hsieh, T. C., Ma, K. H., & Chao, A. (2016). iNEXT: An R package for interpolation and extrapolation in measuring species diversity. *Methods in Ecology and Evolution*, *7*, 1451–1456. https://doi.org/10.1111/2041-210X.12613

Huang, M., Chen, Y., Zhou, W., & Wei, F. (2024). Assessing the response of marine fish communities to climate change and fishing. *Conservation Biology*, *38*(6), e14291. https://doi.org/10.1111/cobi.14291

Jost, L. (2006). Entropy and diversity. *Oikos*, *113*(2), 363–375. https://doi.org/10.1111/j.2006.0030-1299.14714.x

Kouakou, A. K., Collart, P., Perron, T., Kolo, Y., Gay, F., Brauman, A., & Brunel, C. (2025). Soil Microbial Recovery to the Rubber Tree Replanting Process in Ivory Coast. *Microbial Ecology*, *88*(1), 13. https://doi.org/10.1007/s00248-025-02506-3

Leinster, T., & Cobbold, C. (2012). Measuring diversity: The importance of species similarity. *Ecology*, *93*(3), 477–489. https://doi.org/10.1890/10-2402.1

Magurran, A. E. (2004). *Measuring biological diversity*. Blackwell Publishing. ISBN: 978-1-118-68792-5

Mandal, S., Donner, E., Vasileiadis, S., Skinner, W., Smith, E., & Lombi, E. (2018). The effect of biochar feedstock, pyrolysis temperature, and application rate on the reduction of ammonia volatilisation from biochar-amended soil. *Science of The Total Environment*, *627*, 942–950. https://doi.org/10.1016/j.scitotenv.2018.01.312

Marcon, E. (2015). Practical estimation of diversity from abundance data. *HAL*, *01212435*(version 2).

Marcon, E., & Hérault, B. (2015a). Decomposing phylodiversity. *Methods in Ecology and Evolution*, *6*(3), 333–339. https://doi.org/10.1111/2041-210X.12323

Marcon, E., & Hérault, B. (2015b). Entropart, an R package to measure and partition diversity. *Journal of Statistical Software*, *67*(8), 1–26. https://doi.org/10.18637/jss.v067.i08

Marcon, E., & Puech, F. (2023). Mapping distributions in non-homogeneous space with

distance-based methods. *Journal of Spatial Econometrics*, *4*(1), 13. https://doi.org/10.1007/s43071-023-00042-1

Pavoine, S., Marcon, E., & Ricotta, C. (2016). "Equivalent numbers" for species, phylogenetic or functional diversity in a nested hierarchy of multiple scales. *Methods in Ecology and Evolution*, *7*(10), 1152–1163. https://doi.org/10.1111/2041-210X.12591

Poorter, L., Craven, D., Jakovac, C. C., van der Sande, M. T., Amissah, L., Bongers, F., Chazdon, R. L., Farrior, C. E., Kambach, S., Meave, J. A., Muñoz, R., Norden, N., Rüger, N., van Breugel, M., Almeyda Zambrano, A. M., Amani, B., Andrade, J. L., Brancalion, P. H. S., Broadbent, E. N., … Hérault, B. (2021). Multidimensional tropical forest recovery. *Science*, *374*(6573), 1370–1376. https://doi.org/10.1126/science.abh3629

Prunot, S., Cortet, J., Auclerc, A., Henon, N., Jolivet, C., Magne, G., Peres, G., Pouzenc, S., Versavel, C., Vincent, Q., & Hedde, M. (2026). *Imperfect detection and sampling effort shape ground beetle (Carabidae) diversity indicators in large-scale monitoring*. SSRN. https://doi.org/10.2139/ssrn.6175011

Réjou-Méchain, M., Mortier, F., Bastin, J.-F., Cornu, G., Barbier, N., Bayol, N., Bénédet, F., Bry, X., Dauby, G., Deblauwe, V., Doucet, J.-L., Doumenge, C., Fayolle, A., Garcia, C., Kibambe Lubamba, J.-P., Loumeto, J.-J., Ngomanda, A., Ploton, P., Sonké, B., … Gourlet-Fleury, S. (2021). Unveiling African rainforest composition and vulnerability to global change. *Nature*. https://doi.org/10.1038/s41586-021-03483-6

Shannon, C. E. (1948). A mathematical theory of communication. *The Bell System Technical Journal*, *27*(3), 379–423. https://doi.org/10.1002/j.1538-7305.1948.tb01338.x

Shen, G., He, F., Waagepetersen, R., Sun, I.-F., Hao, Z., Chen, Z.-S., & Yu, M. (2013). Quantifying effects of habitat heterogeneity and other clustering processes on spatial distributions of tree species. *Ecology*, *94*(11), 2436–2443. https://doi.org/10.1890/12-1983.1

Shimatani, K. (2001). Multivariate point processes and spatial variation of species diversity. *Forest Ecology and Management*, *142*(1-3), 215–229. https://doi.org/10.1016/s0378-1127(00)00352-2

Taudière, A. (2023). MiscMetabar: An R package to facilitate visualizationand reproducibility in metabarcoding analysis. *Journal of Open Source Software*, *8*(92), 6038. https://doi.org/10.21105/joss.06038

Tsallis, C. (1988). Possible generalization of boltzmann-gibbs statistics. *Journal of Statistical Physics*, *52*(1), 479–487. https://doi.org/10.1007/BF01016429

Whittaker, R. H. (1960). Vegetation of the Siskiyou mountains, Oregon and California. *Ecological Monographs*, *30*(3), 279–338. https://doi.org/10.2307/1943563

Wickham, H. (2011). Testthat: Get started with testing. *The R Journal*, *3*, 5–10. https://doi.org/10.32614/RJ-2011-002

Wickham, H., & Bryan, J. (2023). *R packages* (2nd ed.). O'Reilly Media, Inc. ISBN: 978-1-0981-3494-5

Wickham, H., Danenberg, P., Csárdi, G., & Eugster, M. (2025). *Roxygen2: In-Line Documentation for R* [Manual]. https://doi.org/10.32614/CRAN.package.roxygen2

Wickham, H., Hesselberth, J., Salmon, M., Roy, O., & Brüggemann, S. (2025). *Pkgdown: Make static HTML documentation for a package* [Manual]. https://doi.org/10.32614/CRAN.package.pkgdown

Wiegand, T., Uriarte, M., Kraft, N. J. B., Shen, G., Wang, X., & He, F. (2017). Spatially explicit metrics of species diversity, functional diversity, and phylogenetic diversity: Insights into plant community assembly processes. *Annual Review of Ecology, Evolution, and*

Systematics, *48*(1), 329–351. https://doi.org/10.1146/annurev-ecolsys-110316-022936