

Magrathea v2: A planetary interior modeling platform in C++

David R. Rice^{1,5}[¶], Chenliang Huang²[¶], Robert Royer³[¶], Mangesh Daspute⁴, Krishang Mittal⁵, Baptiste Journaux⁶[¶], Jason H. Steffen³[¶], and Allona Vazan¹[¶]

¹ Astrophysics Research Center (ARCO), Department of Natural Sciences, The Open University of Israel, Raanana 4353701, Israel ² Shanghai Astronomical Observatory, Chinese Academy of Sciences, Shanghai 200030, People's Republic of China ³ Department of Physics and Astronomy, University of Nevada, Las Vegas, 4505 South Maryland Parkway, Las Vegas, NV 89154, USA ⁴ Department of Physics, Ariel University, Ariel 40700, Israel ⁵ Department of Astronomy, University of Wisconsin–Madison, 475 N. Charter Street, Madison, WI 53706, USA ⁶ Department of Earth and Space Science, University of Washington, Seattle, WA 98195, USA [¶] Corresponding author

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Editor: [Tobias Buck](#)  

Reviewers:

- [@Mark-Eb](#)
- [@astro-seanwhy](#)

Submitted: 23 October 2025

Published: unpublished

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

Summary

Magrathea is an open-source C++ code for modeling the internal structure of differentiated planets. The initial release, Huang et al. (2022), introduced the base solver, a modular framework for defining equations of state (EOS) within phase diagrams for each layer, and a plan for expanding the code. Many of those updates are now implemented. Magrathea v2 is a more versatile platform that supports a wider range of compositions, adds new tools for composition retrieval, and makes it easier for users to adapt the code to their own models.

Statement of need

Constraining a planet's composition is essential for understanding its formation and evolution. Observations of mass and radius alone are not sufficient, since many different interiors can yield the same bulk density. Interior structure solvers are therefore essential tools for constraining possible compositions. With observational programs routinely measuring the densities of small to large planets, researchers require codes with models that are transparent and flexible which can adapt as our understanding of planet mineralogy changes.

Magrathea is designed as such a platform. Rather than enforcing a fixed planet model, Magrathea provides a framework in which users can define their own phase diagrams, equations of state (EOS), and thermal profiles. The goal is not one preferred interior model, but a fast and readable code base for building and testing many of them. With the continued expansion of the physics and usability in Version 2.0, Magrathea helps the community keep up with the growing precision of exoplanet observations and experimental constraints on planetary materials.

State of the field

The field now includes several open tools for planet interior structure modeling, with broader summaries and comparisons given in Acuña et al. (2025) and Baumeister et al. (2025). Two open-source options are GASTLI (Acuña et al., 2025) which targets volatile-rich planets with coupled interior–atmosphere modeling and ExoPlex (Unterborn et al., 2023) which focuses on rocky-planet mantle mineralogy.

Magrathea gives users explicit control over phase diagrams and EOS choices in each differentiated layer allowing for diverse planet models from sub-Earth to Neptune-mass planets. Magrathea allows users to swap mineral physics assumptions quickly, run forward models fast enough for large sweeps, and test how interior assumptions propagate into inferred compositions. The combination of modularity and speed from the C++ implementation is the main reason we seek to build on Magrathea.

Software design

The core solver of Magrathea is a one-dimensional, spherically symmetric integrator of the equations of hydrostatic equilibrium, mass continuity, temperature gradient, and equation of state. For a user-defined planet consisting of up to four differentiated layers, the code integrates inward and outward solutions using a shooting-to-fitting-point method with adaptive Runge–Kutta–Fehlberg stepping. The solver returns the radius of the planet, the radii of each compositional boundary, and profiles of pressure, temperature, density, and phase as functions of enclosed mass. Solving one planet takes about one second for most configurations.

A key design choice is modularity. A large variety of EOS forms are supported in `EOS.cpp`, including Birch–Murnaghan, Vinet, Holzapfel, Keane, ideal and van der Waals gases, Debye or Einstein thermal terms, and tabulated EOS. Parameters for each material are defined in a library of more than 70 EOS in `EOSlist.cpp`. Phase diagrams for each layer define which material is used at a given pressure–temperature condition in `phase.cpp`. Alternative phase diagrams are also stored in a library and can be selected at run time.

Magrathea separates the solver from the model. The code offers nine run modes through human-readable `.cfg` files, including a full four-layer solver, isothermal and two-layer modes, bulk runs, two/three-layer composition finder, on-the-fly EOS modification modes, and an MCMC retrieval mode. This range of models and use cases makes Magrathea not just a solver but a platform for exploring interior models.

Major updates in this version

Since the initial release (Huang et al., 2022), Magrathea has undergone expansions in physics, solvers, and usability.

New physical models and materials

- **Default Mantle:** Added upper-mantle polymorphs of Mg_2SiO_4 (forsterite, wadsleyite, ringwoodite) (Dorogokupets et al., 2015), see Figure 1.
- **Default Hydrosphere:** Updated H_2O EOS and phase boundaries for ices (Journaux et al., 2020), liquid and gas (Wagner & Pruß, 2002), and supercritical water (Mazevet et al., 2019), largely inspired by the AQUA package (Haldemann et al., 2020), see Figure 1.
- **Additional Gas EOS:** Including the solar-metallicity hydrogen/helium table from Chabrier & Debras (2021) and van der Waals gases.
- **Carbon Mantles:** EOS and phase diagrams for phases of carbon (Benedict et al., 2014; Lowitzer et al., 2006) and silicon carbide (Miozzi et al., 2018), see Figure 1.
- **EOS library growth:** Including the AQUA table (Haldemann et al., 2020), fcc- and bcc-iron (Dorogokupets et al., 2017), and mantle materials from Stixrude & Lithgow-Bertelloni (2011).

New functionality and solvers

- **Composition finders:**
 - A secant-method routine that determines the mass of a third unknown layer given a target mass, radius, and ratio between the other two layers, looped over layer

ratios and mass-radius posterior draws.

– A Markov chain Monte Carlo routine following Rogers & Seager (2010) and Dorn et al. (2015) for probabilistic composition inference given mass, radius, and associated uncertainties with the Metropolis–Hastings method.

- **Tabulated EOS:** Support for tabulated P - T - ρ - ∇T_S EOS tables using bilinear interpolation.
- **Modular phase diagrams:** Users can store multiple phase-diagram configurations and call them in the configuration file, for example switching between silicate and carbon mantle models without recompiling.

Usability

- **Input handling:** All input parameters were moved to `run/*.cfg` files with descriptive keys.
- **Parallelization:** Bulk runs and composition finder routines can exploit OpenMP in `compfind.cpp` enabling execution with multiple threads.
- **Diagnostics:** More informative error messages are returned when solutions fail to converge.
- **Tutorial and documentation:** A guided set of examples and practice problems resides in the `docs/` folder with online documentation at magrathea.readthedocs.io

Together, these changes make Magrathea v2 a substantially expanded platform rather than a simple update to the default planet model.




Figure 1: New phase diagrams in the code. Left, updated default hydrosphere. Center, default mantle with lower-pressure Mg_2SiO_4 phases. Right, carbon and SiC mantle phase diagrams. On each plot are shown the pressure-temperature conditions inside a one Earth-mass planet with different outer temperatures.

Research impact statement

Magrathea has been used in published work to generate mass-radius diagrams (Taylor et al., 2025), infer interiors of observed planets (Desai et al., 2024; MacDonald et al., 2022), connect theoretical composition from formation to observables (Childs et al., 2023; Dou et al., 2024; Steffen et al., 2025), and test how updated high-pressure EOS measurements and interior assumptions change inferred structures (Huang et al., 2021; Lozovsky, 2026). The new composition finders are used and described in Daspute et al. (2025), Rice et al. (2025), and Kroft et al. (2026).

Beyond its use in published research, Magrathea is already contributing to emerging intercomparison efforts in exoplanet interior modeling. Schulze et al. (2026) compares EOS

and material choices across common rocky-planet models and shows that those choices can change inferred compositions at a level comparable to current observational uncertainties. In that context, open and modular tools like Magrathea help make differences in physical assumptions, EOS choices, and retrieval workflows easier to isolate and test.

Magrathea now covers a broader set of interior assumptions, supports faster composition inference workflows, and makes EOS-level sensitivity studies easier to perform. These are the main research impacts of this version.

AI usage disclosure

Generative AI was not used to implement most of the updates described in this paper. Limited use of Generative AI was explored during development of the MCMC mode. All AI-assisted code changes were reviewed by the authors against the existing code base, and outputs were tested against other non-AI retrieval modes. Generative AI was also used in drafting and reorganizing this manuscript. All scientific claims and software descriptions were checked and revised by the authors against prior publications and the broader literature.

Acknowledgements

We acknowledge the many researchers using the Magrathea code base; we appreciate each contribution to science and the code. We thank Douglas Adams for the timeless stories compiled in “The Hitchhiker’s Guide to the Galaxy” which inspired the name of the code. We acknowledge support from the College of Sciences and the Nevada Center for Astrophysics at the University of Nevada, Las Vegas. A.V. acknowledges support by ISF grants 770/21 and 773/21. C.H. is sponsored by Shanghai Pujiang Program (grant No. 23PJ1414900).

References

- Acuña, L., Kreidberg, L., Zhai, M., Mollière, P., & Fournes, M. (2025). GASTLI: A Python package for coupled interior–atmosphere modelling of volatile-rich planets. *The Journal of Open Source Software*, 10(107), 7288. <https://doi.org/10.21105/joss.07288>
- Baumeister, P., Miozzi, F., Guimond, C. M., Steinmeyer, M.-L., Dorn, C., Karato, S.-I., Bolmont, É., Revol, A., Thamm, A., & Noack, L. (2025). Fundamentals of Interior Modelling and Challenges in the Interpretation of Observed Rocky Exoplanets. *Space Science Reviews*, 221(8), 123. <https://doi.org/10.1007/s11214-025-01248-5>
- Benedict, L. X., Driver, K. P., Hamel, S., Militzer, B., Qi, T., Correa, A. A., Saul, A., & Schwegler, E. (2014). Multiphase equation of state for carbon addressing high pressures and temperatures. *Physical Review B*, 89(22), 224109. <https://doi.org/10.1103/PhysRevB.89.224109>
- Chabrier, G., & Debras, F. (2021). A New Equation of State for Dense Hydrogen-Helium Mixtures. II. Taking into Account Hydrogen-Helium Interactions. *The Astrophysical Journal*, 917(1), 4. <https://doi.org/10.3847/1538-4357/abfc48>
- Childs, A. C., Shakespeare, C., Rice, D. R., Yang, C.-C., & Steffen, J. H. (2023). Composition constraints of the TRAPPIST-1 planets from their formation. *Monthly Notices of the Royal Astronomical Society*, 524(3), 3749–3768. <https://doi.org/10.1093/mnras/stad2110>
- Daspute, M., Wandel, A., Kopparapu, R. K., Perdelwitz, V., Teklu, J. T., & Tal-Or, L. (2025). Potential Interior Structures and Habitability of Super-Earth Exoplanets LHS 1140 b, K2-18 b, TOI-1452 b, and TOI-1468 c. *The Astrophysical Journal*, 979(2), 158. <https://doi.org/10.3847/1538-4357/ad9ba9>

- Desai, A., Turtelboom, E. V., Harada, C. K., Dressing, C. D., Rice, D. R., Murphy, J. M. A., Brinkman, C. L., Chontos, A., Crossfield, I. J. M., Dai, F., Hill, M. L., Fetherolf, T., Giacalone, S., Howard, A. W., Huber, D., Isaacson, H., Kane, S. R., Lubin, J., MacDougall, M. G., ... Winn, J. N. (2024). The TESS-Keck Survey. XVIII. A Sub-Neptune and Spurious Long-period Signal in the TOI-1751 System. *The Astronomical Journal*, 167(5), 194. <https://doi.org/10.3847/1538-3881/ad29ee>
- Dorn, C., Khan, A., Heng, K., Connolly, J. A. D., Alibert, Y., Benz, W., & Tackley, P. (2015). Can we constrain the interior structure of rocky exoplanets from mass and radius measurements? *Astronomy & Astrophysics*, 577, A83. <https://doi.org/10.1051/0004-6361/201424915>
- Dorogokupets, P. I., Dymshits, A. M., Litasov, K. D., & Sokolova, T. S. (2017). Thermodynamics and Equations of State of Iron to 350 GPa and 6000 K. *Scientific Reports*, 7, 41863. <https://doi.org/10.1038/srep41863>
- Dorogokupets, P. I., Dymshits, A. M., Sokolova, T. S., Danilov, B. S., & Litasov, K. D. (2015). The equations of state of forsterite, wadsleyite, ringwoodite, akimotoite, MgSiO₃-perovskite, and postperovskite and phase diagram for the Mg₂SiO₄ system at pressures of up to 130 GPa. *Russian Geology and Geophysics*, 56(1-2), 172–189. <https://doi.org/10.1016/j.rgg.2015.01.011>
- Dou, J., Carter, P. J., & Leinhardt, Z. M. (2024). Formation of super-Mercuries via giant impacts. *Monthly Notices of the Royal Astronomical Society*, 529(3), 2577–2594. <https://doi.org/10.1093/mnras/stae644>
- Haldemann, J., Alibert, Y., Mordasini, C., & Benz, W. (2020). AQUA: a collection of H₂O equations of state for planetary models. *Astronomy & Astrophysics*, 643, A105. <https://doi.org/10.1051/0004-6361/202038367>
- Huang, C., Rice, D. R., Grande, Z. M., Smith, D., Smith, J. S., Boisvert, J. H., Tschauer, O., Salamat, A., & Steffen, J. H. (2021). Implications of an improved water equation of state for water-rich planets. *Monthly Notices of the Royal Astronomical Society*, 503(2), 2825–2832. <https://doi.org/10.1093/mnras/stab645>
- Huang, C., Rice, D. R., & Steffen, J. H. (2022). MAGRATHEA: an open-source spherical symmetric planet interior structure code. *Monthly Notices of the Royal Astronomical Society*, 513(4), 5256–5269. <https://doi.org/10.1093/mnras/stac1133>
- Journaux, B., Brown, J. M., Pakhomova, A., Collings, I. E., Petitgirard, S., Espinoza, P., Boffa Ballaran, T., Vance, S. D., Ott, J., Cova, F., Garbarino, G., & Hanfland, M. (2020). Holistic Approach for Studying Planetary Hydrospheres: Gibbs Representation of Ices Thermodynamics, Elasticity, and the Water Phase Diagram to 2,300 MPa. *Journal of Geophysical Research (Planets)*, 125(1), e06176. <https://doi.org/10.1029/2019JE006176>
- Kroft, M. A., Beatty, T. G., Salzer, J. M., Zwicker, C., Triantafyllides, A., Becker, J., Soares-Furtado, M., Cisewski-Kehe, J., Lissauer, J. J., Armitage, T. S., Livesey, J. R., Narayan, R. S., Widicus Weaver, S., Zhang, K., Bieryla, A., Ciardi, D. R., Clark, C. A., Felsmann, M., Fernandes, R. B., ... Lund, M. B. (2026). GJ 523b is a Massive, 170 Myr-old Mega-Earth, Likely on a Polar Orbit. *arXiv e-Prints*, arXiv:2603.24682. <https://doi.org/10.48550/arXiv.2603.24682>
- Lowitzer, S., Winkler, B., & Tucker, M. (2006). Thermoelastic behavior of graphite from in situ high-pressure high-temperature neutron diffraction. *Physical Review B*, 73(21), 214115. <https://doi.org/10.1103/PhysRevB.73.214115>
- Lozovsky, M. (2026). The Uncertainty in Water Mass Fraction of Wet Planets. *The Astrophysical Journal*, 997(2), 234. <https://doi.org/10.3847/1538-4357/ae2c56>
- MacDonald, M. G., Feil, L., Quinn, T., & Rice, D. (2022). Confirming the 3:2 Resonance Chain of K2-138. *The Astronomical Journal*, 163(4), 162. <https://doi.org/10.3847/1538->

- 207 [3881/ac524c](#)
- 208 Mazevet, S., Licari, A., Chabrier, G., & Potekhin, A. Y. (2019). Ab initio based equation of
209 state of dense water for planetary and exoplanetary modeling. *Astronomy & Astrophysics*,
210 621, A128. <https://doi.org/10.1051/0004-6361/201833963>
- 211 Miozzi, F., Morard, G., Antonangeli, D., Clark, A. N., Mezouar, M., Dorn, C., Rozel, A., &
212 Fiquet, G. (2018). Equation of State of SiC at Extreme Conditions: New Insight Into the
213 Interior of Carbon-Rich Exoplanets. *Journal of Geophysical Research (Planets)*, 123(9),
214 2295–2309. <https://doi.org/10.1029/2018JE005582>
- 215 Rice, D. R., Huang, C., Steffen, J. H., & Vazan, A. (2025). Uncertainties in the Inference
216 of Internal Structure: The Case of TRAPPIST-1 f. *The Astrophysical Journal*, 986(1), 2.
217 <https://doi.org/10.3847/1538-4357/add34b>
- 218 Rogers, L. A., & Seager, S. (2010). A Framework for Quantifying the Degeneracies of
219 Exoplanet Interior Compositions. *The Astrophysical Journal*, 712(2), 974–991. <https://doi.org/10.1088/0004-637X/712/2/974>
- 221 Schulze, J. G., Hinkel, N. R., Panero, W. R., & Unterborn, C. T. (2026). Toward Reliable
222 Interpretations of Small-exoplanet Compositions: Comparisons and Considerations of
223 Equations of State and Materials Used in Common Rocky Planet Models. *The Planetary
224 Science Journal*, 7(2), 30. <https://doi.org/10.3847/PSJ/ae2ea2>
- 225 Steffen, J. H., Shakespeare, C., Royer, R., Rice, D., & Vazan, A. (2025). Effect of Galactic
226 Chemical Evolution on Exoplanet Properties. *The Astrophysical Journal Letters*, 991(2),
227 L33. <https://doi.org/10.3847/2041-8213/ae0457>
- 228 Stixrude, L., & Lithgow-Bertelloni, C. (2011). Thermodynamics of mantle minerals - II. Phase
229 equilibria. *Geophysical Journal International*, 184(3), 1180–1213. <https://doi.org/10.1111/j.1365-246X.2010.04890.x>
- 231 Taylor, J., Radica, M., Chatterjee, R. D., Hammond, M., Meier, T., Aigrain, S., MacDonald,
232 R. J., Albert, L., Benneke, B., Coulombe, L.-P., Cowan, N. B., Dang, L., Doyon, R., Flagg,
233 L., Johnstone, D., Kaltenegger, L., Lafrenière, D., Pelletier, S., Piaulet-Ghorayeb, C., ...
234 Roy, P.-A. (2025). JWST NIRISS transmission spectroscopy of the super-Earth GJ 357b, a
235 favourable target for atmospheric retention. *Monthly Notices of the Royal Astronomical
236 Society*, 540(4), 3677–3692. <https://doi.org/10.1093/mnras/staf894>
- 237 Unterborn, C. T., Desch, S. J., Haldemann, J., Lorenzo, A., Schulze, J. G., Hinkel, N.
238 R., & Panero, W. R. (2023). The Nominal Ranges of Rocky Planet Masses, Radii,
239 Surface Gravities, and Bulk Densities. *The Astrophysical Journal*, 944(1), 42. <https://doi.org/10.3847/1538-4357/aca3b>
- 241 Wagner, W., & Pruß, A. (2002). The IAPWS Formulation 1995 for the Thermodynamic
242 Properties of Ordinary Water Substance for General and Scientific Use. *Journal of Physical
243 and Chemical Reference Data*, 31(2), 387–535. <https://doi.org/10.1063/1.1461829>