The Journal of Open Source Software

Low scaling BSE implementation in the exciting code

Benedikt Maurer [1*] **and Claudia Draxl** [1*]

1 Department of Physics and CSMB Adlershof, Humboldt-Universit"at zu Berlin, Zum Großen Windkanal 2, D-12489 Berlin, Germany * These authors contributed equally.

DOI: 10.xxxxxx/draft

Software

- Review
- Repository
- Archive

Editor: Bonan Zhu

Reviewers:

- @jjkas
- @ruiyiQM
- @EderGio

Submitted: 03 February 2025
Published: unpublished

License
Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License (CC BY 4.0).

Summary

Solving the Bethe-Salpeter Equation (BSE) is essential for understanding excited-state systems, but often challenging to converge or even computationally prohibitive for large systems. We implement a matrix-free BSE solver leveraging Interpolative Separable Density Fitting (ISDF) to interpolate electron-hole interaction kernels together with the Lanczos algorithm for diagonalization, avoiding full matrix setup. The scaling of our implementation is bounded by $\mathcal{O}(N_o N_u N_\mathbf{k} \log N_\mathbf{k})$ and is thus a massive improvement over methods that set up the whole matrix, scaling with at least $\mathcal{O}((N_o N_u N_\mathbf{k})^3)$, where N_o and N_u are the numbers of occupied and unoccupied states, respectively, and $N_\mathbf{k}$ is the number $\mathbf{k}$-points.

Theoretical background

The Bethe-Salpeter Equation (BSE) within many body perturbation theory (MBPT) provides the state-of-the-art frame work for describing light-matter interaction. In particular, it is used to obtain optical absorption spectra, including the effects of excitons, which are bound electron-hole states. By expanding the electron-hole wavefunctions in the transition basis, solving the BSE can be reduced to a Schrödinger like equation. Setting up and diagonalizing the Bethe-Salpeter Hamiltonian (BSH) are the computationally expensive tasks(Vorwerk et al., 2019). The BSH is given as

$$H^{BSH} = D + \gamma V - W\,,$$

where $\gamma = 2$ gives the spin-singlet and $\gamma = 0$ spin-triplet channel, respectively. The diagonal term D is given by the differences of the one-particle energies of the occupied (o) and unoccupied (u) states:

$$D_{ou\mathbf{k},o'u'\mathbf{k}'} = (\varepsilon_{u\mathbf{k}} - \varepsilon_{o\mathbf{k}})\delta_{oo'}\delta_{uu'}\delta_{\mathbf{k}\mathbf{k}'}\,.$$

The matrix elements of the repulsive exchange interaction V and the attractive screened Coulomb interaction W are calculated by solving integrals of the form

$$V_{ou\mathbf{k},o'u'\mathbf{k}'} = \int d^3r \int d^3r' \frac{u_{o\mathbf{k}}(\mathbf{r}) u^*_{u\mathbf{k}}(\mathbf{r}) u^*_{o'\mathbf{k}'}(\mathbf{r}') u_{u'\mathbf{k}'}(\mathbf{r}')}{|\mathbf{r}-\mathbf{r}'|}\,,$$

$$W_{ou\mathbf{k},o'u'\mathbf{k}'} = \int d^3r \int d^3r' u^*_{u\mathbf{k}}(\mathbf{r}) u_{u'\mathbf{k}'}(\mathbf{r}) W_{\mathbf{k}-\mathbf{k}'}(\mathbf{r},\mathbf{r}') u_{o\mathbf{k}}(\mathbf{r}') u^*_{o'\mathbf{k}'}(\mathbf{r}')\,,$$

where $W_{\mathbf{k}-\mathbf{k}'}(\mathbf{r},\mathbf{r}')$ is the statically screened Coulomb potential. $u_{i\mathbf{k}}(\mathbf{r})$ is the periodic part of the one-particle wavefunction of state i at the reciprocal lattice point $\mathbf{k}$. In general, we cannot see, which of the matrix elements will be zero, thus we need to compute all of them. Therefore, setting up the full BSH scales with $\mathcal{O}(N_o^2 N_u^2 N_\mathbf{k}^2)$ and diagonalizing it directly with $\mathcal{O}(N_o^3 N_u^3 N_\mathbf{k}^3)$.

DRAFT

The primary scaling bottleneck stems from evaluating the matrix elements of the interaction kernels. To mitigate this, we reformulate the wavefunction products using Interpolative Separable Density Fitting (ISDF) Lu & Ying (2016). Specifically, we approximate these products on a discrete real-space grid, $\{\mathbf{r}\}$, by expressing them as superpositions of values evaluated on a smaller interpolation grid, $\{\mathbf{r}_\mu\} \subset \{\mathbf{r}\}$:

$$u^*_{i\mathbf{k}}(\mathbf{r})u_{j\mathbf{k}'}(\mathbf{r}) \approx \sum_{\mu=1}^{N_\mu} \zeta_\mu(\mathbf{r})u^*_{i\mathbf{k}}(\mathbf{r}_\mu)u_{j\mathbf{k}}(\mathbf{r}_\mu) \ ,$$

where N_μ is the number of interpolation points, and $\zeta_\mu(\mathbf{r})$ are the expansion coefficients. Due to the tensor product structure of $u^*_{i\mathbf{k}}(\mathbf{r})u_{j\mathbf{k}'}(\mathbf{r})$, $\zeta_\mu(\mathbf{r})$ can be computed efficiently, and the scaling is bounded by $\mathcal{O}(N_\mu^3)$ (Hu et al., 2017). We observe that we can always choose $N_\mu \ll N_o N_u N_\mathbf{k}$, thus ISDF is never a bottleneck. The interpolation points are computed efficiently with centroidal Voronoi tessellation within $\mathcal{O}(N_\mu N_r)$ (Dong et al., 2018).

Inserting ISDF in the equations for the interaction kernels yields

$$V_{ou\mathbf{k},o'u'\mathbf{k}'} \approx \frac{1}{N_k^2}\sum_{\mu=1}^{N_\mu^V}\sum_{\nu=1}^{N_\mu^V} u_{o\mathbf{k}}(\mathbf{r}_\mu^V)u^*_{u\mathbf{k}}(\mathbf{r}_\mu^V)\tilde{V}_{\mu\nu}u^*_{o'\mathbf{k}'}(\mathbf{r}_\nu^V)u_{u'\mathbf{k}'}(\mathbf{r}_\nu^V),$$

$$W_{ou\mathbf{k},o'u'\mathbf{k}'} \approx \frac{1}{N_k^2}\sum_{\mu=1}^{N_\mu^{W_u}}\sum_{\nu=1}^{N_\mu^{W_o}} u^*_{u\mathbf{k}}(\mathbf{r}_\mu^{W_u})u_{u'\mathbf{k}'}(\mathbf{r}_\mu^{W_u})\tilde{W}_{\mu\nu,\mathbf{k}-\mathbf{k}'}u_{o\mathbf{k}}(\mathbf{r}_\nu^{W_o})u^*_{o'\mathbf{k}'}(\mathbf{r}_\nu^{W_o}) \ ,$$

where we have shifted the integration from the wavefunction pairs to the interpolation coefficients such that

$$\tilde{V}_{\mu\nu} = \int_{\Omega^l\times\Omega^l} drdr'\zeta^{*V}_\mu(\mathbf{r})V(\mathbf{r},\mathbf{r}')\zeta^V_\nu(\mathbf{r}') \ ,$$

$$\tilde{W}_{\mu\nu,\mathbf{k}-\mathbf{k}'} = \int_{\Omega^l\times\Omega^l} drdr'\zeta^{*W_u}_\mu(\mathbf{r})W_{\mathbf{k}-\mathbf{k}'}(\mathbf{r},\mathbf{r}')\zeta^{W_o}_\nu(\mathbf{r}') \ .$$

Note that there are three different wavefunction pairings, i.e. $u^*_{o\mathbf{k}}(\mathbf{r})u_{u\mathbf{k}}(\mathbf{r})$ for the exchange kernel (V) and $u^*_{o\mathbf{k}}(\mathbf{r})u_{o'\mathbf{k}'}(\mathbf{r})$, $u^*_{u\mathbf{k}}(\mathbf{r})u_{u'\mathbf{k}'}(\mathbf{r})$ for the screened kernel W. Each pairing requires a separate ISDF calculation, denoted by the superscripts V, W_o, and W_u. Since the number of combinations may vary, the number of interpolation points required may also vary. Reformulating the matrix elements in this way alone does not improve scaling with respect to the system size. To achieve this, we combine it with an iterative solver, here with the Lanczos algorithm. This class of algorithms constructs an approximation to the eigenvalues and eigenvectors by iteratively applying matrix-vector multiplications. Applying the interaction kernels in their interpolated forms to a vector X of dimension $N_o N_u N_\mathbf{k}$ allows efficient computation by rearranging the summations to exploit a separable structure of the kernels. For the exchange kernel we get

$$[V\cdot X]_{ou\mathbf{k}} = \frac{1}{N_\mathbf{k}}\sum_{\mu=1}^{N_\mu^V} u^*_{u\mathbf{k}}(\mathbf{r}_\mu^V)u_{o\mathbf{k}}(\mathbf{r}_\mu^V)\left\{\sum_{\nu=1}^{N_\mu^V}\tilde{V}_{\mu\nu}\left[\sum_{\mathbf{k}'}\left(\sum_{u'}u_{u'\mathbf{k}'}(\mathbf{r}_\nu^V)\left[\sum_{o'}u^*_{o'\mathbf{k}'}(\mathbf{r}_\nu^V)\cdot X_{o'u'\mathbf{k}'}\right]\right)\right]\right\},$$

where we first compute the sums over o', u', and $\mathbf{k}'$ to get a term that depends only on $\mathbf{r}_\nu^V$ with a complexity of $\mathcal{O}(N_\mu^V(N_o N_u N_\mathbf{k} + N_u N_\mathbf{k})$. The remaining sums can be computed with $\mathcal{O}((N_\mu^V)^2 N_\mu^V N_o N_u N_\mathbf{k})$, so the complexity of computing $V\cdot X$ is bounded by $\mathbf{O}((N_\mu^V)^2 + N_\mu N_o N_u N_\mathbf{k})$. Applying the screened kernel to X, after reordering the sums, we get

$$[W\cdot X]_{ou\mathbf{k}} = \frac{1}{N_k}\sum_{\nu=1}^{N_\mu^{W_o}} u_{o\mathbf{k}}(\mathbf{r}_\nu^{W_o})\left\{\sum_{\mu=1}^{N_\mu^{W_u}} u^*_{u\mathbf{k}}(\mathbf{r}_\mu^{W_u})\sum_{\mathbf{k}'}\left[\tilde{W}_{\mu\nu,\mathbf{k}-\mathbf{k}'}\right.\right.$$

DRAFT

$$\times\left(\sum_{u'} u_{u'\mathbf{k}'}(\mathbf{r}_\mu^{W_u})\left[\sum_{o'} u^*_{o'\mathbf{k}'}(\mathbf{r}_\nu^{W_o})X_{o'u'\mathbf{k}'}\right]\right)\Bigg]\Bigg\}\,.$$

Here we exploit the separable structure of the decomposition so that the terms depending on $\mathbf{k}$ and $\mathbf{k}'$ are on the left and right of $\tilde{W}_{\mu\nu,\mathbf{k}-\mathbf{k}'}$. The evaluation of the two innermost sums over o' and u' to $A^{\mathbf{k}'}_{\mu\nu}$ scales with $\mathcal{O}(N_\mu^{W_u}N_oN_uN_\mathbf{k} + N_\mu^{W_o}N_\mu^{W_u}N_uN_\mathbf{k})$. Then the sum over $\mathbf{k}'$ reads as a discrete convolution

$$\sum_{\mathbf{k}'} W_{\mathbf{k}-\mathbf{k}'}A^{\mathbf{k}'}_{\mu\nu}\,,$$

which can be efficiently evaluated with fast Fourier transforms simultaneously within the $\mathcal{O}(N\mathbf{k}\log N\mathbf{k})$ scaling for each $\mu\nu$ pair. The remaining summations scale with $\mathcal{O}(N_\mu^{W_o}N_\mu^{W_u}N_uN_\mathbf{k})$. So the complexity for the computation of $W\cdot X$ is bounded by $\mathcal{O}(N_\mu^{W_u}N_oN_uN_\mathbf{k} + N_\mu^{W_o}N_\mu^{W_u}N_uN_\mathbf{k} + N_\mu^{W_o}N_\mu^{W_u}N_\mathbf{k}\log N_\mathbf{k})$.

Statement of need

Due to the unfavorable scaling of solving the BSE directly, many interesting problems such as complex materials with large unit cells or systems requiring a dense Brioullin-zone sampling are not feasible. Even though Henneke and coworkers (Henneke et al., 2020) have already described the new algorithm and demonstrated the scaling improvement, an easy-to-use and scalable implementation was still missing. We have implemented and fully integrated this approach in the existing BSE infrastructure of the all-electron, full-potential package exciting (Gulans et al., 2014). Users can now easily choose which algorithm they prefer to use and have the full suite of exciton analysis implemented in exciting at hand.

Results

For computing the ISDF, two new parameters, $n_\mathbf{r}$ and c_μ, are introduced. $n_\mathbf{r}$ is the real-space sampling density for $u_{i\mathbf{k}}(\mathbf{r})$ and is defined as

$$n_\mathbf{r} = \frac{N_\mathbf{r}}{\Omega}\,,$$

where $N_\mathbf{r}$ is the number of $\mathbf{r}$-points and Ω the unit cell volume. The sampling is chosen to be regular such that the distance between the sampling points in each lattice direction is as similar as possible. The dimensionless parameter c_μ is used to control the number of interpolation points and is defined as

$$N_\mu = c_\mu\sqrt{\sqrt{N_{\text{pairs}}}}\,,$$

where N_{pairs} refers to the number of wave function pairs for which ISDF is computed. Note that N_{pairs} depends on $N_\mathbf{k}$. The double square-root dependence ensures that the overall scaling remains below $\mathcal{O}(N_\mathbf{k}^2)$. In Fig.(Figure 1) we present, for the example of diamond, the difference in exciton binding energies obtained with the new implementation and a reference calculation. The reference, based on the direct implementation, sets up and diagonalizes the full BSH and depends neither on $n_\mathbf{r}$ nor c_μ. The results are shown as functions of $n_\mathbf{r}$ and c_μ for a small $\mathbf{k}$-grid of $2\times2\times2$. Additionally, we show the spectral similarities compared to the reference calculation as functions of $n_\mathbf{r}$ and c_μ. For both parameters, both properties converge as their values increase. To find the optimal interpolation grid for ISDF, we first converge $n_\mathbf{r}$, then c_μ. In our example, $n_\mathbf{r} = 138$ [a.u.] and $c_\mu = 40.0$ yield converged results. This corresponds to a real-space sampling of $22\times22\times22$ and numbers of interpolation points $N_\mu^V = 202$, $N_\mu^{W_o} = 322$, and $N_\mu^{W_u} = 360$.

DRAFT



Figure 1: Difference of the exciton binding energy and spectral similarity between the new method and the direct method as functions of $n_\mathbf{r}$ (upper panel) and c_μ (lower panel) for diamond on a $2 \times 2 \times 2$ $\mathbf{k}$-grid.

In Fig.(Figure 2) we compare the exciton binding energies and spectra of the new implementation to those of the old implementation for increasing $\mathbf{k}$-grids, while keeping $n_\mathbf{r}$ as well as N_μ^V, $N_\mu^{W_o}$, and $N_\mu^{W_u}$ fixed at the values above. We observe that the results converge as $N_\mathbf{k}$ increases. Thus, the number of interpolation points is asymptotically independent of $N_\mathbf{k}$. A similar behavior of ISDF was observed in (Lu & Ying, 2016).

DRAFT



Figure 2: Difference of the exciton binding energy and spectral similarity between the new method and the direct method as functions of $N_\mathbf{k}$ for diamond.

In Fig.(Figure 3) we show the wall times for solving the BSE with the new and direct implementations for increasing $N_\mathbf{k}$. The new algorithm massively outperforms the direct one, and the speedup increases more than linearly with $N_\mathbf{k}$. We also show the wall times for computing the RPA screening with increasing $N_\mathbf{k}$, which is now clearly the bottleneck in solving the BSE.



Figure 3: Runtimes of the direct (black) and new (red) BSE implementations and the RPA screening (blue) as functions of $N_\mathbf{k}$. The speedup of the algorithm is shown by the gray dashed line.

Altogether, we have implemented a new, low-scaling BSE solver and fully integrated it into the all-electron, full-potential package exciting. We demonstrate that the new implementation yields results equivalent to the direct solution of the BSE but with significantly reduced computational time. Consequently, it enables more precise calculations and facilitates the study of more complex problems.

References

Dong, K., Hu, W., & Lin, L. (2018). Interpolative separable density fitting through centroidal voronoi tessellation with applications to hybrid functional electronic structure calculations. *Journal of Chemical Theory and Computation*, *14*, 1311–1320. https://doi.org/10.1021/acs.jctc.7b01113

Gulans, A., Kontur, S., Meisenbichler, C., Nabok, D., Pavone, P., Rigamonti, S., Sagmeister, S., Werner, U., & Draxl, C. (2014). Exciting: A full-potential all-electron package implementing density-functional theory and many-body perturbation theory. *Journal of Physics: Condensed Matter*, *26*, 363202. https://doi.org/10.1088/0953-8984/26/36/363202

Henneke, F., Lin, L., Vorwerk, C., Draxl, C., Klein, R., & Yang, C. (2020). Fast optical absorption spectra calculations for periodic solid state systems. *Communications in Applied Mathematics and Computational Science*, *15*, 89–113. https://doi.org/10.2140/camcos.2020.15.89

Hu, W., Lin, L., & Yang, C. (2017). Interpolative separable density fitting decomposition for accelerating hybrid density functional calculations with applications to defects in silicon. *Journal of Chemical Theory and Computation*, *13*, 5420–5431. https://doi.org/10.1021/acs.jctc.7b00807

Lu, J., & Ying, L. (2015). Compression of the electron repulsion integral tensor in tensor hypercontraction format with cubic scaling cost. *Journal of Computational Physics*, *302*, 329–335. https://doi.org/10.1016/j.jcp.2015.09.014

Lu, J., & Ying, L. (2016). Fast algorithm for periodic density fitting for bloch waves. *Annals of Mathematical Sciences and Applications*, *1*(2), 321–339. https://doi.org/10.4310/AMSA.2016.v1.n2.a3

Vorwerk, C., Aurich, B., Cocchi, C., & Draxl, C. (2019). Bethe–salpeter equation for absorption and scattering spectroscopy: Implementation in the exciting code. *Electronic Structure*, *1*, 037001–037001. https://doi.org/10.1088/2516-1075/ab3123

DRAFT