The Journal of Open Source Software

mcpy: Canonical & Grand Canonical Monte Carlo for atomistic systems with machine-learning potentials

Riccardo Farris [1,2], **Emanuele Telari** [1], **Konstantin M. Neyman** [1,3], and **Albert Bruix** [1]

1 Institut de Química Teòrica i Computacional (IQTCUB), Universitat de Barcelona, Barcelona, Spain **2** LEITAT Technological Center, Terrassa, Spain **3** Institució Catalana de Recerca i Estudis Avançats (ICREA), Barcelona, Spain

DOI: 10.xxxxxx/draft

Software

- Review
- Repository
- Archive

Editor: Rohit Goswami

Reviewers:

- @isdanni
- @tonigi
- @ptmerz

Submitted: 22 June 2026
Published: unpublished

License
Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License (CC BY 4.0).

Summary

mcpy is a Python package for Monte Carlo (MC) simulations of atomistic systems (surfaces, nanoparticles, fluids, and bulk) at controlled temperature, volume, and particle number or chemical potential. It is built on the Atomic Simulation Environment (ASE) (Larsen et al., 2017), so any ASE-compatible calculator can drive the sampling: density-functional theory codes, classical force fields, or machine-learning interatomic potentials (MLIPs) such as MACE (Batatia et al., 2022). mcpy implements canonical and grand canonical ensembles, as well as replica-exchange approaches (Swendsen & Wang, 1986), in modular run loops and runs with or without local relaxation of trial moves. Following the hybrid scheme of Senftle et al. (2014), insertions and deletions can be relaxed locally before the acceptance test, which makes sampling efficient in grand canonical ensemble simulations of densely packed metallic systems where rigid insertions would almost always be rejected; relaxation can equally be switched off for standard MC of fluids and other systems. The same loop runs on CPUs via MPI or on GPUs through different backends such as NVIDIA Alchemi, which relaxes a batch of replicas in a single evaluation. The package provides modular trial moves (insertion, deletion, displacement, permutation (enabling chemical-ordering optimization in multi-component systems), shake, and short MD runs via Brownian/Langevin moves), configurable cell geometries (periodic box, rectangular sub-slab, spherical region around a nanoparticle, and user-defined cells), global optimization and statistical sampling, together with post-processing utilities that turn raw ensembles into surface and nanoparticle phase diagrams.

Statement of need

Predicting the equilibrium structure and composition of surfaces and nanoparticles under reactive atmospheres requires sampling over candidate structures and compositions (i.e. particle number) in the grand canonical ensemble at near–first-principles accuracy. MLIPs now deliver close-to-DFT accuracy at a small fraction of the cost, making large-scale GCMC of realistic structures increasingly practical. Yet no open-source package combines relaxation-coupled GCMC, replica exchange, an ASE-native MLIP backend, and the cell geometries needed for finite nanoparticles in a single, modular workflow.

mcpy's grand canonical simulations are aimed at computational chemists and materials scientists studying heterogeneous catalysis, oxidation, and gas–surface equilibria, where the stable composition, not just a fixed structure, is the quantity of interest. Although it was developed for these applications, the grand canonical sampling is agnostic to the level of theory or the targeted system: it runs on any system ASE can represent and any ASE-compatible calculator, from molecular fluids to bulk solids. The same workflow therefore extends well beyond the

DRAFT

metallic surfaces and nanoparticles that motivated it.

State of the field

Several established codes perform Monte Carlo sampling of atomistic systems, but each covers only part of the space `mcpy` targets. RASPA2 (Dubbeldam et al., 2016) and its recent C++ rewrite RASPA3 (Ran et al., 2024) are specialized for adsorption and diffusion of molecules in nanoporous materials with classical force fields. DL_MONTE (Brukhno et al., 2021) is a general-purpose Monte Carlo code that supports GCMC of solids and fluids, but with classical interaction models and its own input ecosystem rather than native integration with MLIPs or ASE. LAMMPS (Thompson et al., 2022) provides a GCMC fix, parallel tempering, and machine-learning pair styles, but it neither couples replica exchange to grand canonical sampling nor exposes a relaxation-coupled GCMC in which an arbitrary ASE calculator drives each acceptance test. ASE itself (Larsen et al., 2017) supplies the `Atoms` object and a unified calculator interface spanning DFT codes, classical force fields, and MLIPs, but no grand canonical ensemble built on top of it.

We built `mcpy` as a new package rather than extending any one of these because the capabilities it unifies live in different ecosystems: the calculator-agnostic accuracy of ASE, the grand canonical and replica-exchange machinery of dedicated Monte Carlo codes, and the geometric flexibility needed for finite nanoparticles. Adding MLIP-native, relaxation-coupled GCMC to RASPA2 or DL_MONTE would require reimplementing ASE's calculator abstraction, while wrapping LAMMPS in a relaxation-coupled, ASE-native grand canonical loop with free-volume cell geometries and *ab initio* thermodynamics post-processing would reproduce much of `mcpy`'s MC design around an MD-centric engine. Building directly on ASE instead lets `mcpy` inherit the entire ASE calculator ecosystem and contribute the missing grand canonical layer. That layer is the package's distinct contribution: ASE-native, MLIP-driven, relaxation-coupled GCMC with replica exchange and nanoparticle/surface geometries.

Software design

`mcpy` is organized around four extensible abstractions: *ensembles* that drive the Monte Carlo loop (`GrandCanonicalEnsemble`, `CanonicalEnsemble`, `ReplicaExchange`), *moves* that propose trial configurations (insertion, deletion, displacement, permutation, shake, and Brownian), *cells* that define the sampling region and estimate its free volume (periodic box, rectangular sub-slab, spherical region around a nanoparticle, and user-defined cells), and *calculators* that wrap ASE energy evaluations. This separation reflects the central design choice: by expressing the physics through ASE's `Atoms` and calculator interfaces, `mcpy` trades the raw speed of a monolithic, hand-tuned Monte Carlo kernel for the ability to drive identical sampling code with any ASE-compatible backend, from a DFT code to a MACE potential (Batatia et al., 2022). Because the dominant cost in the target application is the energy evaluation rather than the Monte Carlo bookkeeping, that trade-off is strongly favorable, and it lets a researcher exchange accuracy for cost without rewriting the simulation.

The second design choice is an optional relaxation-coupled acceptance criterion. Following the hybrid scheme of Senftle et al. (2014), each trial insertion or deletion can be relaxed locally before the acceptance test. This adds cost to each step, but in densely packed metallic surfaces and nanoparticles a rigid insertion overlaps neighboring atoms and is almost always rejected; the local relaxation is what makes grand canonical sampling converge at all for these systems, so the per-step cost is the enabling trade-off rather than an overhead. Because the relaxation is delegated to ASE, any of ASE's local-relaxation algorithms can be selected according to its suitability for the chosen system and level of theory. Conversely, the relaxation step can be disabled entirely, recovering standard unbiased Metropolis Monte Carlo that samples the exact Boltzmann distribution, appropriate for fluids and other systems where rigid trial moves are

DRAFT

readily accepted.

The third choice concerns ergodicity and parallelism. Replica exchange (Swendsen & Wang, 1986) across temperatures or chemical potentials improves mixing, and `mcpy` exposes two complementary parallelization paths: over MPI ranks on CPUs, one replica per rank, or on a GPU through the NVIDIA Alchemi backend, which relaxes a batch of replicas together in a single evaluation. The MPI path is portable and dependency-light (`mpi4py` is optional); the GPU path trades that portability for throughput on large systems where batched relaxation dominates. Because the ensemble loop is decoupled from the calculator, the same replica-exchange code runs under either path.

Finally, the modular loop is reused beyond equilibrium sampling: every ensemble doubles as a basin-hopping global optimizer, emitting a running trajectory of strictly improving (minimum-energy) configurations, and replica-exchange runs additionally report the global minimum found across replicas. Compound moves that perturb several atoms per trial make these basin-hopping escapes effective in densely packed systems. Simulations log per-step energies, particle counts, and per-move acceptance ratios, and write extended-XYZ trajectories alongside an optional `minima` trajectory of strictly improving structures. The bundled phase-diagram utilities map sampled ensembles onto *ab initio* thermodynamics phase diagrams (Reuter & Scheffler, 2001), connecting the simulated chemical potential to experimental temperature and partial pressure.

To validate the grand canonical sampling independently of any MLIP, we reproduced two Lennard-Jones reference systems against LAMMPS (Thompson et al., 2022) (Figure 1): a purely repulsive Lennard-Jones fluid in reduced units and real argon in physical (eV) units. Both codes run the same potential, initial state, move mix, and number of trial moves, with four independently seeded runs per chemical potential. The average particle number agrees within statistical error across both unit systems, including the sharp gas–liquid condensation of argon. The script that produces the figure, `benchmark/lammps_lj_isotherms.py`, and the data behind it, `benchmark/lj_isotherms/lj_isotherms.csv`, are part of the repository. Although `mcpy` has so far been applied mainly to metal surfaces and nanoparticles, this benchmark confirms that its grand canonical machinery is correct and general, applicable to atomistic materials beyond the metallic systems that motivated it.

DRAFT

The Journal of Open Source Software




Figure 1: Validation of `mcpy`'s grand canonical sampling against LAMMPS (Thompson et al., 2022) for two Lennard-Jones reference systems. (a) A purely repulsive Lennard-Jones fluid in reduced units ($T^* = 1$). (b) Argon in physical units ($T = 87.79$ K) across the gas–liquid condensation. Top: average particle number $\langle N \rangle$ versus chemical potential μ for `mcpy` (red) and LAMMPS (blue). Bottom: the difference between the two codes, with error bars and a grey band of two combined standard errors over four independent runs per code.

Research impact statement

`mcpy` was developed to study the equilibrium oxidation of metal nanoparticles and surfaces under reactive atmospheres at near-DFT accuracy. The replica-exchange GCMC workflow was built during doctoral research at the Universitat de Barcelona (Farris, 2025) and underpins a forthcoming manuscript on the oxidation thermodynamics of silver nanoparticles, with further application to additional catalytic metal/oxide systems. Simulations run with `mcpy` have contributed to the group's grand canonical modelling of nanostructured catalysts under reaction conditions, presented at recent international conferences on cluster science and on machine learning for materials (Bruix, 2026; Telari et al., 2026). Figure 2 shows a representative result from this workflow: the hydrogenation phase diagram of a 201-atom PdAg nanoparticle, in which `mcpy` recovers the progressive hydrogen loading of the particle as the chemical potential increases. By coupling MLIP-driven energetics with grand canonical sampling and *ab initio* thermodynamics post-processing, `mcpy` predicts stable surface and nanoparticle compositions as a function of temperature and chemical potential, quantities that are otherwise inaccessible to fixed-composition relaxations. The package is open-source, documented with worked examples covering GCMC, replica exchange, and phase-diagram construction, and continuously tested across Python 3.11–3.13, so that the published workflows can be reproduced and extended by other groups.

DRAFT

The Journal of Open Source Software




Figure 2: (a) Hydrogenation phase diagram of a 201-atom palladium–silver nanoparticle ($Pd_{151}Ag_{50}$) obtained with replica-exchange GCMC in `mcpy`, driven by the small AGNESI MACE potential (Batatia et al., 2025). The free energy of formation ΔG (in eV) is shown as a function of the hydrogen chemical potential $\Delta\mu_H$. The top axis converts $\Delta\mu_H$ into $\log_{10}(p_{H_2}/p_0)$ at $T = 300$ K with $p = p_0$ at $\Delta\mu_H = 0$. This neglects the standard chemical potential of H_2 at 300 K (-0.16 eV per H) and the H_2 zero-point energy ($+0.14$ eV per H), so the true pressures are about 0.8 decades higher than the axis shows. The insets show some of the most stable $Pd_{151}Ag_{50}H_z$ structures sampled for different chemical-potential windows, from the bare particle at low μ_H to a hydrogen-loaded $Pd_{151}Ag_{50}H_{123}$ at high pressure, highlighting the change in chemical ordering induced by the increase in hydrogen coverage. The chemical ordering of the bare $Pd_{151}Ag_{50}$ particle was first optimized with a single basin-hopping run; hydrogenated structures were then sampled by replica-exchange GCMC at hydrogen chemical potentials from $\Delta\mu_H = -1$ to -0.35 eV with six temperatures per chemical-potential value, using H insertion, H deletion, and Pd-Ag swap moves. (b) Evolution of the number of inserted oxygen atoms during a replica-exchange GCMC simulation of a Pd_{55} fcc nanoparticle supported on a Al_2O_3 surface. The run consisted of 12 replicas with different temperatures at a fixed $\Delta\mu_O = -0.5$. The structural insets show representative configurations sampled along the trajectories of the different replicas.

Acknowledgements

This work was supported by the Spanish/FEDER Ministerio de Ciencia, Innovacion y Universidades [Grant Nos. PID2021-128217NB-I00, MDM-2017-0767, CEX2021-001202-M, PID2022-140120OA-I00, and RYC2021-032281-I (for A.B.)] as well as by the Generalitat de Catalunya [Grant No. 2021SGR00286]. R. F. thanks the Spanish MICIUN for an FPI PhD grant (MDM-2017-0767-20-2). R. F. further acknowledges his current research group for supporting the continuation and development of this work.

AI usage disclosure

Claude (Opus) assisted with code refactoring, test scaffolding, packaging configuration, and editing of the manuscript and documentation. All AI-assisted outputs were reviewed, edited, and validated by the human authors, who designed the overall code architecture and take full responsibility for the accuracy and originality of the software and all submitted materials.

DRAFT

References

Batatia, I., Benner, P., Chiang, Y., Elena, A. M., Kovács, D. P., & others. (2025). A foundation model for atomistic materials chemistry. *The Journal of Chemical Physics*, *163*(18), 184110. https://doi.org/10.1063/5.0297006

Batatia, I., Kovács, D. P., Simm, G. N. C., Ortner, C., & Csányi, G. (2022). MACE: Higher order equivariant message passing neural networks for fast and accurate force fields. *Advances in Neural Information Processing Systems (NeurIPS)*. https://doi.org/10.48550/arXiv.2206.07697

Bruix, A. (2026). Accelerating the structural and chemical characterization of nanostructured materials under reaction conditions with ML-guided grand canonical global optimization. *Book of Abstracts, Artificial Intelligence for Advanced Materials (AI4AM 2026)*. https://phantomsfoundation.com/AI4AM/2026/Abstracts/AI4AM2026_Bruix_Albert_58.pdf

Brukhno, A. V., Grant, J., Underwood, T. L., Stratford, K., Parker, S. C., Purton, J. A., & Wilding, N. B. (2021). DL_MONTE: A multipurpose code for Monte Carlo simulation. *Molecular Simulation*, *47*(2-3), 131–151. https://doi.org/10.1080/08927022.2019.1569760

Dubbeldam, D., Calero, S., Ellis, D. E., & Snurr, R. Q. (2016). RASPA: Molecular simulation software for adsorption and diffusion in flexible nanoporous materials. *Molecular Simulation*, *42*(2), 81–101. https://doi.org/10.1080/08927022.2015.1010082

Farris, R. (2025). *Development and application of advanced computational methods for predicting the structure and properties of nanoparticles* [PhD thesis, Universitat de Barcelona]. https://hdl.handle.net/10803/697698

Larsen, A. H., Mortensen, J. J., Blomqvist, J., & others. (2017). The atomic simulation environment—a Python library for working with atoms. *Journal of Physics: Condensed Matter*, *29*(27), 273002. https://doi.org/10.1088/1361-648X/aa680e

Ran, Y. A., Sharma, S., Balestra, S. R. G., Li, Z., Calero, S., Vlugt, T. J. H., Snurr, R. Q., & Dubbeldam, D. (2024). RASPA3: A Monte Carlo code for computing adsorption and diffusion in nanoporous materials and thermodynamics properties of fluids. *The Journal of Chemical Physics*, *161*(11), 114106. https://doi.org/10.1063/5.0226249

Reuter, K., & Scheffler, M. (2001). Composition, structure, and stability of RuO_2(110) as a function of oxygen pressure. *Physical Review B*, *65*(3), 035406. https://doi.org/10.1103/PhysRevB.65.035406

Senftle, T. P., Janik, M. J., & Duin, A. C. T. van. (2014). A ReaxFF investigation of hydride formation in palladium nanoclusters via Monte Carlo and molecular dynamics simulations. *The Journal of Physical Chemistry C*, *118*(9), 4967–4981. https://doi.org/10.1021/jp411015a

Swendsen, R. H., & Wang, J.-S. (1986). Replica Monte Carlo simulation of spin-glasses. *Physical Review Letters*, *57*(21), 2607–2609. https://doi.org/10.1103/PhysRevLett.57.2607

Telari, E., Farris, R., Quinlivan Domínguez, J., Neyman, K. M., & Bruix, A. (2026). Modelling nanoparticle transformations in reactive environments. *Book of Abstracts, CSI 2026 – Cluster-Surface Interactions*, 26. https://csi2026.sciencesconf.org/data/pages/Book_Abstracts.pdf

Thompson, A. P., Aktulga, H. M., Berger, R., Bolintineanu, D. S., Brown, W. M., Crozier, P. S., Veld, P. J. in 't, Kohlmeyer, A., Moore, S. G., Nguyen, T. D., Shan, R., Stevens, M. J., Tranchida, J., Trott, C., & Plimpton, S. J. (2022). LAMMPS - a flexible simulation tool for particle-based materials modeling at the atomic, meso, and continuum scales. *Computer Physics Communications*, *271*, 108171. https://doi.org/10.1016/j.cpc.2021.108171

DRAFT