

ODINN.jl: Scientific machine learning glacier modelling

Jordi Bolibar^{1,2}, Facundo Sapienza^{3,4}, Alban Gossard¹, Lisa Girod¹, Mathieu le Séac'h¹, Lucille Gimenes¹, Vivek Gajadhar², Fabien Maussion^{5,6}, Ching-Yao Lai³, Bert Wouters², and Fernando Pérez⁴

¹ Univ. Grenoble Alpes, CNRS, IRD, G-INP, Institut des Géosciences de l'Environnement, Grenoble, France ² Faculty of Civil Engineering and Geosciences, Delft University of Technology, Delft, The Netherlands ³ Department of Geophysics, Stanford University, Stanford, United States ⁴ Department of Statistics, University of California, Berkeley, United States ⁵ Bristol Glaciology Centre, School of Geographical Sciences, University of Bristol, Bristol, UK ⁶ Department of Atmospheric and Cryospheric Sciences, University of Innsbruck, Innsbruck, Austria ¶ Corresponding author

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#)
- [Repository](#)
- [Archive](#)

Editor: [Anjali Sandip](#)

Reviewers:

- [@haakon-e](#)
- [@luraess](#)

Submitted: 22 January 2026

Published: unpublished

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

Summary

ODINN.jl is a glacier model leveraging scientific machine learning (SciML) methods to perform forward and inverse simulations of large-scale glacier evolution. It can simulate both surface mass balance and ice flow dynamics through a modular architecture which enables the user to easily modify model components.

The most unique aspect of ODINN.jl is its differentiability and capabilities of performing all sorts of different hybrid modelling. Since the whole ecosystem is differentiable (where differentiable means the ability to compute model derivatives with respect to parameters (Shen et al., 2023)), we can optimize almost any model component, providing a powerful framework to tackle many scientific problems (Bolibar et al., 2023). ODINN.jl can optimize, separately or together, in a steady-state (time-independent simulation) or transient (time-dependent simulation) way the following:

- The initial or intermediate state of glaciers (i.e. their ice thickness) or the equivalent ice surface velocities.
- Model parameters (e.g. the Glen coefficient A related to ice viscosity in a 2D Shallow Ice Approximation (Hutter, 1983)), in a gridded or scalar format. This can be done for multiple time steps where observations (e.g. ice surface velocities) are available.
- The parameters of a statistical regressor (e.g. a neural network), used to parametrize a subpart or one or more coefficients of an ice flow or surface mass balance mechanistic model. This enables the exploration of empirical laws describing physical processes of glaciers, leveraging Universal Differential Equations (UDEs, Rackauckas et al. (2021)).

For this, it is necessary to differentiate (that is, computing gradients or derivatives) through complex code, including numerical solvers, which is a non-trivial task (Sapienza et al., 2024). We use reverse mode automatic differentiation (AD) based on the adjoint method to achieve this. We have two strategies for computing both the adjoint and the required vector-jacobian products (VJPs): (1) hand-derived adjoint equations, evaluated via AD using Enzyme.jl (Moses et al., 2021), as well as fully manual implementations of the spatially discrete and spatially continuous VJPs; and (2) automatic adjoints using SciMLSensitivity.jl (Rackauckas et al., 2019), available with different AD back-ends for the VJPs computation. These two approaches are complementary, with the manual adjoints being ideal for high-performance tasks by providing more control on the implementation, and serving as a ground truth for benchmarking and testing automatic adjoint methods from SciMLSensitivity.jl.

Beyond all these inverse modelling capabilities, ODINN.jl can also act as a more conventional forward glacier model, simulating glaciers in parallel, and easily customizing different model parametrizations and choices within the simulation. Its high modularity, combined with the easy access to a vast collection of datasets coming from the Open Global Glacier Model (OGGM, Maussion et al. (2019)), makes it very easy to run simulations, even with an average laptop. Multiple ice flow dynamics models can be easily swapped, thanks to a modular architecture (see Software design). Models based on partial differential equations (PDEs) are solved using DifferentialEquations.jl (Rackauckas & Nie, 2017), which provides access to a large amount of numerical solvers. For now, we have implemented a 2D Shallow Ice Approximation (SIA, Hutter (1983)), but in the future we plan to incorporate other models, such as the Shallow Shelf Approximation (SSA, Weis et al. (1999)). Validation of numerical forward simulations is evaluated in the test suite based on exact analytical solutions of the SIA equation (Bueler et al., 2005). Multiple surface mass balance models are available, based on simple temperature-index models. Nonetheless, the main addition of the upcoming version will be the machine learning-based models from the MassBalanceMachine (Sjursen et al., 2025), which will provide further mass balance models.

Statement of need

ODINN.jl addresses the need for a glacier model that combines the physical interpretability of mechanistic approaches with the flexibility and data-assimilation capabilities of data-driven methods (Bolibar et al., 2023). By integrating both paradigms, it enables targeted inverse methods to learn parametrizations of glacier processes, capturing unknown physics while preserving the physically grounded structure of glacier dynamics through differential equations.

While purely mechanistic and purely data-driven glacier models already exist (e.g. Gagliardini et al. (2013), Maussion et al. (2019), Rounce et al. (2023), Bolibar et al. (2022)), they often do not fully exploit growing glacier observations, including surface velocity, ice thickness, topography, surface mass balance, and climate reanalyses. Existing empirical laws are not always directly linked to these observables, so calibration remains challenging. Differentiable programming and functional inversion offer a path forward by enabling new empirical relationships from proxies and testing hypotheses about poorly constrained processes such as basal sliding, creep, and calving.

Improving the representation of these processes is crucial for reliable projections of glacier evolution and impacts on freshwater availability and sea-level rise (IPCC, 2021). To this end, ODINN.jl provides a unified modelling ecosystem supporting advanced inverse calibration and efficient modular forward simulations for large-scale glacier studies.

Developing such a framework places strong demands on scientific software. Inefficient code and irreproducible implementations can restrict progress, emphasizing open-source, community-driven tools that follow modern research software engineering (RSE) practices (Combemale et al., 2023). These tools should support modular design for rapid prototyping and extension of workflows (Nyenah et al., 2024). Julia provides two key advantages: it addresses the two-language problem by combining high-level expressiveness with high-performance (Bezanson et al., 2017), and it enables source-code differentiability for modular gradient-based inverse optimization and hybrid modelling, including UDEs. With ODINN.jl, we aim to create a robust, future-proof framework bridging physical understanding and data-driven discovery. Its modular architecture, testing, and continuous integration (CI) ensure reproducibility and reliability, while its open design enables collaboration across glaciology and Earth system modelling.




Figure 1: Overview of the ODINN.jl ecosystem.

State of the field

The field of large-scale glacier modelling has considerably grown over the last decade, with most models being (or becoming) open-source. While global glacier models such as OGGM (MauSSION et al., 2019), GloGEM (Huss & Hock, 2015) and PyGEM (Rounce et al., 2023) target global-scale efficient simulations of past and future glacier changes and sea-level rise contributions, ODINN.jl has a focus on inverse modelling for glacier physical processes and catchment-scale glacio-hydrological simulations. Moreover, ODINN.jl has the potential to implement more robust glacier physics compared to global models focusing on simplicity and lightweight parametrizations. IGM (Jouvet, 2023), with its neural network-driven ice flow emulator approach has some parallels to ODINN.jl, but the model's focus has mainly been on speeding up simulations for large-scale or paleo simulations. Alternatively, the new ice sheet model DJuice.jl (Moses et al., n.d.) – a Julia translation of the well-known ISSM model – is perhaps the most similar model to ODINN.jl. They both leverage Julia's differentiability capabilities, and aim to exploit them to infer parametrizations and physical processes related to ice flow. Nonetheless, while DJuice.jl is an ice sheet model, ODINN.jl is highly specialized to simulate glaciers (i.e. anything outside Greenland and Antarctica), thanks to its compatibility with OGGM's data preprocessing.

Software design

ODINN.jl is an ecosystem composed of multiple packages, each one handling a specific task:

- Sleipnir.jl: Handles all the basic types, functions and datasets, common through the

whole ecosystem, as well as data management tasks.

- `Muninn.jl`: Handles surface mass balance processes, via different types of models.
- `Huginn.jl`: Handles ice flow dynamics, by solving the ice flow partial differential equations (PDEs) using numerical methods. It can accommodate multiple types of ice flow models.
- `ODINN.jl`: Acts as the interface to the whole ecosystem, and provides the necessary tools to differentiate and optimize any model component. It can be seen as the SciML layer, enabling different types of inverse methods, using hybrid models combining differential equations with data-driven models.

Splitting large Julia ([Bezanson et al., 2017](#)) packages into smaller, focused subpackages is a good practice that enhances maintainability, usability, and collaboration. Modular design simplifies debugging, testing, and updates by isolating functionalities, while users benefit from faster precompilation and reduced memory overhead by loading only the subpackages they need. This approach also lowers the barrier for new contributors, fosters clearer dependency management, and ensures scalability as projects grow, ultimately creating a robust and adaptable software ecosystem. The ODINN ecosystem extends beyond this suite of Julia packages, by leveraging the data preprocessing tools of OGGM. This is done via the auxiliary Python library `Gungnir`, which is responsible for generating all the necessary data to initialize and run the model, such as glacier outlines from the Randolph Glacier Inventory (RGI Consortium (2023), RGI), digital elevation models (DEMs), ice thickness observations from `GlaThiDa` ([GlaThiDa Consortium, 2020](#)), ice surface velocities from different studies ([Millan et al., 2022](#)), and different sources of climate reanalyses and projections ([Eyring et al., 2016](#); [Lange, 2019](#)). This implies that `ODINN.jl`, like OGGM, is virtually capable of simulating any of the ~274,000 glaciers on Earth ([RGI Consortium, 2023](#)).

`ODINN.jl` provides a high-level user-friendly interface, enabling the user to swap and replace most elements of a glacier simulation in a modular fashion. The main elements of a simulation, such as the `Parameters`, a `Model`, and a `Simulation` (i.e. a `Prediction` or an `Inversion`), are all objects that can be easily modified and combined. In a few lines of code, the user can automatically retrieve all necessary information for most glaciers on Earth, compose a `Model` based on a specific combination of surface mass balance and ice flow models, and incorporate data-driven models (e.g. a neural network) to parametrize specific physical processes of any of these components. Both forward and inverse simulations run in parallel using multiprocessing, leveraging Julia's speed and performance. Graphics Processing Unit (GPU) compatibility is still not ready, due to the difficulties of making GPU architectures compatible with automatic differentiation (AD). Nonetheless, it is planned for future versions.

Research impact statement

`ODINN.jl` has evolved through the last five years with code contributions during three postdoctoral positions, one PhD, and three master's internships. It has so far been used to explore the use of UDEs to invert hidden empirical laws in a synthetic glacier setup, where a prescribed rheological law was successfully recovered using a neural network ([Bolibar et al., 2023](#)). This proof-of-concept then served as a backbone to create the current architecture of `ODINN.jl`, finalized with the recent 1.0 release. The main changes and scientific goals of this large software development investment are the capacity to now apply these methods to large-scale remote sensing data for multiple glaciers, which will enable the exploration of new glacier basal sliding laws directly from heterogeneous observations, which remains a long-standing problem in glaciology ([Minchew & Joughin, 2020](#)). The development of the differentiable programming methods in `ODINN.jl` also served as a catalyst to write an exhaustive review paper, together with other key players in this community, on differentiable programming for differential equations ([Sapienza et al., 2024](#)).

Additionally, `ODINN.jl` is currently being used in a newly funded 4-year project, to simulate past and future glacier changes in several catchments in the Andes and the Alps. These model outputs will then be combined with a hydrological model to investigate the impacts of glacier

160 retreat on the hydrological regimes and drought mitigation under different climate change
161 scenarios.

162 With these two research venues, ODINN.jl will continue to be developed and used to pursue
163 both fundamental research on glacier modelling, and applied research to assess the impact of
164 glacier retreat on freshwater availability and drought mitigation.

165 AI usage disclosure

166 Generative AI has been used to generate docstrings, documentation, and to assist in the coding
167 of tests, debugging and implementing small new features over the existing architecture.

168 Acknowledgements

169 We acknowledge the help of Chris Rackauckas for the debugging and discussion of issues
170 related to the SciML Julia ecosystem, Redouane Lguensat for scientific discussions on the first
171 prototype of the model, and Julien le Sommer for scientific discussions around differentiable
172 programming. We thank all the developers of the SciML Julia ecosystem who work in each
173 one of the core libraries used within ODINN.jl. JB acknowledges financial support from the
174 Nederlandse Organisatie voor Wetenschappelijk Onderzoek, Stichting voor de Technische
175 Wetenschappen (Vidi grant 016.Vidi.171.063), a TU Delft Climate Action grant and a PEPR
176 TRACCS grant from the Agence Nationale de la Recherche as part of France 2030 (reference
177 ANR-25-EXTR-0006). FS and CYL were supported by NSF via grant number OPP-2441132 and
178 the Alfred P. Sloan Foundation under grant number FG-2024-21649. FS and FP acknowledge
179 funding from the National Science Foundation (EarthCube programme under awards 1928406
180 and 1928374). AG acknowledges funding from the MIAI cluster and Agence Nationale de la
181 Recherche (ANR) in the context of France 2030 (grant ANR-23-IACL-0006).

182 References

- 183 Bezanson, J., Edelman, A., Karpinski, S., & Shah, V. B. (2017). Julia: A Fresh Approach to
184 Numerical Computing. *SIAM Review*, 59(1), 65–98. <https://doi.org/10.1137/141000671>
- 185 Bolibar, J., Rabatel, A., Gouttevin, I., Zekollari, H., & Galiez, C. (2022). Nonlinear sensitivity
186 of glacier mass balance to future climate change unveiled by deep learning. *Nature*
187 *Communications*, 13(1), 409. <https://doi.org/10.1038/s41467-022-28033-0>
- 188 Bolibar, J., Sapienza, F., Maussion, F., Lguensat, R., Wouters, B., & Pérez, F. (2023). Universal
189 differential equations for glacier ice flow modelling. *Geoscientific Model Development*,
190 16(22), 6671–6687. <https://doi.org/10.5194/gmd-16-6671-2023>
- 191 Bueler, E., Lingle, C. S., Kallen-Brown, J. A., Covey, D. N., & Bowman, L. N. (2005).
192 Exact solutions and verification of numerical models for isothermal ice sheets. *Journal of*
193 *Glaciology*, 51(173), 291–306. <https://doi.org/10.3189/172756505781829449>
- 194 Combemale, B., Gray, J., & Rumpe, B. (2023). Research software engineering and the
195 importance of scientific models. *Software and Systems Modeling*, 22(4), 1081–1083.
196 <https://doi.org/10.1007/s10270-023-01119-z>
- 197 Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., & Taylor, K.
198 E. (2016). Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6)
199 experimental design and organization. *Geoscientific Model Development*, 9(5), 1937–1958.
200 <https://doi.org/10.5194/gmd-9-1937-2016>
- 201 Gagliardini, O., Zwinger, T., Gillet-Chaulet, F., Durand, G., Favier, L., Fleurian, B. de,
202 Greve, R., Malinen, M., Martín, C., Råback, P., Ruokolainen, J., Sacchettini, M., Schäfer,

- 203 M., Seddik, H., & Thies, J. (2013). Capabilities and performance of Elmer/Ice, a new-
 204 generation ice sheet model. *Geoscientific Model Development*, 6(4), 1299–1318. <https://doi.org/10.5194/gmd-6-1299-2013>
 205
- 206 GlaThiDa Consortium. (2020). *Glacier Thickness Database 3.1.0*. World Glacier Monitoring
 207 Service, Zurich, Switzerland.
- 208 Huss, M., & Hock, R. (2015). A new model for global glacier change and sea-level rise.
 209 *Frontiers in Earth Science*, 3. <https://doi.org/10.3389/feart.2015.00054>
- 210 Hutter, K. (1983). *Theoretical Glaciology*. Springer Netherlands. <https://doi.org/10.1007/978-94-015-1167-4>
 211
- 212 IPCC. (2021). *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change: Vols. In Press*. Cambridge University Press. <https://doi.org/10.1017/9781009157896>
 213
 214
- 215 Jouvett, G. (2023). Inversion of a Stokes glacier flow model emulated by deep learning. *Journal of Glaciology*, 69(273), 13–26. <https://doi.org/10.1017/jog.2022.41>
 216
- 217 Lange, S. (2019). *WFDE5 over land merged with ERA5 over the ocean (W5E5)*. GFZ Data
 218 Services. <https://doi.org/10.5880/PIK.2019.023>
- 219 Maussion, F., Butenko, A., Champollion, N., Dusch, M., Eis, J., Fourteau, K., Gregor, P., Jarosch, A. H., Landmann, J., Oesterle, F., Recinos, B., Rothenpieler, T., Vlug, A., Wild, C. T., & Marzeion, B. (2019). The Open Global Glacier Model (OGGM) v1.1. *Geoscientific Model Development*, 12(3), 909–931. <https://doi.org/10.5194/gmd-12-909-2019>
 220
 221
 222
- 223 Millan, R., Mouginit, J., Rabatel, A., & Morlighem, M. (2022). Ice velocity and thickness of the world's glaciers. *Nature Geoscience*, 15(2), 124–129. <https://doi.org/10.1038/s41561-021-00885-z>
 224
 225
- 226 Minchew, B., & Joughin, I. (2020). Toward a universal glacier slip law. *Science*, 368(6486), 29–30. <https://doi.org/10.1126/science.abb3566>
 227
- 228 Moses, W. S., Cheng, G., Churavy, V., Gelbrecht, M., Klöwer, M., Kump, J., Morlighem, M., Williamson, S., Apte, D., Berg, P., Giordano, M., Hill, C., Loose, N., Montoisson, A., Narayanan, S. H. K., Pal, A., Schanen, M., Silvestri, S., Wagner, G. L., & Heimbach, P. (n.d.). *DJ4Earth: Differentiable, and Performance-portable Earth System Modeling via Program Transformations*. Retrieved November 19, 2025, from <https://www.authorea.com/doi/full/10.22541/essoar.176314951.18114616?commit=b41d12401ff9fbfce8bb85c4a3c9cd8681e142a0>
 229
 230
 231
 232
 233
 234
- 235 Moses, W. S., Churavy, V., Paehler, L., Hükelheim, J., Narayanan, S. H. K., Schanen, M., & Doerfert, J. (2021). Reverse-mode automatic differentiation and optimization of GPU kernels via enzyme. *Proceedings of the International Conference for High Performance Computing, Networking, Storage and Analysis*. <https://doi.org/10.1145/3458817.3476165>
 236
 237
 238
- 239 Nyenah, E., Döll, P., Katz, D. S., & Reinecke, R. (2024). Software sustainability of global impact models. *Geoscientific Model Development Discussions*, 2024, 1–29. <https://doi.org/10.5194/gmd-2024-97>
 240
 241
- 242 Rackauckas, C., Innes, M., Ma, Y., Bettencourt, J., White, L., & Dixit, V. (2019). DiffEqFlux.jl - A Julia Library for Neural Differential Equations. *arXiv:1902.02376 [Cs, Stat]*. <http://arxiv.org/abs/1902.02376>
 243
 244
- 245 Rackauckas, C., Ma, Y., Martensen, J., Warner, C., Zubov, K., Supekar, R., Skinner, D., Ramadhan, A., & Edelman, A. (2021). *Universal Differential Equations for Scientific Machine Learning*. arXiv. <https://doi.org/10.48550/arXiv.2001.04385>
 246
 247
- 248 Rackauckas, C., & Nie, Q. (2017). DifferentialEquations.jl – A Performant and Feature-Rich Ecosystem for Solving Differential Equations in Julia. *Journal of Open Research Software*,
 249

- 250 5, 15. <https://doi.org/10.5334/jors.151>
- 251 RGI Consortium. (2023). *Randolph Glacier Inventory - A Dataset of Global Glacier Outlines,*
252 *Version 7.* National Snow; Ice Data Center. <https://doi.org/10.5067/F6JMOVY5NAVZ>
- 253 Rounce, D. R., Hock, R., Maussion, F., Hugonnet, R., Kochtitzky, W., Huss, M., Berthier,
254 E., Brinkerhoff, D., Compagno, L., Copland, L., Farinotti, D., Menounos, B., & McNabb,
255 R. W. (2023). Global glacier change in the 21st century: Every increase in temperature
256 matters. *Science*, 379(6627), 78–83. <https://doi.org/10.1126/science.abo1324>
- 257 Sapienza, F., Bolibar, J., Schäfer, F., Groenke, B., Pal, A., Boussange, V., Heimbach,
258 P., Hooker, G., Pérez, F., Persson, P.-O., & Rackauckas, C. (2024). *Differentiable*
259 *Programming for Differential Equations: A Review.* arXiv. [http://arxiv.org/abs/2406.](http://arxiv.org/abs/2406.09699)
260 [09699](http://arxiv.org/abs/2406.09699)
- 261 Shen, C., Appling, A. P., Gentine, P., Bandai, T., Gupta, H., Tartakovsky, A., Baity-Jesi,
262 M., Fenicia, F., Kifer, D., Li, L., Liu, X., Ren, W., Zheng, Y., Harman, C. J., Clark, M.,
263 Farthing, M., Feng, D., Kumar, P., Aboelyazeed, D., ... Lawson, K. (2023). Differentiable
264 modelling to unify machine learning and physical models for geosciences. *Nature Reviews*
265 *Earth & Environment*, 1–16. <https://doi.org/10.1038/s43017-023-00450-9>
- 266 Sjursen, K. H., Bolibar, J., Meer, M. van der, Andreassen, L. M., Biesheuvel, J. P., Dunse,
267 T., Huss, M., Maussion, F., Rounce, D. R., & Tober, B. (2025). Machine learning
268 improves seasonal mass balance prediction for unmonitored glaciers. *EGUsphere*, 1–39.
269 <https://doi.org/10.5194/egusphere-2025-1206>
- 270 Weis, M., Greve, R., & Hutter, K. (1999). Theory of shallow ice shelves. *Continuum Mechanics*
271 *and Thermodynamics*, 11(1), 15–50. <https://doi.org/10.1007/s001610050102>