The Journal of Open Source Software

pide: Petrophysical Interpretation tools for geoDynamic Exploration.

Sinan Ozaydin [1*¶], **Lu Li** [2*], **Utpal Singh** [1], **Patrice F. Rey** [1], and **Maria Constanza Manassero** [3]

1 School of Geosciences, University of Sydney, Sydney, Australia. **2** School of Earth Sciences, University of Western Australia, Perth, Australia. **3** School of Natural Sciences (Physics), University of Tasmania, Hobart, Australia. ¶ Corresponding author * These authors contributed equally.

DOI: 10.xxxxxx/draft

Software

- Review
- Repository
- Archive

Editor:

Submitted: 10 May 2024
Published: unpublished

License
Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License (CC BY 4.0).

DRAFT

Summary

`pide` is a Python3 library for calculating geophysical parameters (e.g., electrical conductivity, seismic velocity), employing the results from experimental petrology, mineral/rock physics, and thermomechanical modelling studies. `pide` can calculate the theoretical electrical conductivity of any earth material that exists in the literature. `pide` can also calculate seismic velocity utilising the external 'sister' library `SAnTex`. Using these theoretical calculations, users can utilise inversion modules to decode geophysical anomalies compositionally or convert thermomechanical models into geophysical observables. With a given spatial mapping of earth materials, which can preferentially be loaded from a thermomechanical model, `pide` can be used to build synthetic electrical conductivity and seismic velocity models and generate gravity and magnetic anomalies. Moreover, it is built as a modular tool, so users can easily build their functions.

Statement of need

Given the inherent heterogeneity and complexity of Earth systems, geophysical tomographies often yield complex 2D and 3D images that are challenging to interpret. To enhance their interpretations, researchers commonly turn to experimental petrology and mineral physics, covering various geophysical properties, including electrical conductivity, magnetic susceptibility, seismic velocity, and rheology. These properties are sensitive to phase transitions, partial melting, major and trace elements partitioning, mineral solubilities, and phase-mixing models. Numerous specialized tools have been designed to address specific properties, many of which feature graphical interfaces (Abers & Hacker, 2016; Özaydın & Selway, 2020) or are accessible through web-based applications (Pommier & Le-Trong, 2011). In this context, `pide` is a solution fulfilling the need for a versatile library capable of facilitating petrophysical calculations across a range of properties and supporting the creation of specific scientific tools. Beyond this, `pide` aims to host toolkits tailored for specific purposes, such as constructing realistic, petrophysically constrained synthetic models and converting numerical plate tectonic models into synthetic geophysical tomographies.

Library modules and methods

The general workflow diagram of the library can ben seen in Figure 1. The library has three main classes used in these calculations: `pide`, `material` and `model`.

`pide` is the main class in which the electrical conductivity and seismic-related observables

are calculated. In order to achieve this, the relevant parameters have to be defined in the pide object with the associated functions (e.g., composition, water content, interconnectivity). Just using the `pide` class, for instance, the user can make a figure of all calculations of all the olivine electrical conductivity and seismic velocity models for olivine. While experimental parameters for electrical conductivity parameters are defined and calculated within the `pide`, seismic velocities of given earth materials are calculated through `SAnTex` library automatically.

`material` is the class that can be specified as a holder of pre-defined material properties. For instance, one can create a Lherzolite material by mixing specific modal proportions of ol, opx, cpx and gt, how these constituents are interconnected, or how water behave among them. This class allows the user to define some parameters tied to other parameters via user-defined functions. This allows the users to streamline some specific processes in a modular manner.

The `model` class, on the other hand, is where a collection of `material` objects can be appended with specific positions indexed in 3D space. `model` is also where the user can calculate the magnetic and gravitational anomalies solely since these observables are dependent on the position of the materials and assigned magnetic and density parameters only. `pide` can generate synthetic data for magnetic and gravitational anomalies utilising the `harmonica` library (Soler et al., 2019). However, it can only generate synthetic magnetotelluric and seismic velocity models since their forward modelling is not as easy to implement within the pide framework. Currently, pide can generate input files for commonly used magnetotelluric modelling algorithms `ModEM` (Kelbert et al., 2014) and `Mare2DEM` (Key, 2016), and then users can generate synthetic data using the algorithms themselves with the given synthetic model.

`pide` also comes with several tools that can exploit the program architecthure. 'model_modifier' functions, for instance, can be used to calculate certain specific hypothesis tests. `inversion` modules, on the other hand, can be utilised to invert for specific input parameters that fit real geophysical data.



Figure 1: Workflow Chart for pide

Acknowledgements

This study is supported by the Australian Research Council (ARC) Linkage Grant ARC-LP190100146 and ARC DP Grant ARC-DP220100709.

References

Abers, G. A., & Hacker, B. R. (2016). A MATLAB toolbox and excel workbook for calculating the densities, seismic wave speeds, and major element composition of minerals and rocks at pressure and temperature. *Geochemistry, Geophysics, Geosystems*, *17*(2), 616–624. https://doi.org/10.1002/2015GC006171

Kelbert, A., Meqbel, N., Egbert, G. D., & Tandon, K. (2014). ModEM: A modular system for inversion of electromagnetic geophysical data. *Computers & Geosciences*, *66*, 40–53. https://doi.org/10.1016/j.cageo.2014.01.010

Key, K. (2016). MARE2DEM: A 2-d inversion code for controlled-source electromagnetic and magnetotelluric data. *Geophysical Journal International*, *207*(1), 571–588. https://doi.org/10.1093/gji/ggw290

Özaydın, S., & Selway, K. (2020). MATE: An analysis tool for the interpretation of magnetotelluric models of the mantle. *Geochemistry, Geophysics, Geosystems*, *21*(9), e2020GC009126. https://doi.org/10.1029/2020gc009126

Pommier, A., & Le-Trong, E. (2011). "SIGMELTS": A web portal for electrical conductivity calculations in geosciences. *Computers & Geosciences*, *37*(9), 1450–1459. https://doi.org/10.1016/j.cageo.2011.01.002

Soler, S. R., Pesce, A., Gimenez, M. E., & Uieda, L. (2019). Gravitational field calculation in spherical coordinates using variable densities in depth. *Geophysical Journal International*, *218*(3), 2150–2164. https://doi.org/10.1093/gji/ggz277

DRAFT