

HBAT 2: A Python Package to Analyse Hydrogen Bonds and Other Non-covalent Interactions in Macromolecular Structures

Abhishek Tiwari ¹

1 Independent Researcher, New York, United States

DOI: 10.xxxxxx/draft

Software

- [Review](#)
- [Repository](#)
- [Archive](#)

Editor: [✉](#)

Submitted: 15 May 2026

Published: unpublished

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

Summary

HBAT 2 is a Python package for automated analysis of hydrogen bonds and other non-covalent interactions in macromolecular structures available in Protein Data Bank (PDB) format. The software identifies and analyzes traditional hydrogen bonds, weak hydrogen bonds, halogen bonds, X-H... π interactions, π - π stacking, and $n \rightarrow \pi^*$ interactions using geometric criteria. HBAT 2 also detects networks of linked hydrogen bonds and renders them as interactive 2D network visualizations. Originally developed in Perl/Tk and published in 2007 (Tiwari & Panigrahi, 2007), HBAT 2 has been completely rewritten in Python with a modern cross-platform tkinter-based graphical user interface (GUI), web-based interface, command-line interface (CLI), and developer-friendly APIs enabling integration with Jupyter notebooks and custom workflows. The software also provides JavaScript-based 3D visualization available both on the web server and as a widget in Jupyter notebooks, enabling researchers to interactively explore detected interactions without requiring separate visualization software.




Figure 1: The latest update to HBAT 2 uses tkinter to provide a cross-platform graphical user interface (GUI)

Statement of Need

Hydrogen bonds (Desiraju & Steiner, 2001) and other non-covalent interactions, including halogen bonds (Cavallo et al., 2016), are fundamental to the structure, stability, and function of macromolecular systems including proteins, nucleic acids, and their complexes. These interactions determine how biological molecules fold, recognize their targets, and bind ligands. With over 200,000 macromolecular structures archived in the Protein Data Bank (Berman et al., 2000), there is an increasing need for automated tools to systematically analyze these critical interactions across diverse structural types.

The target audience includes structural biologists studying molecular mechanisms, computational chemists designing new drugs, protein engineers improving enzyme properties, and researchers analyzing molecular dynamics simulations. These researchers need to quickly identify and quantify weak interactions in diverse macromolecular structures to answer biological questions about molecular function and evolution.

The original HBAT (Tiwari & Panigrahi, 2007) was developed in Perl/Tk with a Windows-only GUI, limiting its adoption in modern computational environments where researchers use diverse operating systems (Windows, Linux, macOS) and integrate analyses into Python-based scientific workflows. This update addresses these limitations while expanding the types of interactions analyzed.

State of the Field

The landscape of hydrogen bond analysis tools is diverse but fragmented, with most solutions designed for specific use cases or as secondary features within larger systems. Classic tools like HBPLUS (McDonald & Thornton, 1994) and HBExplore (Lindauer et al., 1996) pioneered automated H-bond detection but lack modern interfaces and support for diverse interaction types.

Current tools are typically constrained: PLIP (Salentin et al., 2015) and Arpeggio (Jubb et al., 2017) focus on protein-ligand interactions; HBonanza (Durrant & McCammon, 2011), HBCalculator (Y. Wang & Wang, 2024), and BRIDGE2 (Siemers & Bondar, 2021) are optimized for molecular dynamics trajectories; MDAnalysis (MDAnalysis, 2025), GROMACS (GROMACS, 2025), and AMBER (AMBER, 2025) provide H-bond analysis as secondary features within larger simulation suites. Visualization tools like VMD (VMD, 2025) and ChimeraX (ChimeraX, 2025) excel at interactive exploration but offer limited quantitative analysis. ProteinTools (Ferruz et al., 2021) provides web-based network analysis but lacks cross-platform accessibility.

HBAT 2 uniquely provides a **general-purpose solution** with: (1) detection of diverse interaction types including hydrogen bonds, halogen bonds, π -interactions, and $n \rightarrow \pi^*$ interactions; (2) multiple access modes (desktop GUI, command-line, web server, Python API) supporting diverse research workflows; (3) identification of hydrogen bond networks - both potential cooperativity and anticooperativity chains and water-mediated hydrogen bond networks - in static structures; (4) seamless Jupyter (Thomas et al., 2016) integration; and (5) flexible parameters with domain-specific presets. The modern Python implementation with cross-platform support and web accessibility makes comprehensive interaction analysis available to researchers regardless of computational background or infrastructure.

Software Design

HBAT 2 employs a modular architecture with distinct components for PDB parsing, geometric analysis, statistical computation, and network visualization. The core analysis engine uses efficient nearest-neighbor searching with configurable distance cutoffs, followed by geometric filtering based on distance and angular criteria. This fundamental approach matches the

65 original HBAT (Tiwari & Panigrahi, 2007) but with optimized algorithms for analyzing large
66 macromolecular structures.




Figure 2: HBAT 2 analysis workflow showing the complete pipeline from PDB input to multiple output formats.

67 Key design decisions include: (1) Python for accessibility and integration with scientific
68 ecosystems; (2) integration with PDBFixer (Eastman et al., 2013; Openmm/Pdbfixer, 2025)
69 and OpenBabel (O'Boyle et al., 2008) for automated structure preparation, addressing real-world
70 challenges of missing atoms and incomplete side chains; (3) support for multiple interaction
71 types reflecting advances in weak interaction chemistry (Brandl et al., 2001; Cavallo et al., 2016;
72 Desiraju & Steiner, 2001); (4) preset parameter systems for common experimental conditions
73 (high-resolution X-ray, NMR, membrane proteins, drug design) that balance accessibility for
74 experimentalists with flexibility for computational experts; and (5) multiple output formats
75 (CSV, JSON, network graphics in PNG/SVG/PDF) enabling integration into downstream

76 analysis pipelines.

77 The software analyzes hydrogen bonds (O-H...O, N-H...O, N-H...N, C-H...O), halogen bonds
78 (C-X...Y where X=F,Cl,Br,I), X-H... π interactions, π - π stacking (McGaughey et al., 1998;
79 Vernon et al., n.d.), carbonyl-carbonyl n $\rightarrow$ π^* interactions (Newberry & Raines, 2017; Rahim
80 et al., 2017), and n $\rightarrow$ π^* interactions (Choudhary et al., 2009). Network visualization uses
81 either NetworkX/Matplotlib (Hagberg et al., 2008; Hunter, 2007) or GraphViz (Graphviz,
82 2025), emphasizing cooperativity chains and network topology with customizable layouts and
83 high-resolution export.




Figure 3: C-H... π interaction between A:ALA:20 $\rightarrow$ A:TYR:25 of PDB Structure 6RSA visualized using 3Dmol.js integration in HBAT 2

84 3D Visualization with 3Dmol.js Integration

85 HBAT 2 integrates 3Dmol.js (Rego & Koes, 2014) for interactive 3D molecular visualization in
86 both web and Jupyter notebook environments. The JavaScript-based viewer enables researchers
87 to interactively explore detected interactions with customizable color schemes, allowing rotation,
88 zoom, and inspection of specific interaction geometries. Within Jupyter notebooks, the 3D
89 visualization widget is embedded directly alongside analysis code, bridging automated analysis
90 and manual structural inspection.

91 Research Impact Statement

92 Since its original publication, HBAT has been cited more than 80 times in numerous studies
93 spanning multiple research domains (Tiwari & Panigrahi, 2007). HBAT has been used for:

94 **Structure-based drug design:** Researchers have used HBAT to characterize protein-ligand
95 interactions for anticancer compounds targeting ABL kinase (Díaz-Cervantes et al., 2020),
96 analyze cardiovascular disease receptors with metabolite interactions (Ravindran et al., 2015),

97 predict drug resistance based on hydrogen bond patterns (Zhou et al., 2013), and elucidate
98 anticoagulant interactions (Russo Krauss et al., 2011).

99 **Protein engineering:** HBAT enabled identification of stabilizing C-H... π and C-H...O interactions
100 in thermostable enzyme variants (C.-H. Wang et al., 2020) and characterization of hydrogen
101 bonding networks for functional specificity (Fournier et al., 2014).

102 **Molecular dynamics analysis:** The tool has been used to characterize interactions in viral
103 proteins important for understanding RNA encapsulation (Huang et al., 2009), glycoprotein
104 interactions for cellular quality control (Jayaprakash et al., 2025), and DNA structural dynamics
105 (Barzegar & Tohidifar, 2025).

106 **Comparative structural analysis:** HBAT has enabled systematic mutation analysis identifying
107 disease-associated variants in cancer genes (Abdulazeez et al., 2019; Abdulazeez & Borgio,
108 2016; Kavitha & Mohanapriya, 2024; Khan et al., 2018; Khan & Ansari, 2017).

109 **Recent citations:** Recent applications include biosorption and molecular docking studies utilizing
110 HBAT 2 for hydrogen bond analysis (Er et al., 2026; Karadayı et al., 2026).




Figure 4: An example visualization of hydrogen-bonds network detected by HBAT 2 software for Protein Data Bank (PDB) entry 6RSA

111 HBAT 2 extends this demonstrated impact by providing modern cross-platform access, expanded
112 interaction coverage, and integration with contemporary Python workflows, supporting growing
113 adoption across diverse research applications.

114 Availability

115 HBAT 2 is freely available to download from [GitHub](#) and [PyPI](#) under the MIT license with
116 detailed [user and API documentation](#). The software can be installed via [PyPI](#) (`pip install hbat`) or via [Conda](#) (`conda install -c hbat hbat`), with optional GraphViz integration for
117 advanced visualization features. A [web-based interface](#) is available for immediate browser-based
118 analysis without installation. Interactive [Jupyter notebooks](#) with Google Colab integration are
119 provided for computational workflows.
120

AI Usage Disclosure

AI tools were used for porting original HBA1 from Perl to Python, as well as for refactoring, test scaffolding, and documentation generation for portions of the codebase. AI tools were also used to assist with editing portions of this manuscript. All AI-assisted code and documentation outputs were reviewed, edited, validated, and tested by the human author, who takes full responsibility for the final software and paper. No figures or data were generated by AI.

Acknowledgements

The author thanks the original co-developer Sunil K. Panigrahi and acknowledges the structural biology community for feedback that guided the modernization of HBA1.

References

- AbdulAzeez, S., & Borgio, J. F. (2016). In-Silico Computing of the Most Deleterious nsSNPs in HBA1 Gene. *PLOS ONE*, 11(1), e0147702. <https://doi.org/10.1371/journal.pone.0147702>
- Abdulazeed, S., Sultana, S., Almandil, N. B., Almohazey, D., Bency, B. J., & Borgio, J. F. (2019). The rs61742690 (S783N) single nucleotide polymorphism is a suitable target for disrupting BCL11A-mediated foetal-to-adult globin switching. *PLOS ONE*, 14(2), e0212492. <https://doi.org/10.1371/journal.pone.0212492>
- AMBER. (2025). <https://amberhub.chpc.utah.edu/hbond/>
- Barzegar, A., & Tohidifar, N. (2025). Stabilization of Hoogsteen H-bonds in G-quartet sheets by coordinated K⁺ ion for enhanced efficiency in guanine-rich DNA nanomotor. *BiolImpacts : BI*, 15, 30596. <https://doi.org/10.34172/bi.30596>
- Berman, H. M., Westbrook, J., Feng, Z., Gilliland, G., Bhat, T. N., Weissig, H., Shindyalov, I. N., & Bourne, P. E. (2000). The protein data bank. *Nucleic Acids Research*, 28(1), 235–242.
- Brandl, M., Weiss, M. S., Jabs, A., Sühnel, J., & Hilgenfeld, R. (2001). C-h π -interactions in proteins. *Journal of Molecular Biology*, 307(1), 357–377. <https://doi.org/10.1006/jmbi.2000.4473>
- Cavallo, G., Metrangolo, P., Milani, R., Pilati, T., Priimagi, A., Resnati, G., & Terraneo, G. (2016). The Halogen Bond. *Chemical Reviews*, 116(4), 2478–2601. <https://doi.org/10.1021/acs.chemrev.5b00484>
- Chimerax. (2025). <https://www.cgl.ucsf.edu/chimerax/docs/user/tools/hbonds.html>
- Choudhary, A., Gandla, D., Krow, G. R., & Raines, R. T. (2009). Nature of Amide Carbonyl-Carbonyl Interactions in Proteins. *Journal of the American Chemical Society*, 131(21), 7244–7246. <https://doi.org/10.1021/ja901188y>
- Desiraju, G., & Steiner, T. (2001). *The Weak Hydrogen Bond*. Oxford University Press. <https://doi.org/10.1093/acprof:oso/9780198509707.001.0001>
- Díaz-Cervantes, E., Cortés-García, C. J., Chacón-García, L., & Suárez-Castro, A. (2020). Molecular docking and pharmacophoric modelling of 1,5-disubstituted tetrazoles as inhibitors of two proteins present in cancer, the ABL and the mutated T315I kinase. *In Silico Pharmacology*, 8(1), 6. <https://doi.org/10.1007/s40203-020-00059-6>
- Durrant, J. D., & McCammon, J. A. (2011). HBonanza: A Computer Algorithm for Molecular-Dynamics-Trajectory Hydrogen-Bond Analysis. *Journal of Molecular Graphics & Modelling*, 31, 5–9. <https://doi.org/10.1016/j.jmgm.2011.07.008>

- 163 Eastman, P., Friedrichs, M. S., Chodera, J. D., Radmer, R. J., Bruns, C. M., Ku, J. P.,
164 Beauchamp, K. A., Lane, T. J., Wang, L.-P., Shukla, D., Tye, T., Houston, M., Stich,
165 T., Klein, C., Shirts, M. R., & Pande, V. S. (2013). OpenMM 4: A Reusable, Extensible,
166 Hardware Independent Library for High Performance Molecular Simulation. *Journal of*
167 *Chemical Theory and Computation*, 9(1), 461–469. <https://doi.org/10.1021/ct300857j>
- 168 Er, M. Y., Karadayi, G., Gulsahin, Y., Karadayi, M., Turan, M., Yildirim, E., Agar, G., &
169 Gulluce, M. (2026). Mitigation of bensulfuron-methyl induced multiple stress responses in
170 cucumis sativus l. By biostimulants: A comprehensive metabolic and molecular docking
171 approach. *Journal of Hazardous Materials*, 514, 142962. <https://doi.org/10.1016/j.jhazmat.2026.142962>
- 173 Ferruz, N., Schmidt, S., & Höcker, B. (2021). ProteinTools: A toolkit to analyze protein
174 structures. *Nucleic Acids Research*, 49(W1), W559–W566. [https://doi.org/10.1093/nar/](https://doi.org/10.1093/nar/gkab375)
175 [gkab375](https://doi.org/10.1093/nar/gkab375)
- 176 Fournier, J.-B., Rebuffet, E., Delage, L., Grijol, R., Meslet-Cladière, L., Rzonca, J., Potin, P.,
177 Michel, G., Czjzek, M., & Leblanc, C. (2014). The Vanadium Iodoperoxidase from the
178 Marine Flavobacteriaceae Species *Zobellia galactanivorans* Reveals Novel Molecular and
179 Evolutionary Features of Halide Specificity in the Vanadium Haloperoxidase Enzyme Family.
180 *Applied and Environmental Microbiology*, 80(24), 7561–7573. [https://doi.org/10.1128/](https://doi.org/10.1128/AEM.02430-14)
181 [AEM.02430-14](https://doi.org/10.1128/AEM.02430-14)
- 182 *Graphviz*. (2025). <https://graphviz.org/>.
- 183 *GROMACS*. (2025). <https://manual.gromacs.org/current/onlinehelp/gmx-hbond.html>
- 184 Hagberg, A., Swart, P., & Chult, D. S. (2008). Exploring network structure, dynamics, and
185 function using NetworkX. *Proceedings of the 7th Python in Science Conference*, 11, 11–15.
186 <https://doi.org/10.25080/tcww9851>
- 187 Huang, L., Massa, L., & Karle, J. (2009). Kernel energy method applied to vesicular stomatitis
188 virus nucleoprotein. *Proceedings of the National Academy of Sciences*, 106(6), 1731–1736.
189 <https://doi.org/10.1073/pnas.0811959106>
- 190 Hunter, J. D. (2007). Matplotlib: A 2D graphics environment. *Computing in Science &*
191 *Engineering*, 9(3), 90–95. <https://doi.org/10.1109/MCSE.2007.55>
- 192 Jayaprakash, N. G., Sarkar, D. K., & Surolia, A. (2025). Atomic visualization of flipped-
193 back conformations of high mannose glycans interacting with cargo lectins: An MD
194 simulation perspective. *Proteins: Structure, Function, and Bioinformatics*, 93(1), 112–133.
195 <https://doi.org/10.1002/prot.26556>
- 196 Jubb, H. C., Higuero, A. P., Ochoa-Montano, B., Pitt, W. R., Ascher, D. B., & Blundell,
197 T. L. (2017). Arpeggio: A Web Server for Calculating and Visualising Interatomic
198 Interactions in Protein Structures. *Journal of Molecular Biology*, 429(3), 365–371. <https://doi.org/10.1016/j.jmb.2016.12.004>
- 199 Karadayi, M., Güllüce, E., Aksu, Ş., Gülşahin, Y., Tosun, B., Karadayi, G., & Güllüce, M.
200 (2026). Removal and detoxification of safranin in aqueous solutions using a *betula pendula*
201 roth based biomass: Isotherm, kinetics, thermodynamics of the biosorption process, and
202 molecular insights into the mechanism of safranin ecotoxicity. *Water Environment Research*,
203 98(5), e70393. <https://doi.org/10.1002/wer.70393>
- 205 Kavitha, K., & Mohanapriya, A. (2024). Insights into the structure–function relationship of
206 missense mutations in the human TOP2A protein in ovarian cancer. *Frontiers in Physics*,
207 12. <https://doi.org/10.3389/fphy.2024.1358406>
- 208 Khan, I., & Ansari, I. A. (2017). Prediction of a highly deleterious mutation E17K in
209 AKT-1 gene: An *in silico* approach. *Biochemistry and Biophysics Reports*, 10, 260–266.
210 <https://doi.org/10.1016/j.bbrep.2017.04.013>

- 211 Khan, I., Ansari, I. A., Singh, P., Dass, J. F. P., & Khan, F. (2018). Identification and
212 characterization of functional single nucleotide polymorphisms (SNPs) in Axin 1 gene:
213 A molecular dynamics approach. *Cell Biochemistry and Biophysics*, 76(1), 173–185.
214 <https://doi.org/10.1007/s12013-017-0818-1>
- 215 Lindauer, K., Bendic, C., & Sühnel, J. (1996). HBExplore—a new tool for identifying and
216 analysing hydrogen bonding patterns in biological macromolecules. *Computer Applications*
217 *in the Biosciences: CABIOS*, 12(4), 281–289. [https://doi.org/10.1093/bioinformatics/12.](https://doi.org/10.1093/bioinformatics/12.4.281)
218 [4.281](https://doi.org/10.1093/bioinformatics/12.4.281)
- 219 McDonald, I. K., & Thornton, J. M. (1994). Satisfying hydrogen bonding potential in proteins.
220 *Journal of Molecular Biology*, 238(5), 777–793. <https://doi.org/10.1006/jmbi.1994.1334>
- 221 McGaughey, G. B., Gagné, M., & Rappé, A. K. (1998). π -Stacking Interactions: ALIVE
222 AND WELL IN PROTEINS*. *Journal of Biological Chemistry*, 273(25), 15458–15463.
223 <https://doi.org/10.1074/jbc.273.25.15458>
- 224 MDAnalysis. (2025). [https://docs.mdanalysis.org/1.1.0/documentation_pages/analysis/](https://docs.mdanalysis.org/1.1.0/documentation_pages/analysis/hydrogenbonds.html)
225 [hydrogenbonds.html](https://docs.mdanalysis.org/1.1.0/documentation_pages/analysis/hydrogenbonds.html)
- 226 Newberry, R. W., & Raines, R. T. (2017). The $n \rightarrow \pi^*$ Interaction. *Accounts of Chemical*
227 *Research*, 50(8), 1838–1846. <https://doi.org/10.1021/acs.accounts.7b00121>
- 228 O'Boyle, N. M., Morley, C., & Hutchison, G. R. (2008). Pybel: A Python wrapper for
229 the OpenBabel cheminformatics toolkit. *Chemistry Central Journal*, 2(1), 5. <https://doi.org/10.1186/1752-153X-2-5>
- 230 <https://doi.org/10.1186/1752-153X-2-5>
- 231 Openmm/pdbfixer. (2025). OpenMM. <https://github.com/openmm/pdbfixer>
- 232 Rahim, A., Saha, P., Jha, K. K., Sukumar, N., & Sarma, B. K. (2017). Reciprocal
233 carbonyl–carbonyl interactions in small molecules and proteins. *Nature Communications*,
234 8(1), 78. <https://doi.org/10.1038/s41467-017-00081-x>
- 235 Ravindran, R., Sharma, N., Roy, S., Thakur, A. R., Ganesh, S., Kumar, S., Devi, J., & Rajkumar,
236 J. (2015). Interaction Studies of Withania Somnifera's Key Metabolite Withaferin A with
237 Different Receptors Associated with Cardiovascular Disease. *Current Computer-Aided*
238 *Drug Design*, 11(3), 212–221. <https://doi.org/10.2174/1573409912666151106115848>
- 239 Rego, N., & Koes, D. (2014). 3Dmol.js: Molecular visualization with WebGL. *Bioinformatics*,
240 31(8), 1322–1324. <https://doi.org/10.1093/bioinformatics/btu829>
- 241 Russo Krauss, I., Merlino, A., Giancola, C., Randazzo, A., Mazzarella, L., & Sica, F. (2011).
242 Thrombin–aptamer recognition: A revealed ambiguity. *Nucleic Acids Research*, 39(17),
243 7858–7867. <https://doi.org/10.1093/nar/gkr522>
- 244 Salentin, S., Schreiber, S., Haupt, V. J., Adasme, M. F., & Schroeder, M. (2015). PLIP:
245 Fully automated protein–ligand interaction profiler. *Nucleic Acids Research*, 43(W1),
246 W443–W447. <https://doi.org/10.1093/nar/gkv315>
- 247 Siemers, M., & Bondar, A.-N. (2021). Interactive Interface for Graph-Based Analyses of
248 Dynamic H-Bond Networks: Application to Spike Protein S. *Journal of Chemical Information*
249 *and Modeling*, 61(6), 2998–3014. <https://doi.org/10.1021/acs.jcim.1c00306>
- 250 Thomas, K., Benjamin, R.-K., Fernando, P., Brian, G., Matthias, B., Jonathan, F., Kyle, K.,
251 Jessica, H., Jason, G., Sylvain, C., Paul, I., Damián, A., Safia, A., Carol, W., & Jupyter
252 Development Team. (2016). Jupyter notebooks – a publishing format for reproducible
253 computational workflows. In *Positioning and power in academic publishing: Players, agents*
254 *and agendas*. IOS Press.
- 255 Tiwari, A., & Panigrahi, S. K. (2007). HBAT: A complete package for analysing strong
256 and weak hydrogen bonds in macromolecular crystal structures. *In Silico Biology*, 7(6),
257 651–661. <https://doi.org/10.3233/ISI-2007-00337>

- 258 Vernon, R. M., Chong, P. A., Tsang, B., Kim, T. H., Bah, A., Farber, P., Lin, H., & Forman-
259 Kay, J. D. (n.d.). Pi-Pi contacts are an overlooked protein feature relevant to phase
260 separation. *eLife*, 7, e31486. <https://doi.org/10.7554/eLife.31486>
- 261 VMD. (2025). <https://www.ks.uiuc.edu/Research/vmd/plugins/hbonds/>
- 262 Wang, C.-H., Lu, L.-H., Huang, C., He, B.-F., & Huang, R.-B. (2020). Simultaneously
263 Improved Thermostability and Hydrolytic Pattern of Alpha-Amylase by Engineering Central
264 Beta Strands of TIM Barrel. *Applied Biochemistry and Biotechnology*, 192(1), 57–70.
265 <https://doi.org/10.1007/s12010-020-03308-8>
- 266 Wang, Y., & Wang, Y. (2024). HBCalculator: A Tool for Hydrogen Bond Distribution
267 Calculations in Molecular Dynamics Simulations. *Journal of Chemical Information and*
268 *Modeling*, 64(6), 1772–1777. <https://doi.org/10.1021/acs.jcim.4c00054>
- 269 Zhou, W., Wang, D. D., Yan, H., Wong, M., & Lee, V. (2013). Prediction of anti-EGFR drug
270 resistance base on binding free energy and hydrogen bond analysis. *2013 IEEE Symposium*
271 *on Computational Intelligence in Bioinformatics and Computational Biology (CIBCB)*,
272 193–197. <https://doi.org/10.1109/CIBCB.2013.6595408>

DRAFT