

FEZrs: An Open-Source Python Package for Geospatial Multispectral Image Processing and Remote Sensing

Mahdi Farmahinifarahani¹, Parsa Elmi^{2¶}, Mehdi Nedaei³, Mohammad Kiani faizabadi⁴, Hooman Mirzaee⁴, Parsa Moradi³, Mohammadhossein Yazdanifar⁵, Fariba Khosravani⁶, Fatemeh Najafi⁶, and Mehdi Talkhablou⁶

¹ Department of Earth and Environmental Science, The Chinese University of Hong Kong, Hong Kong
² SAR 2 Faculty of Management, Islamic Azad University, Central Tehran Branch, Tehran, Iran
³ Department of Software Engineering, Shamsipour Technical College, Tehran, Iran
⁴ Faculty of Computer Science, Islamic Azad University, Central Tehran Branch, Tehran, Iran
⁵ Department of Industrial Engineering, Sharif University of Technology, Tehran, Iran
⁶ Faculty of Earth Sciences, Kharazmi University, Tehran, Iran
¶ Corresponding author

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#)
- [Repository](#)
- [Archive](#)

Editor: [Jack Atkinson](#)

Reviewers:

- [@gmarupilla](#)
- [@Digdgeo](#)
- [@martina01natali](#)

Submitted: 10 July 2026

Published: unpublished

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

Summary

The history of remote sensing has evolved from early nineteenth-century balloon photography to modern multispectral and satellite-based earth observation systems, enabling large-scale environmental monitoring and geospatial analysis workflows (Cracknell, 2018). Over time, remote sensing technologies have expanded from aerial observation platforms to advanced satellite (Zhang et al., 2022) and unmanned aerial vehicle (Simic Milas et al., 2018) systems that support scientific research, environmental assessment, and operational geospatial applications.

The increasing availability of multispectral remote sensing imagery from platforms such as Landsat (Williams et al., 2006) and Sentinel (Malenovsky et al., 2012) has accelerated the adoption of spectral analysis methods across a broad range of remote sensing tasks and applications (Montero et al., 2023; Xue & Su, 2017). These developments have expanded remote sensing use in geoscience (Camps-Valls et al., 2021) and mineral resource assessment (Sabins, 1999). The same imagery now also supports environmental engineering, wildlife monitoring (Melesse et al., 2007; Shukla et al., 2024), biology, marine science (Chassot et al., 2011), and agriculture (Sishodia et al., 2020).

As remote sensing datasets continue to grow in scale and complexity, researchers and practitioners increasingly rely on accessible computational tools for image analysis, feature extraction, and spectral interpretation. These workflows often require standardized implementations of widely used spectral indices (Montero et al., 2023) and consistent processing utilities to ensure reproducibility and efficiency in both research and operational applications. FEZrs is designed to simplify remote sensing data processing and interpretation by providing a consistent framework for a wide range of analytical workflows.

Although primarily developed for geoscience applications, FEZrs can also be applied more broadly to multispectral and hyperspectral spectral analysis in diverse fields beyond the geosciences.

Statement of Need

FEZrs is an open-source project for remote sensing image processing and geospatial analysis that provides a Python package for spectral analysis, image enhancement, filtering, spectral index computation, and feature extraction. The package integrates established remote sensing workflows within a unified open-source framework, supporting both research and industrial applications.




Figure 1: Overview of the FEZrs processing workflow. Multispectral remote sensing imagery is processed within the FEZrs framework to generate representative outputs, including (a) K-Means clustering, (b) gamma-corrected RGB enhancement, (c) RGB Image representation, and (d) Support Vector Machine (SVM) classification.

Modern remote sensing studies frequently rely on multiple analytical approaches. FEZrs implements a collection of widely used spectral indices (Montero et al., 2023), including the Normalized Difference Vegetation Index (NDVI) (Huang et al., 2021), Normalized Difference Water Index (NDWI) (McFeeters, 1996), Soil Adjusted Vegetation Index (SAVI) (Huete, 1988), aerosol free vegetation index (AFRI) (Karnieli et al., 2001), Bare Soil Index (BSI) (Rikimaru et al., 2002), and Urban Index (UI) (Deng & Wu, 2012). These indices are widely used for vegetation monitoring, water-resource assessment, environmental monitoring, land-surface characterization, and urban analysis.




Figure 2: Representative spectral index outputs generated by FEZrs, including (a) Soil Adjusted Vegetation Index (SAVI), (b) Urban Index (UI), (c) Normalized Difference Vegetation Index (NDVI), and (d) Normalized Difference Water Index (NDWI). All outputs were computed from the same multispectral dataset using the corresponding FEZrs implementations.

53 The package also incorporates image enhancement and filtering techniques, including Gaussian
54 smoothing (Lindeberg, 2024), median filtering (Justusson, 1981), Laplacian (Merris, 1994),
55 and Sobel operators (Chang et al., 2023), which are widely used for noise reduction, edge
56 detection, and image interpretation in remote sensing applications. Texture-based analysis
57 is supported through the Gray-Level Co-occurrence Matrix (GLCM) (Utaminingrum et al.,
58 2023), a widely adopted approach for characterizing spatial patterns in land-cover mapping
59 and environmental studies. FEZrs further provides Principal Component Analysis (PCA) for
60 dimensionality reduction and exploratory analysis of multispectral data (Jolliffe & Cadima,
61 2016), as well as K-Means clustering for unsupervised image classification and pattern discovery
62 (Sinaga & Yang, 2020). In addition, Support Vector Machine (SVM) classification is included as
63 a supervised machine-learning approach for land-cover classification and feature discrimination
64 in remote sensing datasets (Yue et al., 2003). spectral profile analysis enables the investigation
65 of spectral responses across wavelengths and supports multispectral interpretation workflows.

Change Detection	Filters	Image Enhancement	Spectral Indices	Additional Tools
Burn Indices	Gaussian Laplacian	Original Float	AFRI BI	KMeans GLCM

Change Detection	Filters	Image Enhancement	Spectral Indices	Additional Tools
MagDir	Mean	Equalize	NDVI	HSV
SubDiv	Median	Adaptive	NDWI	IRHSV
Time	Sobel	Gamma	SAVI	Mosaic
		LogAdjust	UI	PCA
		SigmoidAdjust		SpectralProfile
		OriginalRGB		SVM
		EqualizeRGB		
		AdaptiveRGB		
		GammaRGB		

Table 1: provides an overview of the implemented FEZrs modules, enabling users to perform common remote sensing processing tasks, spectral analysis, feature extraction, and image-based analysis workflows within a single Python package.

Although these methods are widely utilized, there is still considerable value in open-source tools that bring together essential remote sensing functionalities within a single, coherent analytical platform. While FEZrs was developed with geoscience applications in mind, its functionality is equally applicable to multispectral and hyperspectral data analysis across a variety of scientific and technical domains.

State of the field

FEZrs is built on general-purpose libraries (OpenCV (Xie & Lu, 2013), scikit-image (Walt et al., 2014), scikit-learn (Kramer, 2016), NumPy (Harris et al., 2020), matplotlib (Hunter, 2007), pandas (McKinney, 2010), Rasterio (Gillies & others, 2013), GDAL (Warmerdam, 2008), and GeoPandas (Jordahl et al., 2020)) rather than competing with them. The relevant comparison is with domain-level remote sensing packages that already implement overlapping methods.

The closest neighbour is scikit-eo (Tarazona et al., 2024), a JOSS package that covers spectral indices, PCA, supervised classification including SVM, data fusion, and accuracy assessment. spyindex and the Awesome Spectral Indices catalogue (Montero et al., 2023) standardise more than two hundred index definitions; FEZrs implements a small, documented subset of those indices rather than a catalogue. Ndv2Gif (García Díaz, 2026) provides multi-seasonal vegetation analysis on Google Earth Engine, a cloud time-series workflow that is complementary to FEZrs's local raster calculators. Other tools occupy adjacent parts of the same space: earthpy and EOReader for scene handling, rioxtarray for labelled rasters, RSGISLib and Orfeo Toolbox for processing pipelines, and torchgeo for deep-learning workflows.

Those packages are capable, and FEZrs is not a replacement for any of them. What it adds is a uniform calculator API: thirty-seven heterogeneous operations (indices, filters, enhancement, GLCM, PCA, clustering, change detection, SVM) behind one three-line calling pattern, each able to write a publication-ready figure via `execute()` and a georeferenced GeoTIFF via `to_raster()` without plotting or I/O code from the user. That uniformity is aimed at teaching and at rapid exploratory work on multispectral scenes, where assembling a bespoke toolchain for each step is the usual cost.

Package	Scope	What FEZrs adds
scikit-eo	Spectral indices, PCA, SVM, fusion, accuracy assessment	Same methods behind one calculator interface, plus publication figures

Package	Scope	What FEZrs adds
spyndex / Awesome Spectral Indices Ndv2Gif	Catalogue of 200+ index definitions Multi-seasonal vegetation analysis on Google Earth Engine	A small teaching set with <code>execute()</code> / <code>to_raster()</code> export Complementary cloud time-series workflows; FEZrs operates on local rasters
Rasterio, GDAL, OpenCV, scikit-image, scikit-learn	I/O, image processing, and ML building blocks	Domain workflows layered on top of these libraries
earthpy, EOReader, rioarray, RSGISLib, Orfeo Toolbox, torchgeo	Scene I/O, preprocessing pipelines, deep learning	Complementary; FEZrs does not replace them

Table 2: Domain-level packages with overlapping scope, and the narrower contribution FEZrs makes relative to each.

Software design

FEZrs is organized around a shared abstract base class (`BaseTool`) that standardizes the lifecycle of every analysis module: input validation, processing, visualization customization, and export. Each calculator (for example, `NDVICalculator`, `GaussianCalculator`, or `KMeansCalculator`) inherits this interface, accepts band file paths through a common file-handling layer, and exposes a small set of methods such as `execute` and `histogram_export`. This pattern keeps the public API uniform across spectral indices, filters, enhancement tools, change-detection utilities, and machine-learning modules, reducing cognitive overhead when composing multi-step workflows.

The package is modular by domain: (`spectral_indices`, `filters`, `image_enhancement`, `change_detection`, `clustering`, `glcm`, `pca`, `svm`, and others), while shared utilities handle band-path typing, file I/O, and histogram support. Raster inputs are typically multi-band geospatial imagery. Results can be exported two ways: as rendered figures via `execute()`, for inspection and reporting, and as georeferenced GeoTIFF rasters via `to_raster()`, which preserve full numerical precision along with the coordinate reference system and affine transform of the source imagery, and are therefore suitable for GIS overlay, zonal statistics, and multitemporal analysis. The design deliberately favors composition of independent calculators over a single monolithic pipeline object, so researchers can select only the methods required for a given study while still benefiting from consistent validation and export behavior.

Research Impact Statement

Since its release on PyPI in February 2025, FEZrs has received more than 100,000 downloads according to PePy, indicating substantial distribution of the package, with download statistics potentially including traffic from PyPI mirrors. FEZrs is openly available through GitHub, PyPI, and the Anaconda ecosystem as an open-source Python package.

FEZrs has been employed in research on remote sensing and geospatial analysis. It has been used for principal component analysis (PCA) of Landsat 9 multispectral imagery (Farmahini Farahani et al., 2024), comparison of K-means clustering and spectral indices for land use and land cover characterization (Rezaei & Farmahini Farahani, 2024), and a comparative assessment of SVM, K-means, and spectral indices for land cover classification in engineering geology (Talkhablou et al., 2025). These published applications were conducted by members

of the FEZrs development team and demonstrate the software's use in practical research workflows. FEZrs has also supported geospatial analysis of the Torud Playa area in Isfahan, Iran, providing an additional example of its application in field-based remote sensing and geospatial workflows. In addition, FEZrs received the Merit Award (Ideator Category) at the 9th Iran National Young Scientists Festival.

The project provides comprehensive documentation, practical examples, a test suite, and a modular architecture to support reproducible research and operational geospatial workflows. Detailed installation instructions for both pip (`pip install fezrs`) and conda (`conda install -c FEZtool fezrs`) are available in the FEZrs Installation Guide.

Acknowledgements

The authors would like to express their sincere appreciation to the global open-source developer community. The collective efforts, shared knowledge, and freely available tools developed by this community have provided the essential foundation that made this project possible. Their commitment to collaboration, transparency, and innovation continues to advance technological progress and enables researchers to build upon and improve existing work. The authors gratefully acknowledge these invaluable contributions.

AI Usage Disclosure

While the core software design, implementation, and development decisions of FEZrs were made by the authors, AI-assisted tools were used to assist with unit testing, validation workflows, and manuscript refinement. All AI-assisted outputs were independently reviewed, verified, and validated by the authors.

Conflict of interest

The authors declare no conflict of interest.

References

- Camps-Valls, G., Tuia, D., Zhu, X. X., & Reichstein, M. (Eds.). (2021). *Deep learning for the Earth sciences: A comprehensive approach to remote sensing, climate science and geosciences*. John Wiley & Sons. <https://doi.org/10.1002/9781119646181>
- Chang, Q., Li, X., Li, Y., & Miyazaki, J. (2023). Multi-directional Sobel operator kernel on GPUs. *Journal of Parallel and Distributed Computing*, 177, 160–170. <https://doi.org/10.1016/j.jpdc.2023.03.004>
- Chassot, E., Bonhommeau, S., Reygondeau, G., Nieto, K., Polovina, J. J., Huret, M., Dulvy, N. K., & Demarcq, H. (2011). Satellite remote sensing for an ecosystem approach to fisheries management. *ICES Journal of Marine Science*, 68(4), 651–666. <https://doi.org/10.1093/icesjms/fsq195>
- Cracknell, A. P. (2018). The development of remote sensing in the last 40 years. *International Journal of Remote Sensing*, 39(23), 8387–8427. <https://doi.org/10.1080/01431161.2018.1550919>
- Deng, C., & Wu, C. (2012). BCI: A biophysical composition index for remote sensing of urban environments. *Remote Sensing of Environment*, 127, 247–259. <https://doi.org/10.1016/j.rse.2012.09.009>

- 167 Farmahini Farahani, M., Rezaei, K., Karamollahi, S., & Jafari Moghadam, F. (2024).
 168 Application of PCA algorithm in remote sensing using python programming language (case
 169 study: Landsat 9 multispectral images). *8th Symposium of Sedimentological Society
 170 of Iran*. [https://www.researchgate.net/publication/377838524_Application_of_PCA_](https://www.researchgate.net/publication/377838524_Application_of_PCA_algorithm_in_remote_sensing_using_Python_programming_language_case_study_Landsat_9_multispectral_images)
 171 [algorithm_in_remote_sensing_using_Python_programming_language_case_study_](https://www.researchgate.net/publication/377838524_Application_of_PCA_algorithm_in_remote_sensing_using_Python_programming_language_case_study_Landsat_9_multispectral_images)
 172 [Landsat_9_multispectral_images](https://www.researchgate.net/publication/377838524_Application_of_PCA_algorithm_in_remote_sensing_using_Python_programming_language_case_study_Landsat_9_multispectral_images)
- 173 García Díaz, D. (2026). Ndzi2Gif: A Python package for multi-seasonal remote sensing
 174 analysis with Google Earth Engine. *Journal of Open Source Software*, 11(124), 10654.
 175 <https://doi.org/10.21105/joss.10654>
- 176 Gillies, S., & others. (2013). *Rasterio: Geospatial raster I/O for Python programmers*. Mapbox.
 177 <https://github.com/rasterio/rasterio>
- 178 Harris, C. R., Millman, K. J., Walt, S. J. van der, Gommers, R., Virtanen, P., Cournapeau, D.,
 179 Wieser, E., Taylor, J., Berg, S., Smith, N. J., Kern, R., Picus, M., Hoyer, S., Kerkwijk,
 180 M. H. van, Brett, M., Haldane, A., Río, J. F. del, Wiebe, M., Peterson, P., ... Oliphant,
 181 T. E. (2020). Array programming with NumPy. *Nature*, 585(7825), 357–362. <https://doi.org/10.1038/s41586-020-2649-2>
- 182
- 183 Huang, S., Tang, L., Hupy, J. P., Wang, Y., & Shao, G. (2021). A commentary review on the
 184 use of normalized difference vegetation index (NDVI) in the era of popular remote sensing.
 185 *Journal of Forestry Research*, 32(1), 1–6. <https://doi.org/10.1007/s11676-020-01176-w>
- 186 Huete, A. R. (1988). A soil-adjusted vegetation index (SAVI). *Remote Sensing of Environment*,
 187 26(3), 295–309. [https://doi.org/10.1016/0034-4257\(88\)90106-X](https://doi.org/10.1016/0034-4257(88)90106-X)
- 188 Hunter, J. D. (2007). Matplotlib: A 2D graphics environment. *Computing in Science &
 189 Engineering*, 9(3), 90–95. <https://doi.org/10.1109/MCSE.2007.55>
- 190 Jolliffe, I. T., & Cadima, J. (2016). Principal component analysis: A review and recent
 191 developments. *Philosophical Transactions of the Royal Society A*, 374(2065), 20150202.
 192 <https://doi.org/10.1098/rsta.2015.0202>
- 193 Jordahl, K., Van den Bossche, J., Wasserman, J., McBride, J., Fleischmann, M., Gerard, J.,
 194 Tratner, J., Perry, M., Farmer, C., Hjelle, G. A., Gillies, S., Cochran, M., Bartos, M.,
 195 Culbertson, L., Eubank, N., Bilogur, A., & maxalbert. (2020). *geopandas/geopandas*:
 196 *v0.7.0* (Version 0.7.0). Zenodo. <https://doi.org/10.5281/zenodo.3669853>
- 197 Justusson, B. I. (1981). Median filtering: Statistical properties. In *Two-dimensional digital
 198 signal processing II* (Vol. 43). Springer. <https://doi.org/10.1007/BFb0057597>
- 199 Karnieli, A., Kaufman, Y. J., Remer, L., & Wald, A. (2001). Aerosol free vegetation index
 200 (AFRI) for remote sensing of vegetation. *Remote Sensing of Environment*, 77(1), 10–21.
 201 [https://doi.org/10.1016/S0034-4257\(01\)00190-0](https://doi.org/10.1016/S0034-4257(01)00190-0)
- 202 Kramer, O. (2016). Scikit-learn. In *Machine learning for evolution strategies* (pp. 45–53).
 203 Springer International Publishing. https://doi.org/10.1007/978-3-319-33383-0_5
- 204 Lindeberg, T. (2024). Discrete approximations of Gaussian smoothing and Gaussian derivatives.
 205 *Journal of Mathematical Imaging and Vision*, 66(5), 759–800. <https://doi.org/10.1007/s10851-024-01196-9>
- 206
- 207 Malenovský, Z., Rott, H., Cihlar, J., Schaepman, M. E., García-Santos, G., Fernandes, R., &
 208 Berger, M. (2012). Sentinels for science: Potential of Sentinel-1, -2, and -3 missions for
 209 scientific observations of ocean, cryosphere, and land. *Remote Sensing of Environment*,
 210 120, 91–101. <https://doi.org/10.1016/j.rse.2011.09.026>
- 211 McFeeters, S. K. (1996). The use of the Normalized Difference Water Index (NDWI) in
 212 the delineation of open water features. *International Journal of Remote Sensing*, 17(7),
 213 1425–1432. <https://doi.org/10.1080/01431169608948714>

- McKinney, W. (2010). Data structures for statistical computing in Python. In S. van der Walt & J. Millman (Eds.), *Proceedings of the 9th python in science conference* (pp. 56–61). <https://doi.org/10.25080/Majora-92bf1922-00a>
- Melesse, A. M., Weng, Q., Thenkabail, P. S., & Senay, G. B. (2007). Remote sensing sensors and applications in environmental resources mapping and modelling. *Sensors*, 7(12), 3209–3241. <https://doi.org/10.3390/s7123209>
- Merris, R. (1994). Laplacian matrices of graphs: A survey. *Linear Algebra and Its Applications*, 197, 143–176. [https://doi.org/10.1016/0024-3795\(94\)90486-3](https://doi.org/10.1016/0024-3795(94)90486-3)
- Montero, D., Aybar, C., Mahecha, M. D., Martinuzzi, F., Ramoelo, A., & Campos-Taberner, M. (2023). A standardized catalogue of spectral indices to advance the use of remote sensing in Earth system research. *Scientific Data*, 10, 197. <https://doi.org/10.1038/s41597-023-02096-0>
- Rezaei, K., & Farmahini Farahani, M. (2024). Comparing the performance of the k-means clustering method and spectral indices in characterizing land use and land cover using remote sensing with python (case study: Landsat 8 multi-spectral images). *2nd International Conference for Iranian Geography and Earth Sciences Students*. https://www.researchgate.net/publication/378307658_Comparing_the_performance_of_the_k-means_clustering_method_and_spectral_indices_in_characterizing_land_use_and_land_cover_using_remote_sensing_with_Python_case_study_Landsat_8_multi-spectral_images
- Rikimaru, A., Roy, P. S., & Miyatake, S. (2002). Tropical forest cover density mapping. *Tropical Ecology*, 43(1), 39–47.
- Sabins, F. F. (1999). Remote sensing for mineral exploration. *Ore Geology Reviews*, 14(3-4), 157–183. [https://doi.org/10.1016/s0169-1368\(99\)00007-4](https://doi.org/10.1016/s0169-1368(99)00007-4)
- Shukla, V., Morelli, L., Remondino, F., Micheli, A., Tuia, D., & Risse, B. (2024). Towards estimation of 3D poses and shapes of animals from oblique drone imagery. *International Archives of the Photogrammetry, Remote Sensing and Spatial Information Sciences*, 48, 379–386. <https://doi.org/10.5194/isprs-archives-xxviii-2-2024-379-2024>
- Simic Milas, A., Cracknell, A. P., & Warner, T. A. (2018). Drones—the third generation source of remote sensing data. *International Journal of Remote Sensing*, 39(21), 7125–7137. <https://doi.org/10.1080/01431161.2018.1523832>
- Sinaga, K. P., & Yang, M.-S. (2020). Unsupervised K-means clustering algorithm. *IEEE Access*, 8, 80716–80727. <https://doi.org/10.1109/ACCESS.2020.2988796>
- Sishodia, R. P., Ray, R. L., & Singh, S. K. (2020). Applications of remote sensing in precision agriculture: A review. *Remote Sensing*, 12(19), 3136. <https://doi.org/10.3390/rs12193136>
- Talkhablou, M., Farmahini Farahani, M., & Siah Mansouri, S. (2025). A python-based framework for land cover classification in engineering geology: A comparative assessment of SVM, k-means, and spectral indices. *Kharazmi Journal of Earth Sciences*, 11(1), 201–219. <https://doi.org/10.22034/KJES.2025.11.1.103081>
- Tarazona, Y., Benitez-Paez, F., Nowosad, J., Drenkhan, F., & Timaná, M. E. (2024). Scikit-eo: A Python package for remote sensing data analysis. *Journal of Open Source Software*, 9(99), 6692. <https://doi.org/10.21105/joss.06692>
- Utaminingrum, F., Alqadri, A. M., Somawirata, I. K., Karim, C., Septiarini, A., Lin, C. Y., & Shih, T. K. (2023). Feature selection of gray-level co-occurrence matrix using genetic algorithm with extreme learning machine classification for early detection of pole roads. *Results in Engineering*, 20, 101437. <https://doi.org/10.1016/j.rineng.2023.101437>
- Walt, S. van der, Schönberger, J. L., Nunez-Iglesias, J., Boulogne, F., Warner, J. D., Yager, N., Gouillart, E., Yu, T., & contributors, the scikit-image. (2014). scikit-image: Image

- 262 processing in Python. *PeerJ*, 2, e453. <https://doi.org/10.7717/peerj.453>
- 263 Warmerdam, F. (2008). The geospatial data abstraction library. In G. B. Hall & M. G. Leahy
264 (Eds.), *Open source approaches in spatial data handling* (pp. 87–104). Springer Berlin
265 Heidelberg. https://doi.org/10.1007/978-3-540-74831-1_5
- 266 Williams, D. L., Goward, S., & Arvidson, T. (2006). Landsat. *Photogrammetric Engineering*
267 *& Remote Sensing*, 72(10), 1171–1178. <https://doi.org/10.14358/pers.72.10.1171>
- 268 Xie, G., & Lu, W. (2013). Image edge detection based on OpenCV. *International Journal of*
269 *Electronics and Electrical Engineering*, 1(2), 104–106. [https://doi.org/10.12720/ijeee.1.2.](https://doi.org/10.12720/ijeee.1.2.104-106)
270 [104-106](https://doi.org/10.12720/ijeee.1.2.104-106)
- 271 Xue, J., & Su, B. (2017). Significant remote sensing vegetation indices: A review of
272 developments and applications. *Journal of Sensors*, 2017, 1353691. [https://doi.org/](https://doi.org/10.1155/2017/1353691)
273 [10.1155/2017/1353691](https://doi.org/10.1155/2017/1353691)
- 274 Yue, S., Li, P., & Hao, P. (2003). SVM classification: Its contents and challenges. *Applied*
275 *Mathematics—A Journal of Chinese Universities*, 18(3), 332–342. [https://doi.org/10.1007/](https://doi.org/10.1007/s11766-003-0059-5)
276 [s11766-003-0059-5](https://doi.org/10.1007/s11766-003-0059-5)
- 277 Zhang, B., Wu, Y., Zhao, B., Chanussot, J., Hong, D., Yao, J., & Gao, L. (2022). Progress
278 and challenges in intelligent remote sensing satellite systems. *IEEE Journal of Selected*
279 *Topics in Applied Earth Observations and Remote Sensing*, 15, 1814–1822. [https://doi.](https://doi.org/10.1109/jstars.2022.3148139)
280 [org/10.1109/jstars.2022.3148139](https://doi.org/10.1109/jstars.2022.3148139)