The Journal of Open Source Software

FELINE: A tool to detect emission line galaxies in 3d data

Martin Wendt *[1], **Marvin Henschel** [2], and **Oskar Fjonn Soth** [2]

1 Institute of Physics and Astronomy, University of Potsdam, 14476 Potsdam, Germany **2** Institute of Computer Science and Computational Science, University of Potsdam, 14476 Potsdam, Germany

DOI: 10.xxxxxx/draft

Software

- Review
- Repository
- Archive

Editor: Ivelina Momcheva

Reviewers:

- @smutch
- @bronreichardtchu

Submitted: 12 September 2024
Published: unpublished

License
Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License (CC BY 4.0).

In partnership with

AMERICAN ASTRONOMICAL SOCIETY

This article and software are linked with research article DOI 10.3847/xxxxx <- update this with the DOI from AAS once you know it., published in the Astrophysical Journal <- The name of the AAS journal..

DRAFT

Summary

The detection and classification of objects in astrophysical data has been a key task since the earliest days of astronomy. Over the past decade, the volume of newly observed data has increased dramatically. The advent of integral field unit spectrographs (IFUs), which produce 3D data cubes, has shifted the focus from classical single-target observations to much broader fields of view captured in a single exposure. Simple flux-level peak detection algorithms based on thresholding are prone to either missing many potential real objects or, as a trade-off, producing an abundance of false positives.

Statement of need

The *F*ind *E*mission *LINE*s tool `FELINE` combines a fully parallelized galaxy line template matching with the matched filter approach for individual emission features of LSDcat (Herenz, 2023; Herenz & Wisotzki, 2017).

The VLT/MUSE (R. Bacon et al., 2010, 2014) 3D spectrograph creates $\sim$ 90,000 medium resolution spectra arranged in a 300 $\times$ 300 spatial grid. These data cubes have typical sizes of 3-6 GiB per exposure, the sheer amount of data asks for automated processes to support the scientists. The `FELINE` algorithm evaluates the likelihood of emission lines at specific positions in each spectrum of the data cube. It does this by probing all possible combinations of up to 14 typical emission features, including Hα, Hβ, Hγ, Hδ, [OII], [OIII], [NII], [SII], and [NeIII], for the redshift range of interest ($0.4 < z < 1.4$). This extensive analysis leads to approximately 230,000,000,000 iterations.

Science field

The signal-to-noise cube generated after matched filtering with a 3D emission line template reflects the probability of an emission line at a given spatial and spectral position, which is significantly boosted by the filtering process. As a result, galaxies with multiple weak emission features can be detected with a significance that substantially exceeds the significance of each individual contributing line. This approach is particularly successful for galaxies that show no or little continuum flux in the data, and therefore would generally go undetected in imaging data alone.

`FELINE` was used for the galaxy catalogs of the MEGAFLOW survey in (Bouché et al., 2024; Cherrey et al., 2024; Langan et al., 2023; Schroetter et al., 2024).

*mwendt@astro.physik.uni-potsdam.de

Implementation

The tool uses a brute-force search through the parameter space. Due to the size of the parameter space, the language of implementation was chosen as C for computational efficiency. This approach demonstrates the success of filtering the data with expected templates for individual emission lines, rather than testing full physical models of galaxies (including simulated continuum and temperature-broadened emission lines) against the raw observed data. This reduces the individual models to a single position at which the likelihood of a line is being probed.

For each set of parameters (spatial position in the cube, redshift, and line composition), the `FELINE` algorithm returns the value of the highest-scoring combination, along with its corresponding redshift and line composition.

The data cube contains 300 x 300 spectra, each of which is relatively small ($<$ 64KB). The algorithm performs 512 x 5000 iterations on each spectrum, returning only 3 values: the quality of the best match, the redshift of the best match, and the line combination of the best match. Importantly, the outer 300 x 300 iterations are completely independent of each other.

To take advantage of this independence, the code utilizes full parallelization of the outer loop using `OpenMP`, with most variables shared due to their independence. As a result, FELINE scales quite well with the number of CPU cores. Runtimes for the `FELINE` code on the provided 2.8 GB example cube (Roland Bacon et al., 2023) (CC BY-NC-SA 4.0):

Device	Cores	Runtime in seconds
AMD_EPYC_7542	1	1150
AMD_EPYC_7542	4	282
AMD_EPYC_7542	8	141
AMD_EPYC_7542	16	71
NVIDIA A100 GPU	64	27

Another major improvement in execution time was accomplished by re-arranging the data to maximize the amount of cache hits. Initially, the cube data is stored as a series of images, i.e., 300 x 300 spatial data points arranged in an array of 4,000 in spectral dimension. The algorithm works on spectral which would be strongly interleaved by ~360 KB for consecutive data points and the full spectrum exceeding a range of 1 GiB. As a preprocessing step, the data cube is re-arranged as a spatial grid of full spectra.

That arrangement further motivated an implementation of `FELINE` in `CUDA` to utilize GPUs for parallelization. Typical full size MUSE data cubes can be fully loaded into the GPU memory of any modern `CUDA` capable GPU. We provide a working implementation that produces identical results to the `FELINE` C variant.

Optionally, FELINE plots the three return parameters in real time via SDL surface along with storing them on disk.



Shown are from left to right the quality of the best match, the corresponding redshift of the best match and its template. A fourth panel shows the number of lines that contributed to

DRAFT

the most successful model for ease of human readability (it reflects the number of set bits in the best model value).

We provide a python framework to further visualize and verify the FELINE detections.



Figure 1: Plot generated from the FELINE result.

Acknowledgements

We acknowledge the work on LSDcat by Christian Herenz

References

Bacon, R., Accardo, M., Adjali, L., Anwand, H., Bauer, S., Biswas, I., Blaizot, J., Boudon, D., Brau-Nogue, S., & Brinchmann. (2010). The MUSE second-generation VLT instrument. In I. S. McLean, S. K. Ramsay, & H. Takami (Eds.), *Ground-based and airborne instrumentation for astronomy III* (Vol. 7735, p. 773508). https://doi.org/10.1117/12.856027

Bacon, Roland, Brinchmann, J., Conseil, S., Maseda, M., Nanayakkara, T., Wendt, M., Bacher, R., Mary, D., Weilbacher, P. M., Krajnović, D., Boogaard, L., Bouché, N., Contini, T., Epinat, B., Feltre, A., Guo, Y., Herenz, C., Kollatschny, W., Kusakabe, H., … Zoutendijk, S. L. (2023). The MUSE Hubble Ultra Deep Field surveys: Data release II. *670*, A4. https://doi.org/10.1051/0004-6361/202244187

Bacon, R., Vernet, J., Borisova, E., Bouché, N., Brinchmann, J., Carollo, M., Carton, D., Caruana, J., Cerda, S., Contini, T., Franx, M., Girard, M., Guerou, A., Haddad, N., Hau, G., Herenz, C., Herrera, J. C., Husemann, B., Husser, T.-O., … Zins, G. (2014). MUSE Commissioning. *The Messenger*, *157*, 13–16.

Bouché, N. F., Wendt, M., Zabl, J., Cherrey, M., Schroetter, I., Langan, I., Muzahid, S., Schaye, J., Epinat, B., Wisotzki, L., Contini, T., Richard, J., Bacon, R., & Weilbacher, P. M. (2024). MusE GAs FLOw and Wind (MEGAFLOW) XII. Rationale and design of a MgII survey of the cool circum-galactic medium with MUSE and UVES: The MEGAFLOW Survey. *arXiv e-Prints*, arXiv:2411.07014. https://arxiv.org/abs/2411.07014

Cherrey, M., Bouché, N. F., Zabl, J., Schroetter, I., Wendt, M., Langan, I., Richard, J., Schaye, J., Mercier, W., Epinat, B., & Contini, T. (2024). MusE GAs FLOw and Wind

DRAFT

(MEGAFLOW) X. The cool gas and covering fraction of Mg II in galaxy groups. *528*(1), 481–498. https://doi.org/10.1093/mnras/stad3764

Herenz, E. C. (2023). Revisiting the emission line source detection problem in integral field spectroscopic data. *Astronomische Nachrichten*, *344*(5), e20220091. https://doi.org/10.1002/asna.20220091

Herenz, E. C., & Wisotzki, L. (2017). LSDCat: Detection and cataloguing of emission-line sources in integral-field spectroscopy datacubes. *602*, A111. https://doi.org/10.1051/0004-6361/201629507

Langan, I., Zabl, J., Bouché, N. F., Ginolfi, M., Popping, G., Schroetter, I., Wendt, M., Schaye, J., Boogaard, L., Freundlich, J., Richard, J., Matthee, J., Mercier, W., Contini, T., Guo, Y., & Cherrey, M. (2023). MusE GAs FLOw and Wind (MEGAFLOW) IX. The impact of gas flows on the relations between the mass, star formation rate, and metallicity of galaxies. *521*(1), 546–557. https://doi.org/10.1093/mnras/stad357

Schroetter, I., Bouché, N. F., Zabl, J., Wendt, M., Cherrey, M., Langan, I., Schaye, J., & Contini, T. (2024). MusE GAs FLOw and Wind (MEGAFLOW). XI. Scaling relations between outflows and host galaxy properties. *687*, A39. https://doi.org/10.1051/0004-6361/202348725

DRAFT