

Parcels v4: An Xarray-aligned, flexible Lagrangian simulation framework for geoscience

Nick Hodgskin¹, Joe Schoonover², and Erik van Sebille¹

¹ Institute for Marine and Atmospheric Research, Utrecht University, the Netherlands ² Fluid Numerics, Hickory, NC, USA ¶ Corresponding author

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#)
- [Repository](#)
- [Archive](#)

Editor: [↗](#)

Submitted: 28 September 2026

Published: unpublished

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

Summary

Parcels ([Delandmeter & van Sebille, 2019](#); [Lange & van Sebille, 2017](#)) is a highly customisable Lagrangian simulation framework. Version 4 of the software is a major update which overhauls the software internals to natively leverage Xarray ([Hoyer & Hamman, 2017](#)) dataset objects, and adds support for simulating particles on unstructured grids. Xarray support makes Parcels compatible with many new data formats (e.g., Zarr, Icechunk) and execution modes (e.g., streaming data from cloud buckets or other data providers - such as the [Copernicus Marine Data Store](#)), while support for unstructured grid datasets enables simulations on (combinations of) different grid geometries. Version 4 of Parcels also provides support for custom interpolators (surfacing to scientists even more control over the numerics of their simulation), and trajectory output in Parquet format.

Statement of need

The first release of Parcels was in July 2017. Since then, the geoscience modelling space has changed. A significant portion of the geospatial science community has shifted from creating their own scripts for manipulating NetCDF data, to using open source software - like Xarray - for working with multidimensional climate data. Major benefits of Xarray in particular include (a) providing a single in-memory, metadata-rich representation of a full NetCDF-like dataset, (b) its flexibility to manipulate datasets in other data-formats (e.g., Zarr, HDF5), and (c) its abstractions that facilitate easy data ingestion from network data sources.

Writing software that natively works from Xarray datasets provides a powerful abstraction layer allowing downstream developers to create software that works across data formats and data ingestion paradigms. This is particularly important as climate datasets are continually increasing in resolution and size, affecting how the data is stored. Datasets are increasingly being stored in the cloud using more modern data formats, such as Zarr.

Another interesting change is that climate modellers are increasingly providing model output on different grid geometries, which have attractive features compared to conventional structured grids. Running natively on these grid geometries, which can be represented as unstructured grids, without re-interpolation allows researchers to run simulations that fully capture the details from the original dataset.

Finally, an important point is the cross-domain generalisability of Lagrangian particle tracking tools within the geosciences. Many of these tools originate from a specific domain, and (as evidenced by documentation, API naming, and input data support) cater to a userbase within the same domain. This is despite the fact the numerics of the tools themselves is largely identical between the domains. This results in both fractured communities, and toolsets whose usage mirrors the blurry boundaries of the domains themselves. Having cross-domain tooling

41 for Lagrangian particle tracking promotes cross-collaboration, software quality, building of
42 better standards, and the overall impact of the software in users scientific workflows.

43 State of the field

44 Parcels is not the only open-source Lagrangian particle tracking framework for research. Other
45 notable frameworks for oceanographic applications include OpenDrift (Dagestad et al., 2018),
46 TRACMASS (Döös et al., 2017), Ariane (Blanke & Raynaud, 1997), Drifters.jl (Forget, 2021),
47 TrackMPD (Jalón-Rojas et al., 2019), Ichthyop (Barrier et al., 2026), and the Connectivity
48 Modelling System (Paris et al., 2013) – each with their own strengths and limitations. OpenDrift
49 is very good for Search-and-Rescue simulations and operational forecasting, TRACMASS and
50 Ariane excel in deterministic (analytical) particle tracking on structured grids, Drifters.jl is useful
51 for advanced diagnostics, TrackMPD is designed specifically for marine plastic applications, and
52 Ichthyop and the Connectivity Modelling System are especially useful for biological applications.
53 Furthermore, many of the hydrodynamic models also have their (online) particle tracking
54 modules, but these are often tightly coupled to the specific model and lack the flexibility
55 offered by standalone frameworks like Parcels. In the atmosphere, FlexPart (Stohl et al., 1998)
56 is a widely used Lagrangian particle tracking framework. However, none of these packages
57 come with the flexibility in terms of structured and unstructured grid support, and custom
58 kernels and interpolators, that Parcels offers.

59 Software design

60 When working with output from a single circulation model, version 4 of Parcels assumes data
61 (i.e., the field data and mesh data) is all contained in a single Xarray Dataset object that has
62 appropriate metadata. This required metadata includes certain CF-convention metadata (Eaton
63 et al., 2020), providing information about important variables/coordinates and their units,
64 as well SGRID or UGRID metadata, providing grid geometry information for structured grid
65 data and unstructured grid data respectively. This metadata rich object can either be opened
66 directly from disk/a data store, or is constructed by the user. Parcels provides “converter”
67 functions for various circulation models which understand model-specific conventions, that
68 then attach this metadata appropriately. This overall approach allows the internals of Parcels
69 to assume a certain dataset structure and metadata richness, while still allowing the software
70 to transparently facilitate users who come with metadata-poor model data. The diagram in
71 Figure 1 illustrates potential code paths when working from model data to a fully constructed
72 FieldSet. If a user wants to run a simulation with fields from different models, they load each
73 model data into its own FieldSet and then combine the FieldSets together into a single FieldSet
74 simply with an addition.




Figure 1: Code path for FieldSet ingestion in v4 of Parcels

A strength of earlier Parcels versions has been the ability for users to write custom “kernels” which encode particle actions over the course of a simulation. This flexibility has enabled users to model a wide range of physical phenomena, from plastic pollution to plankton and fish larvae. Parcels version 4 adds custom interpolators, allowing users to also have control on how field data is interpolated at particle positions. Users can use a range of pre-packaged interpolators, or write their own, and set them on a field-by-field basis overriding the default linear interpolators.

Version 4 of Parcels also improves generalisability to other domains, such as atmospheric or cryospheric particle tracking. It allows the encoding of various input datasets, relaxes domain specific requirements, replaces domain specific terminology, and updates the branding surrounding the software.

Parcels version 4 also changes the output format from Zarr to Parquet, aligning better with the tabular nature of particle trajectory output.

Example use-case

A key use-case of the Parcels version 4 is the combining of various model data which not only have different sources, but also have very different grid geometries. In this section we present an example simulation focused on the Dutch coast.

We combine flow data from Deltares’ 3D DCSM-FM model, SWAN Wave model data from KNMI, and wind model data from Copernicusmarine. The 3D DCSM-FM model is an unstructured model, SWAN has a structured curvilinear grid, and Copernicusmarine provides data on a structured rectilinear grid geometry. We seed particles along the Dutch coast, and within estuaries facing the North Sea. We advect the particles for a total of a month (from

2025-11-01 to 2025-12-01) with a timestep of 10 minutes. Figure 2 shows the varied types of data and grid geometries that we combine, along with the resulting particle tracks.




Figure 2: Trajectories of almost 5,000 particles seeded near the port of Rotterdam in The Netherlands. Trajectories are coloured by start date, and final locations of the particles are marked by circles. The mesh of the unstructured flow model (DCSM-FM) is shown in grey, the mesh of the rectilinear Copernicusmarine wind model is shown in red, and the mesh of the curvilinear KNMI wave model is shown in blue.

Research impact statement

Parcels has been cited in over 330 peer reviewed scientific papers so far mostly within the field of oceanography.

AI usage disclosure

Large Language Models (Claude Opus 4.6, ChatGPT , GitHub Copilot) have been used in a guided manner for code generation, documentation, refactoring, and testing. All LLM written code and documentation has been verified by the authors. This manuscript was fully written by humans.

Acknowledgements

This work has been funded via Nederlandse Organisatie voor Wetenschappelijk Onderzoek, Exacte en Natuurwetenschappen (VI.C.222.025) as part of the project “Tracing Marine Macroplastics by Unraveling the Ocean’s Multiscale Transport Processes” and via the

111 Warmworld ELPHE project, and compute on the SURF Research Cloud via project
112 EINF-15719.

113 References

- 114 Barrier, N., Verley, P., Lett, C., & Mullon, C. (2026). *Ichthyop: A Lagrangian tool for modelling*
115 *ichthyoplankton dynamics*. Zenodo. <https://doi.org/10.5281/ZENODO.4243813>
- 116 Blanke, B., & Raynaud, S. (1997). Kinematics of the Pacific Equatorial Undercurrent: An
117 Eulerian and Lagrangian approach from GCM results. *Journal of Physical Oceanography*,
118 27(6), 1038–1053.
- 119 Dagestad, K.-F., Röhrs, J., Breivik, Ø., & Ådlandsvik, B. (2018). OpenDrift v1.0: A generic
120 framework for trajectory modelling. *Geoscientific Model Development*, 11(4), 1405–1420.
121 <https://doi.org/10.5194/gmd-11-1405-2018>
- 122 Delandmeter, P., & van Sebille, E. (2019). The Parcels v2.0 Lagrangian framework: new
123 field interpolation schemes. *Geoscientific Model Development*, 12(8), 3571–3584. <https://doi.org/10.5194/gmd-12-3571-2019>
- 125 Döös, K., Jönsson, B., & Kjellsson, J. (2017). Evaluation of oceanic and atmospheric trajectory
126 schemes in the TRACMASS trajectory model v6. 0. *Geoscientific Model Development*, 10,
127 1733–1749. <https://doi.org/10.5194/gmd-10-1733-2017>
- 128 Eaton, B., Gregory, J., Drach, B., Taylor, K., Hankin, S., Caron, J., Signell, R., Bentley,
129 P., Rappa, G., Höck, H., Pamment, A., Jukes, M., Raspaud, M., Horne, R., Whiteaker,
130 T., Blodgett, D., Zender, C., & Lee, D. (2020). *NetCDF Climate and Forecast (CF)*
131 *Metadata Conventions v1.8*. CF Conventions Committee. [http://cfconventions.org/Data/](http://cfconventions.org/Data/cf-conventions/cf-conventions-1.8/cf-conventions.html)
132 [cf-conventions/cf-conventions-1.8/cf-conventions.html](http://cfconventions.org/Data/cf-conventions/cf-conventions-1.8/cf-conventions.html)
- 133 Forget, G. (2021). IndividualDisplacements.jl: A julia package to simulate and study particle
134 displacements within the climate system. *Journal of Open Source Software*, 6(60), 2813.
135 <https://doi.org/10.21105/joss.02813>
- 136 Hoyer, S., & Hamman, J. (2017). Xarray: N-D labeled arrays and datasets in python. *J. Open*
137 *Res. Softw.*, 5(1), 10.
- 138 Jalón-Rojas, I., Wang, X. H., & Fredj, E. (2019). A 3D numerical model to Track Marine
139 Plastic Debris (TrackMPD): Sensitivity of microplastic trajectories and fates to particle
140 dynamical properties and physical processes. *Marine Pollution Bulletin*, 141, 256–272.
141 <https://doi.org/10.1016/j.marpolbul.2019.02.052>
- 142 Lange, M., & van Sebille, E. (2017). Parcels v0.9: prototyping a Lagrangian ocean analysis
143 framework for the petascale age. *Geoscientific Model Development*, 10(11), 4175–4186.
144 <https://doi.org/10.5194/gmd-10-4175-2017>
- 145 Paris, C. B., Helgers, J., Sebille, E. van, & Srinivasan, A. (2013). Connectivity Modeling
146 System: A probabilistic modeling tool for the multi-scale tracking of biotic and abiotic
147 variability in the ocean. *Environmental Modelling & Software*, 42(C), 47–54. <https://doi.org/10.1016/j.envsoft.2012.12.006>
- 149 Stohl, A., Hittenberger, M., & Wotawa, G. (1998). Validation of the lagrangian particle
150 dispersion model FLEXPART against large-scale tracer experiment data. *Atmospheric*
151 *Environment*, 32(24), 4245–4264. [https://doi.org/10.1016/S1352-2310\(98\)00184-8](https://doi.org/10.1016/S1352-2310(98)00184-8)