The Journal of Open Source Software

AquaCrop.jl: A Process-Based Model of Crop Growth

Gabriel Díaz Iturry[1,2], **Marco C. Matthies**[1,2], **Guy Pe'er**[1,2], **and Daniel Vedder**[1,2,3¶]

1 Helmholtz Centre for Environmental Research - UFZ **2** German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig **3** Friedrich-Schiller-Universität Jena ¶ Corresponding author

DOI: 10.xxxxxx/draft

Software

- Review
- Repository
- Archive

Editor: Ethan White

Reviewers:

- @julienmalard
- @tonyewong

Submitted: 21 January 2025
Published: unpublished

License
Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License (CC BY 4.0).

DRAFT

Summary

AquaCrop is a simulation model that forecasts the growth and yield of crop plants under different environmental and management conditions (Steduto et al., 2009). As a process-based model, it can be used to optimise farm management, forecast regional yields, or study climate change impacts and sustainable farming practices. Originally developed by the Food and Agricultural Organization of the United Nations (FAO), it has been widely applied in agricultural research.

Here, we present an expanded reimplementation of the model in Julia (Bezanson et al., 2017), focussing on improving its interoperability with other software and models. With `AquaCrop.jl`, we want to make AquaCrop available to the growing number of environmental modellers working in Julia, and contribute to the creation of integrated, interdisciplinary models in the environmental sciences.

Statement of need

All agriculture is dependent on the growth of plants, which provide food, fodder, fibre, and other resources. Forecasting and optimising crop growth is therefore vital to address global food system challenges, including widespread malnutrition, agriculture-related environmental degradation, and global impacts of climate change (Foley et al., 2011). In this context, process-based crop models can be used to inform decision-making. These models use information about environmental parameters (e.g. temperature, rainfall, soil quality) and knowledge of plant biology to simulate crop growth over time and estimate yield.

AquaCrop is one such model, which has been used to model numerous crops worldwide (Mialyk et al., 2024), and is known to produce reliable estimates of crop phenology and yield (Kostková et al., 2021). First implemented in Delphi, it was later open-sourced in a Fortran version (de Roos et al., 2021; Remote Sensing and Data Assimilation Research Group, 2021/2024). There are also versions available in Matlab, Python, and R, although currently these are not up-to-date with the most recent version of the original model (Camargo Rodriguez & Ober, 2019; Foster et al., 2017; Kelly & Foster, 2021).

`AquaCrop.jl` expands this portfolio to contribute to the emerging ecosystem of environmental research software in Julia. To our knowledge, this is the first process-based crop model available in this language. Our purpose is also to provide a package that can be readily integrated into other scientific software, in order to facilitate the creation of multidisciplinary models of socio-environmental systems (Cabral et al., 2023; Vedder, Fischer, et al., 2024).

Specifically, we developed the package to use it as a component within `Persefone.jl`, a process-based model of agricultural ecosystems (Vedder, Matthies, et al., 2024). The aim of this model is to study the impact that agricultural management and policy has on biodiversity, for which the growth of crop plants is an important mediating factor.

Comparison to original implementation

The core code of `AquaCrop.jl` was translated verbatim from the original Fortran implementation, which allows us to quickly integrate changes and updates to the original `AquaCrop` code. `AquaCrop.jl` supports the original input file formats, and is tested to ensure its output conforms to that of the original software.

On top of this core code, we developed a wrapper layer with an API that improves the interoperability of `AquaCrop.jl` when used as a package with other software. First, we added support for standardised input and output file formats (TOML and CSV), and for loading input data from memory rather than disk (for example using output from a coupled model). Second, we bundled all state variables for a simulation in one struct (`AquaCropField`), thereby eliminating global state and allowing multiple simulations to be carried out in parallel, as well as making serialisation and data transfer easier. Third, we enabled the model to be updated one day at a time, rather than being executed as a singe batch job. This allows state variables to be inspected and changed on the go, which makes it possible to use the package for bidirectional model coupling as well as interactively.

Overall, we leave the scientific core of the model unchanged, but make it easier for environmental modellers using Julia to integrate the model into their own work, interface with other libraries for the Julia ecosystem, or adapt the model to suit their needs.

Examples

Multiple tutorials for different use cases are provided in the documentation. A simple demonstration of a basic run using the data from the `AquaCrop.jl/test/testcase` directory is shown here:

```julia
using AquaCrop
using CairoMakie
using Unitful

# First, we specify the input file format:
# NormalFileRun(): use the original AquaCrop file format
# TomlFileRun(): use TOML and CSV formatting
# NoFileRun(): provide input data via the API
runtype = NormalFileRun();

# Then specify the directory containing the necessary input files
parentdir = AquaCrop.test_dir;

# Now we can do a simulation run and plot the results
outputs = basic_run(; runtype=runtype, parentdir=parentdir);
function plot_basic_out(cropfield, cols)
    x = cropfield[!, "Date"]
    aux_sz = round(Int, sqrt(length(cols)))
    f = Figure()
    for (i, coli) in enumerate(keys(cols))
        ii, jj = divrem(i-1, aux_sz)
        ax = Axis(f[ii, jj],
                title = cols[coli][1],
                xlabel = "Date",
                ylabel = cols[coli][2]
                )
        lines!(ax, x, ustrip.(cropfield[!, coli]))
```

DRAFT

```
        ax.xticklabelrotation = π/4
        ax.xticklabelsize = 8
        ax.yticklabelsize = 8
    end
    return f
end

f = plot_basic_out(outputs[:dayout],
    Dict("CC"=>["Canopy Cover","%"],
         "Tavg"=>["Temperature","K"],
         "Biomass"=>["Biomass","ton/ha"],
         "Rain"=>["Rainfall","mm"]))
```



Figure 1: Simulated canopy cover and biomass of an alfalfa crop over time, together with daily temperature and rainfall.

The resulting graph is shown in Figure 1. Canopy cover increases during the growing season, with regular harvests taking place. Biomass is shown accumulated over the whole season.

Figure 2 shows a simulation of the growth of beans (*Vicia faba*) based on environmental data from Thuringia, Germany, with historical yield data shown for comparison. This showcases that when well parameterised, `AquaCrop.jl` forecasts the development of yields over time quite reliably.



Figure 2: Simulated yield of beans (*Vicia faba*) compared to observed yields in Thuringia, Germany.

Acknowledgements

GDI, MM, and DV are funded through the project CAP4GI by the Federal Ministry of Education and Research (BMBF), within the framework of the Strategy, Research for Sustainability (FONA, www.fona.de/en) as part of its Social-Ecological Research funding priority, funding no. 01UT2102A. Responsibility for the content of this publication lies with the authors. MM, GP, and DV gratefully acknowledge the support of iDiv, funded by the German Research Foundation (DFG–FZT 118, 202548816).

References

Bezanson, J., Edelman, A., Karpinski, S., & Shah, V. B. (2017). Julia: A fresh approach to numerical computing. *SIAM Review*, *59*(1), 65–98. https://doi.org/10.1137/141000671

Cabral, J. S., Mendoza-Ponce, A., da Silva, A. P., Oberpriller, J., Mimet, A., Kieslinger, J., Berger, T., Blechschmidt, J., Brönner, M., Classen, A., Fallert, S., Hartig, F., Hof, C., Hoffmann, M., Knoke, T., Krause, A., Lewerentz, A., Pohle, P., Raeder, U., … Zurell, D. (2023). The road to integrate climate change projections with regional land-use–biodiversity models. *People and Nature*, *6*(5), 1716–1741. https://doi.org/10.1002/pan3.10472

Camargo Rodriguez, A. V., & Ober, E. S. (2019). AquaCropR: Crop Growth Model for R. *Agronomy*, *9*(7, 7), 378. https://doi.org/10.3390/agronomy9070378

de Roos, S., De Lannoy, G. J. M., & Raes, D. (2021). Performance analysis of regional AquaCrop (v6.1) biomass and surface soil moisture simulations using satellite and in situ observations. *Geoscientific Model Development*, *14*(12), 7309–7328. https://doi.org/10.5194/gmd-14-7309-2021

Foley, J. A., Ramankutty, N., Brauman, K. A., Cassidy, E. S., Gerber, J. S., Johnston, M., Mueller, N. D., O'Connell, C., Ray, D. K., West, P. C., Balzer, C., Bennett, E. M., Carpenter, S. R., Hill, J., Monfreda, C., Polasky, S., Rockström, J., Sheehan, J., Siebert, S., … Zaks, D. P. M. (2011). Solutions for a cultivated planet. *Nature*, *478*(7369), 337–342. https://doi.org/10.1038/nature10452

Foster, T., Brozović, N., Butler, A. P., Neale, C. M. U., Raes, D., Steduto, P., Fereres, E., & Hsiao, T. C. (2017). AquaCrop-OS: An open source version of FAO's crop water productivity model. *Agricultural Water Management*, *181*, 18–22. https://doi.org/10.1016/j.agwat.2016.11.015

Kelly, T. D., & Foster, T. (2021). AquaCrop-OSPy: Bridging the gap between research and practice in crop-water modeling. *Agricultural Water Management*, *254*, 106976. https://doi.org/10.1016/j.agwat.2021.106976

Kostková, M., Hlavinka, P., Pohanková, E., Kersebaum, K. C., Nendel, C., Gobin, A., Olesen, J. E., Ferrise, R., Dibari, C., Takáč, J., Topaj, A., Medvedev, S., Hoffmann, M. P., Stella, T., Balek, J., Ruiz-Ramos, M., Rodríguez, A., Hoogenboom, G., Shelia, V., … Trnka, M. (2021). Performance of 13 crop simulation models and their ensemble for simulating four field crops in Central Europe. *The Journal of Agricultural Science*, *159*(1-2), 69–89. https://doi.org/10.1017/S0021859621000216

Mialyk, O., Schyns, J. F., Booij, M. J., Su, H., Hogeboom, R. J., & Berger, M. (2024). Water footprints and crop water use of 175 individual crops for 1990–2019 simulated with a global crop model. *Scientific Data*, *11*(1), 206. https://doi.org/10.1038/s41597-024-03051-3

Remote Sensing and Data Assimilation Research Group. (2024). *AquaCrop repository* (Version 7.2). KU Leuven. https://github.com/KUL-RSDA/AquaCrop (Original work published 2021)

Steduto, P., Hsiao, T. C., Raes, D., & Fereres, E. (2009). AquaCrop—The FAO Crop Model to Simulate Yield Response to Water: I. Concepts and Underlying Principles. *Agronomy Journal*, *101*(3), 426–437. https://doi.org/10.2134/agronj2008.0139s

Vedder, D., Fischer, S. M., Wiegand, K., & Pe'er, G. (2024). Developing multidisciplinary mechanistic models: Challenges and approaches. *Socio-Environmental Systems Modelling*, *6*, 18701. https://doi.org/10.18174/sesmo.18701

Vedder, D., Matthies, M., & Pe'er, G. (2024). *Persefone.jl* (Version 0.6.0). German Centre of Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig. https://persefone-model.eu/

DRAFT