The Journal of Open Source Software

gaia: An R package to estimate crop yield responses to temperature and precipitation

Mengqi Zhao [1], **Stephanie Waldhoff** [2], **Claudia Tebaldi** [2], **and Abigail Snyder** [2]

1 Pacific Northwest National Laboratory, Richland, WA, USA **2** Pacific Northwest National Laboratory, Joint Global Change Research Institute, College Park, MD, USA

DOI: 10.xxxxxx/draft

Software

- Review
- Repository
- Archive

Editor: Taher Chegini

Reviewers:

- @klau506
- @jrising

Submitted: 14 October 2024
Published: unpublished

License
Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License (CC BY 4.0).

DRAFT

Summary

gaia is an open-source R package designed to estimate crop yield shocks in response to annual weather variations and CO2 concentrations at the country scale for 12 major crops. This innovative tool streamlines the workflow from raw climate data processing to projections of annual shocks to crop yields at the country level, using the response surfaces developed and documented in Waldhoff et al. (2020), an empirical econometric model that leverages historical weather, CO_2, and crop yield data for robust empirical fitting for 12 crops. gaia uses these response surfaces with bias-corrected, gridded monthly temperature and precipitation projections (e.g., from the Coupled Model Intercomparison Project Phase 6 (CMIP6, O'Neill et al. (2016)) and Inter-Sectoral Impact Model Intercomparison Project (ISIMIP, Warszawski et al. (2014))) to project shocks that can be applied to agricultural productivity changes at the country-level for use in economic models. The historical and future projections use gridded, country-crop specific monthly growing season precipitation and temperature data, aggregated to the national level, weighted by cropland area derived from MIRCA (Portmann et al., 2010). These annual, country, crop-specific yield shocks can be aggregated to different definitions of regions, crop commodities, and time periods, as needed by specific economic models. gaia serves as a lightweight, powerful model that can aid exploration of crop yields responses under a broad range of future climate projections, enhancing human-Earth system analysis capabilities.

Statement of need

Agricultural production is highly responsive to annual weather patterns, which are becoming increasingly variable in response to climate change (e.g. Iizumi & Ramankutty (2016)). Agricultural markets are global, with significant volumes of international trade, and yield shocks due to this variability in one region will therefore impact others. In addition, agricultural production is closely connected with energy and water sectors. Improving the representation of agricultural production and their yield responses to changing weather patterns in global multi-sector dynamics models, such as the Global Change Analysis Model (GCAM, Calvin et al. (2019), Joint Global Change Research Institute (2023)), is key to better understanding economic impacts of climate change that account for human-Earth system interactions.

Most economic, multisectoral modeling of climate change impacts on agriculture address only the long-term trends in temperature and precipitation effects on crop yields for a few key crops (Lampe et al., 2014). There is a need to include projections of the impacts of inter-annual weather changes on yield responses for multiple crop under a wide range of climate projections (Ray et al., 2015). However, the current crop modeling landscape lacks globally comprehensive, computationally efficient, validated models capable of the aforementioned needs (Waldhoff

et al., 2020). To address the gap, we have developed gaia, which uses the empirical model described in Waldhoff et al. (2020). gaia produces projections of both future annual variations and long-term trends in yield changes at the country-level for a wide range of crops (cassava, cotton, maize, rice, potatoes, sorghum, soybean, sugar beet, sugarcane, sunflower, and wheat) using gridded monthly temperature and precipitation inputs for any future climate scenario.

States of the Field

The exploration of effects of future climate change on agricultural production is crucial for providing insights into global food, energy, water, and economic development. Various crop models, including process-based and empirical models, have been developed to simulate crop yields under different climate scenarios (Chapagain et al., 2022; Rauff & Bello, 2015). Müller et al. (2017) conducted a global gridded crop model (GGCM) intercomparison experiment involving 14 process-based crop models, which allow users to simulate diverse crop management options, soil types, and weather conditions. However, GGCMs require large amounts of site-specific data for calibration, making extension to many crops and regions more challenging (Müller et al., 2017). In addition, these models can take significant time to run and the yield shock projections are often available for only a limited number of climate change scenarios.

These considerations may limit the usefulness of yield changes from process models as inputs to global economic, multisector models, which include a wider range of crops and many future climate scenarios. gaia achieves this balance by employing computationally efficient statistical methods to model 12 major crops globally, while maintaining the sensitivity to capture both gradual climate changes and interannual variability impacts on crop yields. In addition, users can estimate yield shocks for additional crops by providing the appropriate input data.

Functionality

The primary functionality of gaia is encapsulated in the yield_impact wrapper function that streamlines the entire workflow shown in Figure 1. The modular design also facilitates comprehensive diagnostic outputs, enhancing the tool's utility for researchers and decision makers. Users can also execute individual functions to work through the main steps of the process. Detailed instructions on gaia can be accessed at https://jgcri.github.io/gaia.

1. `weighted_climate`: Processes CMIP-ISIMIP climate NetCDF data and calculates cropland-weighted precipitation and temperature at the country level, differentiated by crop type and irrigation type. The function accepts both daily or monthly climate data that are consistent with the CMIP-ISIMIP NetCDF data format
2. `crop_calenders`: Generates crop planting months for each country and crop based on crop calendar data (Sacks et al., 2010).
3. `data_aggregation`: Calculates crop growing seasons using climate variables processed by `weighted_climate` and crop calendars for both historical and projected periods. This function prepares climate and yield data for subsequent model fitting.
4. `yield_regression`: Performs regression analysis fitted with historical annual crop yields, monthly growing season temperature and precipitation, CO_2 concentrations, GDP per capita, and year. The default econometric model applied in gaia is from Waldhoff et al. (2020). User can specify alternative formulas that are consistent with the data processed in `data_aggregation`.
5. `yield_shock_projection`: Projects yield shocks for future climate scenarios using the fitted model and temperature, precipitation, and CO_2 projections from the climate scenario.
6. `gcam_agprodchange`: Remaps country-level yield shocks to GCAM-required spatial scales (i.e., region, basin, technology intersections), based on harvested areas, and aggregates crops to GCAM commodities. This function applies the projected shocks to GCAM

DRAFT

scenario agricultural productivity growth rates (the unit used to project future yields in GCAM) and creates ready-to-use XML outputs for GCAM.



Figure 1: The gaia workflow showing the functions and the corresponding outputs of modeling crop yield shocks to weather variables using an empirical econometric model.

Acknowledgements

This research was supported by the US Department of Energy, Office of Science, as part of research in MultiSector Dynamics, Earth and Environmental System Modeling Program. The Pacific Northwest National Laboratory is operated for DOE by Battelle Memorial Institute under contract DE-AC05-76RL01830. The views and opinions expressed in this paper are those of the authors alone.

References

Calvin, K., Patel, P., Clarke, L., Asrar, G., Bond-Lamberty, B., Cui, R. Y., Di Vittorio, A., Dorheim, K., Edmonds, J., Hartin, C., Hejazi, M., Horowitz, R., Iyer, G., Kyle, P., Kim, S., Link, R., McJeon, H., Smith, S. J., Snyder, A., … Wise, M. (2019). GCAM v5.1: Representing the linkages between energy, water, land, climate, and economic systems. *Geoscientific Model Development*, *12*(2), 677–698. https://doi.org/10.5194/gmd-12-677-2019

Chapagain, R., Remenyi, T. A., Harris, R. M. B., Mohammed, C. L., Huth, N., Wallach, D., Rezaei, E. E., & Ojeda, J. J. (2022). Decomposing crop model uncertainty: A systematic review. *Field Crops Research*, *279*, 108448. https://doi.org/10.1016/j.fcr.2022.108448

Iizumi, T., & Ramankutty, N. (2016). Changes in yield variability of major crops for 1981–2010 explained by climate change. *Environmental Research Letters*, *11*(3), 034003. https://doi.org/10.1088/1748-9326/11/3/034003

Joint Global Change Research Institute. (2023). *GCAM Documentation* (Version 7.0). https://github.com/JGCRI/gcam-doc

Lampe, M. von, Willenbockel, D., Ahammad, H., Blanc, E., Cai, Y., Calvin, K., Fujimori, S., Hasegawa, T., Havlik, P., Heyhoe, E., Kyle, P., Lotze-Campen, H., Mason d'Croz, D.,

Nelson, G. C., Sands, R. D., Schmitz, C., Tabeau, A., Valin, H., Mensbrugghe, D. van der, & Meijl, H. van. (2014). Why do global long-term scenarios for agriculture differ? An overview of the AgMIP global economic model intercomparison. *Agricultural Economics*, *45*(1), 3–20. https://doi.org/10.1111/agec.12086

Müller, C., Elliott, J., Chryssanthacopoulos, J., Arneth, A., Balkovic, J., Ciais, P., Deryng, D., Folberth, C., Glotter, M., Hoek, S., Iizumi, T., Izaurralde, R. C., Jones, C., Khabarov, N., Lawrence, P., Liu, W., Olin, S., Pugh, T. A. M., Ray, D. K., … Yang, H. (2017). Global gridded crop model evaluation: Benchmarking, skills, deficiencies and implications. *Geoscientific Model Development*, *10*(4), 1403–1422. https://doi.org/10.5194/gmd-10-1403-2017

O'Neill, B. C., Tebaldi, C., Vuuren, D. P. van, Eyring, V., Friedlingstein, P., Hurtt, G., Knutti, R., Kriegler, E., Lamarque, J.-F., Lowe, J., Meehl, G. A., Moss, R., Riahi, K., & Sanderson, B. M. (2016). The scenario model intercomparison project (ScenarioMIP) for CMIP6. *Geoscientific Model Development*, *9*(9), 3461–3482. https://doi.org/10.5194/gmd-9-3461-2016

Portmann, F. T., Siebert, S., & Döll, P. (2010). MIRCA2000—global monthly irrigated and rainfed crop areas around the year 2000: A new high-resolution data set for agricultural and hydrological modeling. *Global Biogeochemical Cycles*, *24*(1). https://doi.org/10.1029/2008GB003435

Rauff, K. O., & Bello, R. (2015). A review of crop growth simulation models as tools for agricultural meteorology. *Agricultural Sciences*, *6*(9), 1098–1105. https://doi.org/10.4236/as.2015.69105

Ray, D. K., Gerber, J. S., MacDonald, G. K., & West, P. C. (2015). Climate variation explains a third of global crop yield variability. *Nature Communications*, *6*(1), 5989. https://doi.org/10.1038/ncomms6989

Sacks, W. J., Deryng, D., Foley, J. A., & Ramankutty, N. (2010). Crop planting dates: An analysis of global patterns. *Global Ecology and Biogeography*, *19*(5), 607–620. https://doi.org/10.1111/j.1466-8238.2010.00551.x

Waldhoff, S. T., Wing, I. S., Edmonds, J., Leng, G., & Zhang, X. (2020). Future climate impacts on global agricultural yields over the 21st century. *Environmental Research Letters*, *15*(11), 114010. https://doi.org/10.1088/1748-9326/abadcb

Warszawski, L., Frieler, K., Huber, V., Piontek, F., Serdeczny, O., & Schewe, J. (2014). The inter-sectoral impact model intercomparison project (ISI–MIP): Project framework. *Proceedings of the National Academy of Sciences*, *111*(9), 3228–3232. https://doi.org/10.1073/pnas.1312330110

DRAFT