The Journal of Open Source Software

GWSurrogate: A Python package for gravitational wave surrogate models

Scott E. Field [1,6*], **Vijay Varma** [1*], **Jonathan Blackman** [2], **Bhooshan Gadre** [3], **Chad R. Galley** [4], **Tousif Islam** [1,5], **Keefe Mitman** [7], **Michael Pürrer** [6], **Adhrit Ravichandran** [1], **Mark A. Scheel** [7], **Leo C. Stein** [8], and **Jooheon Yoo** [9]

1 Department of Mathematics and Center for Scientific Computing & Data Science Research, University of Massachusetts, Dartmouth, MA 02747, USA **2** Theorem Partners LLC, San Mateo, California 94402, USA **3** Institute for Gravitational and Subatomic Physics (GRASP), Utrecht University, 3584 CC Utrecht, The Netherlands **4** Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA 91109, USA **5** Kavli Institute for Theoretical Physics, University of California Santa Barbara, CA 93106, USA **6** Department of Physics and Center for Computational Research, East Hall, University of Rhode Island, Kingston, RI 02881 **7** Theoretical Astrophysics 350-17, California Institute of Technology, Pasadena, California 91125, USA **8** Department of Physics and Astronomy, The University of Mississippi, University, MS 38677, USA **9** Cornell Center for Astrophysics and Planetary Science, Cornell University, Ithaca, New York 14853, USA * These authors contributed equally.

DOI: 10.xxxxxx/draft

Software

- Review
- Repository
- Archive

Editor:

Submitted: 03 June 2024
Published: unpublished

License
Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License (CC BY 4.0).

DRAFT

Summary

Gravitational waves are ripples in space-time caused by the motion of massive objects. One of the most astrophysically important sources of gravitational radiation is caused by two orbiting compact objects, such as black holes and neutron stars, that slowly inspiral and merge. The motion of these massive objects generates gravitational waves that radiate to the far field where gravitational-wave detectors can observe them. Complicated partial or ordinary differential equations govern the entire process.

Traditionally, the dynamics of compact binary systems and the emitted gravitational waves have been computed by expensive simulation codes that can take days to months to run. A key simulation output is the gravitational wave signal for a particular set of parameter values describing the system, such as the black holes' masses and spins. The computed signal is required for a diverse range of multiple-query applications, such as template bank generation for searches, parameter estimation, mock data analysis, studies of model bias, and tests of general relativity, to name a few. In such settings, the high-fidelity signal computed from differential equations is often too slow to be directly used.

The Journal of Open Source Software



Figure 1: Example gravitational wave prediction from a surrogate model compared with numerical relativity for a precessing binary black hole system. This particular numerical relativity simulation took 70,881 CPU-hours (about 1.75 months using 56 cores on the supercomputer Frontera), while the surrogate model can be evaluated in about 100 milliseconds.

Surrogate models offer a practical way to dramatically accelerate model evaluation while retaining the high-fidelity accuracy of the expensive simulation code; an example is shown in Fig. 1. Surrogate models can be constructed in various ways, but what separates these models from other modeling frameworks is that they are primarily data-driven. Given a training set of gravitational waveform data sampling the parameter space, a model is built by following three steps:

1. Feature extraction: the waveform is decomposed into *data pieces* that are simple to model,
2. Dimensionality reduction: each data piece is approximated by a low-dimensional vector space, which reduces the degrees of freedom we need to model, and
3. Regression: fitting and regression techniques are applied to the low-dimensional representation of each data piece over the parameter space defining the model.

These model-building steps result in an HDF5 file defining the surrogate model's data and structure, which is stored on Zenodo. The GWSurrogate package provides access to these HDF5 files through its catalog interface, and all available models and their HDF5 files can be found in `gwsurrogate.catalog.list()`. For a recent overview of surrogate modeling as used in gravitational wave astrophysics, please see Section 5 of Afshordi & others (2023).

The development of `GWSurrogate` is hosted on GitHub and distributed through both PyPI and Conda. Quick start guides are found on the project's homepage while model-specific documentation is described through a collection of model-specific Jupyter notebooks. Automated testing is run on GitHub Actions.

Statement of need

`GWSurrogate` is a Python package that provides an easy to use interface to gravitational wave surrogate models. More specifically, `GWSurrogate` gravitational wave models provide evaluation of

$$h_{\rm S}(t,\theta,\phi;\Lambda) = \sum_{\ell=2}^{\infty}\sum_{m=-\ell}^{\ell} h_{\rm S}^{\ell m}(t;\Lambda)\ {}^{-2}Y_{\ell m}(\theta,\phi)\,,$$

where ${}^{-2}Y_{\ell m}$ are the spin$=-2$ weighted spherical harmonics and Λ describes the model's parameterization. The surrogate model provides fast evaluations for the modes, $h_{\rm S}^{\ell m}$. As described more fully in the documentation, the high-level API allows users direct access to the modes $\{h_{\rm S}^{\ell m}(t)\}$ (as a Python dictionary) or assembles the sum $h_{\rm S}(t,\theta,\phi)$ at a particular

DRAFT

location (θ, ϕ). The models implemented in `GWSurrogate` are intended to be used in production data analysis efforts. As such,

- computationally expensive operations (e.g., interpolation onto uniform time grids) are implemented by wrapping low-level C code for speed, whereas `GWSurrogate` provides a user-friendly interface to the high-level waveform evaluation API,
- models implemented in `GWSurrogate` follow the waveform convention choices of the LIGO-Virgo-Kagra collaboration, thus ensuring that downstream data analysis codes can use `GWSurrogate` models without needing to worry about different conventions, and
- `GWSurrogate` models can be directly evaluated in either physical units (often used in data analysis studies) and dimensionless units (often used in theoretical studies) where all dimensioned quantities are expressed in terms of the system's total mass.

Currently, there are 15 supported surrogate models (Blackman et al. (2015), O'Shaughnessy et al. (2017), Blackman et al. (2017), Varma, Field, Scheel, Blackman, Kidder, et al. (2019), Yoo et al. (2023), Yoo et al. (2022), Varma, Field, Scheel, Blackman, Gerosa, et al. (2019), Barkett et al. (2020), Rifat et al. (2020), Islam et al. (2022), Field et al. (2014)). These models vary in their duration, included physical effects (e.g. nonlinear memory, tidal forces, harmonic modes retained, eccentricity, mass ratio extent, precession effects, etc), and underlying solution method (e.g. Effective One Body, numerical relativity, and black hole perturbation theory). Details about all models can be found by doing `gwsurrogate.catalog.list(verbose=True)`, while the `GWSurrogate` homepage summarizes the state-of-the-art models for each particular problem. Certain models allow for additional functionality such as returning the dynamics of the binary black hole; these special features are described further in model-specific example notebooks.

Acknowledgements

We acknowledge our many close collaborators for their contribution to the development of surrogate models. We further acknowledge the community of `GWSurrogate` users who have contributed pull requests and opened issues, including Kevin Barkett, Mike Boyle, Collin Capano, Dwyer Deighan, Raffi Enficiaud, Oliver Jennrich, Gaurav Khanna, Duncan Macleod, Alex Nitz, Seth Olsen, Swati Singh, and Avi Vajpeyi. `GWSurrogate` has been developed over the past 10 years with continued support from the National Science Foundation, most recently through NSF grants PHY-2110496, PHY-2309301, and DMS-2309609. This work was also supported in part by the Sherman Fairchild Foundation, by NSF Grants PHY-2207342 and OAC-2209655 at Cornell, and by NSF Grants PHY-2309211, PHY-2309231, and OAC-2209656 at Caltech.

References

Afshordi, N., & others. (2023). *Waveform Modelling for the Laser Interferometer Space Antenna*. https://arxiv.org/abs/2311.01300

Barkett, K., Chen, Y., Scheel, M. A., & Varma, V. (2020). Gravitational waveforms of binary neutron star inspirals using post-Newtonian tidal splicing. *Phys. Rev. D*, *102*(2), 024031. https://doi.org/10.1103/PhysRevD.102.024031

Blackman, J., Field, S. E., Galley, C. R., Szilágyi, B., Scheel, M. A., Tiglio, M., & Hemberger, D. A. (2015). Fast and Accurate Prediction of Numerical Relativity Waveforms from Binary Black Hole Coalescences Using Surrogate Models. *Phys. Rev. Lett.*, *115*(12), 121102. https://doi.org/10.1103/PhysRevLett.115.121102

Blackman, J., Field, S. E., Scheel, M. A., Galley, C. R., Hemberger, D. A., Schmidt, P., & Smith, R. (2017). A Surrogate Model of Gravitational Waveforms from Numerical

DRAFT

Relativity Simulations of Precessing Binary Black Hole Mergers. *Phys. Rev. D*, *95*(10), 104023. https://doi.org/10.1103/PhysRevD.95.104023

Field, S. E., Galley, C. R., Hesthaven, J. S., Kaye, J., & Tiglio, M. (2014). Fast prediction and evaluation of gravitational waveforms using surrogate models. *Phys. Rev. X*, *4*(3), 031006. https://doi.org/10.1103/PhysRevX.4.031006

Islam, T., Field, S. E., Hughes, S. A., Khanna, G., Varma, V., Giesler, M., Scheel, M. A., Kidder, L. E., & Pfeiffer, H. P. (2022). Surrogate model for gravitational wave signals from nonspinning, comparable-to large-mass-ratio black hole binaries built on black hole perturbation theory waveforms calibrated to numerical relativity. *Phys. Rev. D*, *106*(10), 104025. https://doi.org/10.1103/PhysRevD.106.104025

O'Shaughnessy, R., Blackman, J., & Field, S. E. (2017). An architecture for efficient gravitational wave parameter estimation with multimodal linear surrogate models. *Class. Quant. Grav.*, *34*(14), 144002. https://doi.org/10.1088/1361-6382/aa7649

Rifat, N. E. M., Field, S. E., Khanna, G., & Varma, V. (2020). Surrogate model for gravitational wave signals from comparable and large-mass-ratio black hole binaries. *Phys. Rev. D*, *101*(8), 081502. https://doi.org/10.1103/PhysRevD.101.081502

Varma, V., Field, S. E., Scheel, M. A., Blackman, J., Gerosa, D., Stein, L. C., Kidder, L. E., & Pfeiffer, H. P. (2019). Surrogate models for precessing binary black hole simulations with unequal masses. *Phys. Rev. Research.*, *1*, 033015. https://doi.org/10.1103/PhysRevResearch.1.033015

Varma, V., Field, S. E., Scheel, M. A., Blackman, J., Kidder, L. E., & Pfeiffer, H. P. (2019). Surrogate model of hybridized numerical relativity binary black hole waveforms. *Phys. Rev. D*, *99*(6), 064045. https://doi.org/10.1103/PhysRevD.99.064045

Yoo, J., Mitman, K., Varma, V., Boyle, M., Field, S. E., Deppe, N., Hébert, F., Kidder, L. E., Moxon, J., Pfeiffer, H. P., & others. (2023). Numerical relativity surrogate model with memory effects and post-Newtonian hybridization. *Phys. Rev. D*, *108*(6), 064027. https://doi.org/10.1103/PhysRevD.108.064027

Yoo, J., Varma, V., Giesler, M., Scheel, M. A., Haster, C.-J., Pfeiffer, H. P., Kidder, L. E., & Boyle, M. (2022). Targeted large mass ratio numerical relativity surrogate waveform model for GW190814. *Phys. Rev. D*, *106*(4), 044001. https://doi.org/10.1103/PhysRevD.106.044001

DRAFT