

SEPARATE: Storm Event Partitioning And Rainfall Analytics for Tipping-bucket rain gauge data Evaluation

Brendan P. Murphy ^{1*} and Scott R. David ^{2*}

¹ School of Environmental Science, Simon Fraser University, Canada ² Department of Watershed Sciences, Utah State University, United States of America  Corresponding author * These authors contributed equally.

DOI: [10.21105/joss.08851](https://doi.org/10.21105/joss.08851)

Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Editor: [Taher Chegini](#) 

Reviewers:

- [@jrkasprzyk](#)
- [@dkedar7](#)

Submitted: 11 April 2025

Published: 23 March 2026

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

Summary

Here, we present SEPARATE: an open-source, GUI-based software package designed to provide users with a fast, reliable, and automated means to post-process tipping bucket rain gauge (TBRG) records. SEPARATE stands for ‘Storm Event Partitioning And Rainfall Analytics for Tipping-bucket rain gauge data Evaluation’. The software identifies and partitions storm events using the widely accepted fixed minimum inter-event time (MIT) method ([D. Dunkerley, 2008](#); [Restrepo-Posada & Eagleson, 1982](#); [Staley et al., 2013](#)), analyzes rainfall metrics for each storm, and provides organized outputs with metadata and visualization options. Detailed descriptions of the methods can be found in the supplement and user manual documents in the GitHub repository.



Figure 1: Application logo designed by B. Murphy.

The complete open-source code is included as part of this publication, but our Python-based tool has also been packaged into a convenient GUI-based application for both Microsoft Windows and macOS users. The downloadable installer includes all dependencies, allowing SEPARATE to be installed and run without any additional installations. The GUI was developed using PySimpleGUI v4.60.4 ([B., 2024](#)), compiled with PyInstaller v6.3.0 ([Hartley & PyInstaller Development Team, 2024](#)), and packaged using Inno Setup v6.2.2 ([Laan, 2024](#)) for Windows and create-dmg for macOS ([Tarantsov & contributors, 2024](#)).




Figure 2: When using the installer for Windows or MacOS, SEPARATE runs as a stand-alone desktop application with the graphical user interface (GUI) shown here. All data inputs and parameter selection are managed through drop-down menus, checkboxes, etc., and the tool is initialized by clicking 'Separate Storms'. A progress bar and pop-up window allow users to know when their analysis is complete.

During installation (56 MB for Windows; 171 MB for Mac), users select a directory and optionally add a shortcut to their Desktop (Figure 1). Upon launching the application, users are presented with a GUI organized into three sections (Inputs, Partitioning Criteria, and Output Options) with data entry boxes, buttons, checkboxes, and drop-down menus (Figure 2). SEPARATE requires no programming environment to run and provides a step-by-step process to input rainfall records and then select all needed parameters. If any required inputs are missing or conflicting options are selected, error messages will appear. Tooltip buttons in the GUI open pop-ups with explanations, and a detailed instruction manual (.pdf) is included with the download. Additional methodological details are provided in a supplementary file.

Statement of Need

Accurate characterization of rainfall data is essential in environmental research. Assessing characteristics of independent rainfall events (“storms”) is critical for many applications,

including but not limited to the evaluation of rainfall-runoff thresholds (Kean et al., 2011; Staley et al., 2013), hydroclimatic dynamics (Canham et al., 2025; Slater et al., 2021), sediment transport (DeLong et al., 2018; Dick et al., 1997), soil erosion (D. L. Dunkerley, 2019; Robichaud et al., 2008), watershed sediment dynamics (Murphy et al., 2019), natural hazards (Cannon et al., 2010; Gartner et al., 2014; McGuire et al., 2021; Peres & Cancelliere, 2014; Rengers et al., 2024; Staley et al., 2017), landscape evolution (Tucker & Bras, 2000), and water resource management (Ward et al., 2011). Storm metrics of interest vary by application but often include time of arrival, duration, magnitude, or peak rainfall rates (Collar et al., 2025). While some regions have high-fidelity rainfall data, this is not true worldwide. Even in areas with more robust data coverage, remote sensing techniques, such as radar, perform poorly in steep, complex terrain. Yet, high-gradient landscapes are the focus for lots of hydrogeomorphic research, especially natural hazards. Consequently, ground-based instrumentation is still heavily relied upon (Segoni et al., 2018).

The tipping bucket rain gauge (TBRG) remains one of the most widely used instruments, given its simplicity, ruggedness, affordability, and ability to provide continuous rainfall records. However, TBRG's raw timestamped records must be post-processed to derive meaningful meteorological data. Most TBRG systems include only basic software for launching and downloading data, rarely offering post-processing features, and especially not for storm separation or analysis. Third-party software is also limited. Some open-access options offer event-based rainfall analysis, such as the EPA Storm Water Management Model (SWMM; (Rossman, 2015)) or NetSTORM (Heineman, 2004), however they are often embedded within larger software packages, making them difficult to operationalize for storm separation alone. Open-source programmatic tools also exist for TBRG processing, such as the USGS Rainmaker R package (Corsi & Carvin, 2023), but these options represent a barrier to entry to anyone without programming experience.

A review by Segoni et al. (2018) of 107 recent landslide studies indirectly revealed how software limitations may be critically affecting research quality and reproducibility. While nearly 80% of studies relied on rain gauges, none reportedly used existing software to analyze their rainfall data. The majority relied on manual processing and extraction and another 20% developed purpose-built (but largely unpublished) scripts. One reviewed study even noted how manual rainfall data processing “is a hard and time-consuming activity” (Giannecchini et al., 2016). Troublingly, Segoni et al. (2018) also reported that 13% of studies relied on “expert judgement” alone to analyze rainfall records. Further, Dunkerley (D. Dunkerley, 2008) (in a different review) noted that few researchers provide any basis for selected parameters when separating storms in continuous rainfall records. Without accessible, easy-to-use software, researchers and practitioners who often rely heavily on rainfall data, but may not have specific expertise in hydrometrics, often appear to be relying on manual, custom, undocumented, and/or subjective methods for analyzing rain gauge data. SEPARATE was designed to help fill and address this critical gap.

References

- B., M. (2024). *PySimpleGUI: Python GUIs for humans*. PyPI. <https://pypi.org/project/PySimpleGUI/>
- Canham, H. A., Lane, B., Phillips, C. B., & Murphy, B. P. (2025). Leveraging a time-series event separation method to disentangle time-varying hydrologic controls on streamflow—application to wildfire-affected catchments. *Hydrology and Earth System Sciences*, 29(1), 27–43. <https://doi.org/10.5194/hess-29-27-2025>
- Cannon, S. H., Gartner, J. E., Rupert, M. G., Michael, J. A., Rea, A. H., & Parrett, C. (2010). Predicting the probability and volume of post wildfire debris flows in the intermountain western united states. *Bulletin*, 122(1-2), 127–144. <https://doi.org/10.1130/b26459.1>
- Collar, N. M., Moody, J. A., & Ebel, B. A. (2025). Rainfall as a driver of post-wildfire

- flooding and debris flows: A review and synthesis. *Earth-Science Reviews*, 260, 104990. <https://doi.org/10.1016/j.earscirev.2024.104990>
- Corsi, S., & Carvin, R. (2023). *Rainmaker: Rainfall event processing*. <https://code.usgs.gov/water/analysis-tools/Rainmaker>
- DeLong, S. B., Youberg, A. M., DeLong, W. M., & Murphy, B. P. (2018). Post-wildfire landscape change and erosional processes from repeat terrestrial lidar in a steep headwater catchment, chiricahua mountains, arizona, USA. *Geomorphology*, 300, 13–30. <https://doi.org/10.1016/j.geomorph.2017.09.028>
- Dick, G. S., Anderson, R. S., & Sampson, D. E. (1997). Controls on flash flood magnitude and hydrograph shape, upper blue hills badlands, utah. *Geology*, 25(1), 45–48. [https://doi.org/10.1130/0091-7613\(1997\)025%3C0045:coffma%3E2.3.co;2](https://doi.org/10.1130/0091-7613(1997)025%3C0045:coffma%3E2.3.co;2)
- Dunkerley, D. (2008). Identifying individual rain events from pluviograph records: A review with analysis of data from an australian dryland site. *Hydrological Processes*, 22(26), 5024–5036. <https://doi.org/10.1002/hyp.7122>
- Dunkerley, D. L. (2019). Rainfall intensity bursts and the erosion of soils: An analysis highlighting the need for high temporal resolution rainfall data for research under current and future climates. *Earth Surface Dynamics*, 7(2), 345–360. <https://doi.org/10.5194/esurf-7-345-2019>
- Gartner, J. E., Cannon, S. H., & Santi, P. M. (2014). Empirical models for predicting volumes of sediment deposited by debris flows and sediment-laden floods in the transverse ranges of southern california. *Engineering Geology*, 176, 45–56. <https://doi.org/10.1016/j.enggeo.2014.04.008>
- Gianecchini, R., Galanti, Y., Avanzi, G. D. A., & Barsanti, M. (2016). Probabilistic rainfall thresholds for triggering debris flows in a human-modified landscape. *Geomorphology*, 257, 94–107. <https://doi.org/10.1016/j.geomorph.2015.12.012>
- Hartley, G., & PyInstaller Development Team, the. (2024). *PyInstaller: Freeze python applications into standalone executables*. PyInstaller.org. <https://www.pyinstaller.org/>
- Heineman, M. C. (2004). NetSTORM-a computer program for rainfall-runoff simulation and precipitation analysis. *Critical Transitions in Water and Environmental Resources Management*, 1–14. [https://doi.org/10.1061/40737\(2004\)395](https://doi.org/10.1061/40737(2004)395)
- Kean, J. W., Staley, D. M., & Cannon, S. H. (2011). In situ measurements of post-fire debris flows in southern california: Comparisons of the timing and magnitude of 24 debris-flow events with rainfall and soil moisture conditions. *Journal of Geophysical Research: Earth Surface*, 116(F4). <https://doi.org/10.1029/2011jf002005>
- Laan, M. (2024). *Inno setup: Script-based installation system created in delphi*. <https://jrsoftware.org/isinfo.php>
- McGuire, L. A., Rengers, F. K., Oakley, N., Kean, J. W., Staley, D. M., Tang, H., Orla-Barile, M. de, & Youberg, A. M. (2021). Time since burning and rainfall characteristics impact post-fire debris-flow initiation and magnitude. *Environmental & Engineering Geoscience*, 27(1), 43–56. <https://doi.org/10.2113/eeg-d-20-00029>
- Murphy, B. P., Czuba, J. A., & Belmont, P. (2019). Post-wildfire sediment cascades: A modeling framework linking debris flow generation and network-scale sediment routing. *Earth Surface Processes and Landforms*, 44(11), 2126–2140. <https://doi.org/10.1002/esp.4635>
- Peres, D. J., & Cancelliere, A. (2014). Derivation and evaluation of landslide-triggering thresholds by a monte carlo approach. *Hydrology and Earth System Sciences*, 18(12), 4913–4931. <https://doi.org/10.5194/hess-18-4913-2014>

- Rengers, F. K., Bower, S., Knapp, A., Kean, J. W., vonLembke, D. W., Thomas, M. A., Kostelnik, J., Barnhart, K. R., Bethel, M., Gartner, J. E., Hille, M., Staley, D. M., Anderson, J. K., Roberts, E. K., DeLong, S. B., Lane, B., Ridgway, P., & Murphy, B. P. (2024). Evaluating post-wildfire debris-flow rainfall thresholds and volume models at the 2020 grizzly creek fire in glenwood canyon, colorado, USA. *Natural Hazards and Earth System Sciences*, 24(6), 2093–2114. <https://doi.org/10.5194/nhess-24-2093-2024>
- Restrepo-Posada, P. J., & Eagleson, P. S. (1982). Identification of independent rainstorms. *Journal of Hydrology*, 55(1-4), 303–319. [https://doi.org/10.1016/0022-1694\(82\)90136-6](https://doi.org/10.1016/0022-1694(82)90136-6)
- Robichaud, P. R., Pierson, F. B., Brown, R. E., & Wagenbrenner, J. W. (2008). Measuring effectiveness of three postfire hillslope erosion barrier treatments, western montana, USA. *Hydrological Processes: An International Journal*, 22(2), 159–170. <https://doi.org/10.1002/hyp.6558>
- Rossman, L. A. (2015). *Storm water management model user's manual, version 5.1 (EPA-600/r-14/413b)*. National Risk Management Research Laboratory, Office of Research; Development, US Environmental Protection Agency.
- Segoni, S., Piciullo, L., & Gariano, S. L. (2018). A review of the recent literature on rainfall thresholds for landslide occurrence. *Landslides*, 15(8), 1483–1501. <https://doi.org/10.1007/s10346-018-0966-4>
- Slater, L. J., Anderson, B., Buechel, M., Dadson, S., Han, S., Harrigan, S., Kelder, T., Kowal, K., Lees, T., Matthews, T., Murphy, C., & Wilby, R. L. (2021). Nonstationary weather and water extremes: A review of methods for their detection, attribution, and management. *Hydrology and Earth System Sciences*, 25(7), 3897–3935. <https://doi.org/10.5194/hess-25-3897-2021>
- Staley, D. M., Kean, J. W., Cannon, S. H., Schmidt, K. M., & Laber, J. L. (2013). Objective definition of rainfall intensity–duration thresholds for the initiation of post-fire debris flows in southern california. *Landslides*, 10, 547–562. <https://doi.org/10.1007/s10346-012-0341-9>
- Staley, D. M., Negri, J. A., Kean, J. W., Laber, J. L., Tillery, A. C., & Youberg, A. M. (2017). Prediction of spatially explicit rainfall intensity–duration thresholds for post-fire debris-flow generation in the western united states. *Geomorphology*, 278, 149–162. <https://doi.org/10.1016/j.geomorph.2016.10.019>
- Tarantsov, A., & contributors. (2024). *Create-dmg: A shell script to build fancy DMG images on macOS*. <https://github.com/create-dmg/create-dmg>.
- Tucker, G. E., & Bras, R. L. (2000). A stochastic approach to modeling the role of rainfall variability in drainage basin evolution. *Water Resources Research*, 36(7), 1953–1964. <https://doi.org/10.1029/2000wr900065>
- Ward, E., Buytaert, W., Peaver, L., & Wheeler, H. (2011). Evaluation of precipitation products over complex mountainous terrain: A water resources perspective. *Advances in Water Resources*, 34(10), 1222–1231. <https://doi.org/10.1016/j.advwatres.2011.05.007>