The Journal of Open Source Software

anadrofish: R package for modeling anadromous fish population responses to dams

Daniel S. Stich [1,2¶], **Joshua D. Hardesty** [1], **Nicholas T. Jordan** [1], **Samuel G. Roy** [3], **Timothy F. Sheehan** [4], **Shawn D. Snyder** [5], and **Joseph D. Zydlewski** [6,5]

1 Biology Department, State University of New York at Oneonta, NY 13280 USA **2** Biological Field Station, State University of New York at Oneonta, Cooperstown, NY 13326 USA **3** Maine Geological Survey, Department of Agriculture, Conservation and Forestry, Augusta, ME, 04333 USA **4** National Oceanic and Atmospheric Administration, National Marine Fisheries Service, Northeast Fisheries Science Center, Woods Hole, MA 02543 USA **5** Department of Wildlife, Fisheries, and Conservation Biology, University of Maine, Orono, ME, 04469 USA **6** U.S. Geological Survey Maine Cooperative Fish and Wildlife Research Unit, Orono, ME, 04469 USA ¶ Corresponding author

DOI: 10.xxxxxx/draft

Software

- Review
- Repository
- Archive

Editor:

Submitted: 09 June 2025
Published: unpublished

License
Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License (CC BY 4.0).

DRAFT

USGS Disclaimer

This draft manuscript is distributed solely for the purpose of peer review. Its content is deliberative and predecisional, so it must not be disclosed or released by reviewers. Because the manuscript has not yet been approved for publication by the U.S. Geological Survey (USGS), it does not represent any official USGS finding or policy.

Summary

Diadromous fishes world-wide experienced precipitous declines during the 19th and 20th centuries due to a combination of overfishing, pollution, and freshwater habitat loss through construction of dams (Limburg & Waldman, 2009). Following wide-spread fishing closures and large-scale remediation of many historical pollution sources, dams in coastal rivers remain as the largest tractable impediment to population recovery for many of these species (Waldman & Quinn, 2022). In some cases, dams reduce access to as much as 95% of freshwater and rearing habitat (Hall et al., 2011). These effects are especially pronounced for species that rely on long-distance migrations to spawning and rearing habitat upstream of barriers such as various alosines (herrings; e.g., American shad *Alosa sapidissima*, alewife *A. pseudoharengus*, and blueback herring *A. aestivalis* (Noonan et al., 2012)) and salmonines (trout and salmon; e.g., *Salmo* spp. (Parrish et al., 1998) and *Oncorhynchus* spp. (Quiñones et al., 2015).

Because of their complex life-histories and use of discrete habitats within ontogenetic stages and seasons, traditional stock assessment tools often do not suffice for population assessment. This has historically necessitated species- and system-specific approaches to fisheries stock assessment and management strategy evaluation (Barber et al., 2018; e.g., Nieland et al., 2015; Roy et al., 2018; Stich et al., 2019). Therefore, we created the `anadrofish` package (Stich, Hardesty, et al., 2025) for R (R Core Team, 2025) to provide a generalized approach that also allows broader application to novel species, systems, and scenarios.

Statement of need

`anadrofish` is an R package that can be used to model anadromous fish populations, including responses to dams and other fishery management considerations (e.g., fishery harvest rates).

We created it for use by fisheries scientists and managers interested in development of life-history theory, stock assessment, or decision making. The R package was developed for use in the 2020 Atlantic States Marine Fisheries Commission (ASMFC) American shad benchmark stock assessment (Atlantic States Marine Fisheries Commission, 2020) and the 2024 ASMFC river herring benchmark stock assessment (Atlantic States Marine Fisheries Commission, 2024). It has also been used recently to answer coast-wide ecological research questions (Zydlewski et al., 2021) and evaluate local or regional fishery management scenarios.

The package includes a variety of helper functions for simulating life-history events and population dynamics as well as built-in data sets from empirical studies and population assessments for the native North American range of multiple species of fish from Florida, USA through maritime Canada. It generalizes similar routines implemented in other species- or population-specific modeling tools (Barber et al., 2018; Stich et al., 2019; Stich, Nieland, et al., 2025), is faster than other empirical life-history models (e.g., Stich & Gilligan-Lunda, 2024), and it can be modified to use outputs of common empirical fisheries analysis from other R packages such as `fishStan` (Erikson et al., 2022) or `FSA` (Ogle et al., 2025). The package can also be used to simulate new populations in arbitrary watersheds. Default outputs are readily integrated into tidy workflows through packages in the `tidyverse` (Wickham et al., 2019), including plotting and visualization of results in `ggplot2` (Wickham, 2016).

The `sim_pop()` function provides high-level functionality for end users interested in investigating theoretical questions or applied population management scenarios. This function links various helper functions and built-in datasets to simulate populations through time and space. By default, it includes functionality for 169 American shad, 222 alewife, and 238 blueback herring populations. Additionally, optional arguments to this function such as `custom_habitat`, and corresponding `custom_habitat_template()` helper function provide customization options or ability to extend to arbitrarily defined scales or novel systems. This flexibility has the potential to broaden the application of these modeling approaches to novel species and geographies or restoration scenarios for management strategy evaluation. The next steps for this project will include addition of generalized homing and straying routines that can be used to customize probability of fish migration to specific portions of empirical or derived watersheds, and creation of methods that allow for multiyear freshwater residence to accommodate a wider range of life histories, including catadromous fishes that grow in freshwater and spawn in the marine environment.

Acknowledgements

We acknowledge contributions from M. Bailey, E. Gilligan-Lunda, J. Kipp, and K. Sprankle during initial package development for American shad, and contributions by K. Drew, W. Eakin, and B. Gahagan to package updates for alewife and blueback herring. We thank numerous ASMFC Technical Committee representatives and regional biologists for input during coast-wide stock assessments. This work was funded by the NOAA National Marine Fisheries Service Northeast Fisheries Science Center (award number NMFS-NEFSC-199 MOU), with additional support from the State University of New York Oneonta Biological Field Station, in kind support from the U.S. Geological Survey Maine Cooperative Fish and Wildlife Research Unit, and the University of Maine. We thank the reviewers for helpful guidance and comments that greatly improved the quality of this paper and the open-source software package. Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

References

Atlantic States Marine Fisheries Commission. (2020). *American shad benchmark stock assessment and peer-review report - 2020*. Atlantic States Marine Fisheries Commission.

https://asmfc.org/wp-content/uploads/2025/01/AmShadBenchmarkStockAssessment_PeerReviewReport_2020_web.pdf

Atlantic States Marine Fisheries Commission. (2024). *River herring benchmark stock assessment and peer-review report - 2024*. Atlantic States Marine Fisheries Commission. https://asmfc.org/wp-content/uploads/2025/01/RiverHerringAssessment_PeerReviewReport_2024.pdf

Barber, B. L., Gibson, A. J., O'Malley, A. J., & Zydlewski, J. (2018). Does what goes up also come down? Using a recruitment model to balance alewife nutrient import and export. *Marine and Coastal Fisheries*, *10*(2), 236–254. https://doi.org/10.1002/mcf2.10021

Erikson, R. A., Stich, D. S., & Hebert, J. L. (2022). fishStan: Hierarchical Bayesian models for fisheries. *Journal of Open Source Software*, *7*(71), 3444. https://doi.org/10.21105/joss.03444

Hall, C. J., Jordaan, A., & Frisk, M. G. (2011). The historic influence of dams on diadromous fish habitat with a focus on river herring and hydrologic longitudinal connectivity. *Landscape Ecology*, *26*(1), 95–107. https://doi.org/10.1525/bio.2009.59.11.7

Limburg, K. E., & Waldman, J. R. (2009). Dramatic declines in North Atlantic diadromous fishes. *BioScience*, *59*(11), 955–965. https://doi.org/10.1525/bio.2009.59.11.7

Nieland, J. L., Sheehan, T. F., & Saunders, R. (2015). Assessing demographic effects of dams on diadromous fish: A case study for Atlantic salmon in the Penobscot River, Maine. *International Council for the Exploration of the Sea Journal of Marine Science*, *72*(8), 2423–2437. https://doi.org/10.1093/icesjms/fsv083

Noonan, M. J., Grant, J. W., & Jackson, C. D. (2012). A quantitative assessment of fish passage efficiency. *Fish and Fisheries*, *13*(4), 450–464. https://doi.org/doi.org/10.1111/j.1467-2979.2011.00445.x

Ogle, D. H., Doll, J. C., Powell, W., & Dinno, A. (2025). *FSA: Fisheries stock analysis*. https://doi.org/10.5281/zenodo.597719

Parrish, D. L., Behnke, R. J., Gephard, S. R., McCormick, S. D., & Reeves, G. H. (1998). Why aren't there more Atlantic salmon (*Salmo salar*)? *Canadian Journal of Fisheries and Aquatic Sciences*, *55*(S1), 281–287. https://doi.org/10.1139/d98-012

Quiñones, R. M., Grantham, T. E., Harvey, B. N., Kiernan, J. D., Klasson, M., Wintzer, A. P., & Moyle, P. B. (2015). Dam removal and anadromous salmonid (*oncorhynchus* spp.) conservation in California. *Reviews in Fish Biology and Fisheries*, *25*, 195–215. https://doi.org/10.1007/s11160-014-9359-5

R Core Team. (2025). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing. https://www.R-project.org/

Roy, S. G., Uchida, E., de Souza, S. P., Blachly, B., Fox, E., Gardner, K., Gold, A. J., Jansujwicz, J. S., Klein, McGreavy, B., Mo, W., Smith, S. M. C., Vogler, K., E. Wilson, Zydlewski, J., & D., H. (2018). A multiscale approach to balance trade-offs among dam infrastructure, river restoration, and cost. *Proceedings of the National Academy of Sciences*, *115*(47), 12069–12074. https://doi.org/10.1073/pnas.1807437115

Stich, D. S., & Gilligan-Lunda, E. (2024). *shadia: Dam passage performance standard models for R*. https://doi.org/10.5281/zenodo.13375033

Stich, D. S., Hardesty, J. D., Jordan, N. T., Roy, S. G., Sheehan, T. F., Snyder, S. D., & Zydlewski, J. D. (2025). *Anadrofish: Anadromous fish population responses to habitat changes*. https://doi.org/https://doi.org/10.5281/zenodo.14285894

Stich, D. S., Nieland, J. L., & Sheehan, T. F. (2025). dia: An R package for the National Oceanic and Atmospheric Administration dam impact analysis. *Journal of Open Source*

DRAFT

Software, *10*(105), 7475. https://doi.org/10.21105/joss.07475

Stich, D. S., Sheehan, T. F., & Zydlewski, J. D. (2019). A dam passage performance standard model for American shad. *Canadian Journal of Fisheries and Aquatic Sciences*, *76*(5), 762–779. https://doi.org/10.1139/cjfas-2018-0008

Waldman, J. R., & Quinn, T. P. (2022). North American diadromous fishes: Drivers of decline and potential for recovery in the Anthropocene. *Science Advances*, *8*(4), eabl5486. https://doi.org/10.1126/sciadv.abl5486

Wickham, H. (2016). *ggplot2: Elegant graphics for data analysis*. Springer-Verlag New York. https://doi.org/10.32614/CRAN.package.ggplot2

Wickham, H., Averick, M., Bryan, J., Chang, W., D'Agostino McGowan, L., François, R., Grolemund, G., Hayes, A., Henry, L., Hester, J., Kuhn, M., Lin Pedersen, T., Miller, E., Bache, S. M., Müller, Ooms, J., Robinson, D., Seidel, D. P., Spinu, V., … Yutani, H. (2019). Welcome to the tidyverse. *Journal of Open Source Software*, *4*(43), 1686. https://doi.org/10.21105/joss.01686

Zydlewski, J., Stich, D. S., Roy, S., Bailey, M., Sheehan, T., & Sprankle, K. (2021). What have we lost? Modeling dam impacts on American shad populations through their native range. *Frontiers in Marine Science*, *8*. https://doi.org/10.3389/fmars.2021.734213

DRAFT