The Journal of Open Source Software

GASTLI: A Python package for coupled interior–atmosphere modelling of volatile-rich planets

Lorena Acuña [1¶], **Laura Kreidberg** [1], **Meng Zhai** [1,2], **Paul Mollière** [1], and **Morgan Fouesneau** [1]

1 Max Planck Institute for Astronomy, Königstuhl 17, 69117 Heidelberg, Germany **2** Chinese Academy of Sciences South America Center for Astronomy (CASSACA), National Astronomical Observatories, Chinese Academy of Sciences, Beijing 100101, PR China ¶ Corresponding author

DOI: 10.xxxxxx/draft

Software

- Review
- Repository
- Archive

Editor: Josh Borrow

Reviewers:

- @ivalaginja
- @MartianColonist

Submitted: 08 August 2024
Published: unpublished

License
Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License (CC BY 4.0).

Summary

Gaseous exoplanets are far more diverse in mass, density and temperature than their Solar System counterparts. These range from temperate sub-Neptunes whose masses and sizes are below that of the Solar System ice giants, to extremely irradiated, Jupiter-mass exoplanets known as hot Jupiters. Their bulk composition is key to understanding their formation and evolution pathways, including their dominant formation mechanism (Thorngren et al., 2016) and interactions between the irradiation of the host star and the planetary atmosphere (Owen & Wu, 2013). Their building blocks consist of rocks and water (metals), and hydrogen and helium (H/He), which are present in a wide variety of mass fractions in the exoplanet population. For exoplanets that contain H/He, the planetary radius (or density) not only depends on mass, but also on the temperature due to the distance from its host star, its age, the mass of the core, and the fraction of H/He in its envelope. The inference of bulk composition from observational data – mass, radius, equilibrium temperature, age and atmospheric metallicity – requires non-trivial numerical methods that solve the equations of planetary interior structure. Additionally, interior models need to incorporate the effects of processes occurring in the planet atmosphere, such as radiative transfer, and the thermodynamical properties of the different materials, provided by the equation of state (EOS).

`GASTLI` (GAS gianT modeL for Interiors) is a Python package for interior structure modelling of volatile-rich exoplanets. The package combines a FORTRAN back end with a Python-based user interface. This enables fast convergence of the interior structure to a solution that fulfills the user input, while having a flexible and user-friendly interface. `GASTLI` was specifically designed to compute a wider range of combinations of core mass and H/He content in the envelope compared to existing open-source interior models. GASTLI also incorporates state-of-the-art thermodynamical data and EOS for rock, water and H/He. One of its key features is the coupling class, which couples the interior structure model to a grid of atmospheric temperature profiles to determine self-consistently the temperature of the outermost layer. GASTLI is released with a default atmospheric grid that assumes equilibrium chemistry and a clear atmosphere at varying internal temperatures computed with petitCODE (Mollière et al., 2015, 2017). Nonetheless, the flexible interface enables the user to use their own custom atmospheric grid, which is useful to explore the effect of the processes assumed in the upper atmosphere in the interior, such as the type of chemistry (equilibrium or disequilibrium), cloud implementations or 3D global circulation (GCM).

DRAFT

The Journal of Open Source Software

Statement of need

Interior structure codes are essential to carry out planet composition analyses and retrievals on mass and radius data. Two open-source interior structure models are currently available: MESA (Paxton et al., 2011, 2013, 2015, 2018, 2019) and MAGRATHEA (Huang et al., 2022). MESA is a well-known stellar interior structure software that has been applied to sub-Neptunes and gas giants (Chen & Rogers, 2016; Müller & Helled, 2021). The compositions of the envelope are limited to low metal contents due to the default EOSs, which is particularly limiting for planets with Neptune's mass or lower. Therefore, the implementation of alternative EOS in MESA are necessary, which is not straightforward for the user given the complexity of the code, and the input feature based on FORTRAN inlists. The computational time of one simple forward model is at least 1 minute. This renders MESA difficult to couple to atmospheric models, which can have a significant effect on the output of the interior calculations (Linder et al., 2019; Lopez & Fortney, 2014; Poser et al., 2019; Poser & Redmer, 2024). The second open-source planet interior model, MAGRATHEA, is a computationally fast code based on C++. Its layer structure (ideal gas EOS on top of liquid/ice water layer) makes its adaptation for applications to sub-Neptunes and gas giants a challenge to new users because these type of planets require the treatment of nonideal, high-pressure H/He envelopes.

The GASTLI Python package

The interior model module is contained in the `GASTLI.int_planet` class. The interior structure equations for hydrostatic equilibrium, Gauss's theorem for gravitational acceleration, convection, and mass conservation are solved by combining integration by the trapezoidal rule with an iteration scheme over all profiles (pressure, temperature, gravity and density). In each iteration, the trapezoidal integrations are carried for each point in a 1D grid that represents the radius, from the center of the planet to the outermost boundary of the top layer. The density is calculated by evaluating the EOS for rock (Lyon, 1992; Miguel et al., 2022), water (Mazevet et al., 2019) and H/He (Chabrier & Debras, 2021; Howard & Guillot, 2023) at the pressure and temperature conditions of each point in the spatial grid. The iteration sequence is considered to have converged when the difference between iterations is less than a given precision that can be modified by the user. The recommended and default value for this precision is 10^{-5}. The iteration scheme can also be stopped at a given number of iterations defined by the user, with a default of 30 maximum iterations, and a maximum possible value of 99 iterations. The boundary conditions to solve these equations include the top layer's pressure and temperature, which can be defined by the user. In addition to the profile convergence, the interior module ensures that the surface boundary conditions and the input mass are satisfied. A complete description of the interior module integration scheme can be found in Brugger (2018), and has been widely used (Acuña et al., 2021, 2022, 2023; Aguichine et al., 2021; Brugger et al., 2016, 2017; Mousis et al., 2020; Vivien et al., 2022). The new feature of the interior module includes the optimization of several of its routines, such as the trapezoidal integration and the evaluation of the EOS's density tables, and the parallelization of the calculation of the interior structure profiles with OpenMP in FORTRAN. One single computation of the interior module (function `calc_radius` in the `GASTLI.int_planet` class) for the default precision and number of iterations takes 1.89 seconds in an Apple M2 CPU, with parallelization on 8 cores (four cores operating at 3.49 GHz and the other four cores operating at 2.42 GHz).

The class `atm_models_interp` encloses the function that interpolates the atmospheric grid to obtain the temperature profile and interior boundary temperature. In addition, this class also provides the functions that calculate the atmospheric thickness and the atmosphere density profile. The latter is obtained by evaluating the EOS used by the interior module for H/He, and the AQUA water EOS (Haldemann et al., 2020). The atmospheric grid class and the interior module class are then combined in the `Coupling` class. This class self-consistently couples the interior module and the atmosphere grids to calculate the boundary temperature

DRAFT

of the outermost layer of the interior model. See Acuña et al. (2021) for a detailed description of the coupling algorithm, which has been used in previous work (Acuña et al., 2022, 2023; Aguichine et al., 2021). Finally, the class `Thermal_evolution` contains the functions necessary to evaluate a sequence of coupled interior-atmosphere models at different internal temperatures and solve the luminosity equation to obtain the internal temperatures and radii as a function of age (Fortney et al., 2013; Poser et al., 2019). The complete validation of GASTLI against mass-radius relationships obtained with interior models used in previous work (Fortney et al., 2013; Müller & Helled, 2021) can be found in Acuña et al. (2024).

Acknowledgements

We thank Yamila Miguel for sharing the reformatted EOS tables for silicate (dry sand) from the SESAME database. We acknowledge the support of the Data Science Group at the Max Planck Institute for Astronomy (MPIA) and especially Ivelina Momcheva for her invaluable assistance in optimizing the software and developing the GASTLI Python package. We thank Magali Deleuil, Olivier Mousis, Artyom Aguichine and Bastien Brugger for their contributions to the development of Marseille's Super-Earth Interior (MSEI) model, on which GASTLI's development is based.

References

Acuña, L., Deleuil, M., & Mousis, O. (2023). Interior-atmosphere modelling to assess the observability of rocky planets with JWST. *677*, A14. https://doi.org/10.1051/0004-6361/202245736

Acuña, L., Deleuil, M., Mousis, O., Marcq, E., Levesque, M., & Aguichine, A. (2021). Characterisation of the hydrospheres of TRAPPIST-1 planets. *647*, A53. https://doi.org/10.1051/0004-6361/202039885

Acuña, L., Kreidberg, L., Zhai, M., & Mollière, P. (2024). GASTLI. An open-source coupled interior-atmosphere model to unveil gas-giant composition. *688*, A60. https://doi.org/10.1051/0004-6361/202450559

Acuña, L., Lopez, T. A., Morel, T., Deleuil, M., Mousis, O., Aguichine, A., Marcq, E., & Santerne, A. (2022). Water content trends in K2-138 and other low-mass multi-planetary systems. *660*, A102. https://doi.org/10.1051/0004-6361/202142374

Aguichine, A., Mousis, O., Deleuil, M., & Marcq, E. (2021). Mass-Radius Relationships for Irradiated Ocean Planets. *914*(2), 84. https://doi.org/10.3847/1538-4357/abfa99

Brugger, B. (2018). *Structure interne et minéralogie des exoplanètes terrestres de faible masse* [PhD thesis]. http://www.theses.fr/2018AIXM0324/document

Brugger, B., Mousis, O., Deleuil, M., & Deschamps, F. (2017). Constraints on Super-Earth Interiors from Stellar Abundances. *850*(1), 93. https://doi.org/10.3847/1538-4357/aa965a

Brugger, B., Mousis, O., Deleuil, M., & Lunine, J. I. (2016). Possible Internal Structures and Compositions of Proxima Centauri b. *831*(2), L16. https://doi.org/10.3847/2041-8205/831/2/L16

Chabrier, G., & Debras, F. (2021). A New Equation of State for Dense Hydrogen-Helium Mixtures. II. Taking into Account Hydrogen-Helium Interactions. *917*(1), 4. https://doi.org/10.3847/1538-4357/abfc48

Chen, H., & Rogers, L. A. (2016). Evolutionary Analysis of Gaseous Sub-Neptune-mass Planets with MESA. *831*(2), 180. https://doi.org/10.3847/0004-637X/831/2/180

Fortney, J. J., Mordasini, C., Nettelmann, N., Kempton, E. M.-R., Greene, T. P., & Zahnle,

DRAFT

K. (2013). A Framework for Characterizing the Atmospheres of Low-mass Low-density Transiting Planets. *775*(1), 80. https://doi.org/10.1088/0004-637X/775/1/80

Haldemann, J., Alibert, Y., Mordasini, C., & Benz, W. (2020). AQUA: a collection of H_2O equations of state for planetary models. *643*, A105. https://doi.org/10.1051/0004-6361/202038367

Howard, S., & Guillot, T. (2023). Accounting for non-ideal mixing effects in the hydrogen-helium equation of state. *arXiv e-Prints*, arXiv:2302.07902. https://doi.org/10.48550/arXiv.2302.07902

Huang, C., Rice, D. R., & Steffen, J. H. (2022). MAGRATHEA: an open-source spherical symmetric planet interior structure code. *513*(4), 5256–5269. https://doi.org/10.1093/mnras/stac1133

Linder, E. F., Mordasini, C., Mollière, P., Marleau, G.-D., Malik, M., Quanz, S. P., & Meyer, M. R. (2019). Evolutionary models of cold and low-mass planets: cooling curves, magnitudes, and detectability. *623*, A85. https://doi.org/10.1051/0004-6361/201833873

Lopez, E. D., & Fortney, J. J. (2014). Understanding the Mass-Radius Relation for Sub-neptunes: Radius as a Proxy for Composition. *792*(1), 1. https://doi.org/10.1088/0004-637X/792/1/1

Lyon, S. P. (1992). Sesame: The los alamos national laboratory equation of state database. *Los Alamos National Laboratory Report LA-UR-92-3407*.

Mazevet, S., Licari, A., Chabrier, G., & Potekhin, A. Y. (2019). Ab initio based equation of state of dense water for planetary and exoplanetary modeling. *621*, A128. https://doi.org/10.1051/0004-6361/201833963

Miguel, Y., Bazot, M., Guillot, T., Howard, S., Galanti, E., Kaspi, Y., Hubbard, W. B., Militzer, B., Helled, R., Atreya, S. K., Connerney, J. E. P., Durante, D., Kulowski, L., Lunine, J. I., Stevenson, D., & Bolton, S. (2022). Jupiter's inhomogeneous envelope. *662*, A18. https://doi.org/10.1051/0004-6361/202243207

Mollière, P., van Boekel, R., Bouwman, J., Henning, Th., Lagage, P.-O., & Min, M. (2017). Observing transiting planets with JWST. Prime targets and their synthetic spectral observations. *600*, A10. https://doi.org/10.1051/0004-6361/201629800

Mollière, P., van Boekel, R., Dullemond, C., Henning, Th., & Mordasini, C. (2015). Model Atmospheres of Irradiated Exoplanets: The Influence of Stellar Parameters, Metallicity, and the C/O Ratio. *813*(1), 47. https://doi.org/10.1088/0004-637X/813/1/47

Mousis, O., Deleuil, M., Aguichine, A., Marcq, E., Naar, J., Aguirre, L. A., Brugger, B., & Gonçalves, T. (2020). Irradiated Ocean Planets Bridge Super-Earth and Sub-Neptune Populations. *896*(2), L22. https://doi.org/10.3847/2041-8213/ab9530

Müller, S., & Helled, R. (2021). Synthetic evolution tracks of giant planets. *507*(2), 2094–2102. https://doi.org/10.1093/mnras/stab2250

Owen, J. E., & Wu, Y. (2013). Kepler Planets: A Tale of Evaporation. *775*(2), 105. https://doi.org/10.1088/0004-637X/775/2/105

Paxton, B., Bildsten, L., Dotter, A., Herwig, F., Lesaffre, P., & Timmes, F. (2011). Modules for Experiments in Stellar Astrophysics (MESA). *192*(1), 3. https://doi.org/10.1088/0067-0049/192/1/3

Paxton, B., Cantiello, M., Arras, P., Bildsten, L., Brown, E. F., Dotter, A., Mankovich, C., Montgomery, M. H., Stello, D., Timmes, F. X., & Townsend, R. (2013). Modules for Experiments in Stellar Astrophysics (MESA): Planets, Oscillations, Rotation, and Massive Stars. *208*(1), 4. https://doi.org/10.1088/0067-0049/208/1/4

Paxton, B., Marchant, P., Schwab, J., Bauer, E. B., Bildsten, L., Cantiello, M., Dessart, L.,

Farmer, R., Hu, H., Langer, N., Townsend, R. H. D., Townsley, D. M., & Timmes, F. X. (2015). Modules for Experiments in Stellar Astrophysics (MESA): Binaries, Pulsations, and Explosions. *220*(1), 15. https://doi.org/10.1088/0067-0049/220/1/15

Paxton, B., Schwab, J., Bauer, E. B., Bildsten, L., Blinnikov, S., Duffell, P., Farmer, R., Goldberg, J. A., Marchant, P., Sorokina, E., Thoul, A., Townsend, R. H. D., & Timmes, F. X. (2018). Modules for Experiments in Stellar Astrophysics (MESA): Convective Boundaries, Element Diffusion, and Massive Star Explosions. *234*(2), 34. https://doi.org/10.3847/1538-4365/aaa5a8

Paxton, B., Smolec, R., Schwab, J., Gautschy, A., Bildsten, L., Cantiello, M., Dotter, A., Farmer, R., Goldberg, J. A., Jermyn, A. S., Kanbur, S. M., Marchant, P., Thoul, A., Townsend, R. H. D., Wolf, W. M., Zhang, M., & Timmes, F. X. (2019). Modules for Experiments in Stellar Astrophysics (MESA): Pulsating Variable Stars, Rotation, Convective Boundaries, and Energy Conservation. *243*(1), 10. https://doi.org/10.3847/1538-4365/ab2241

Poser, A. J., Nettelmann, N., & Redmer, R. (2019). The Effect of Clouds as an Additional Opacity Source on the Inferred Metallicity of Giant Exoplanets. *Atmosphere*, *10*(11), 664. https://doi.org/10.3390/atmos10110664

Poser, A. J., & Redmer, R. (2024). The effect of cloudy atmospheres on the thermal evolution of warm giant planets from an interior modelling perspective. *arXiv e-Prints*, arXiv:2402.19466. https://doi.org/10.48550/arXiv.2402.19466

Thorngren, D. P., Fortney, J. J., Murray-Clay, R. A., & Lopez, E. D. (2016). The Mass-Metallicity Relation for Giant Planets. *831*(1), 64. https://doi.org/10.3847/0004-637X/831/1/64

Vivien, H. G., Aguichine, A., Mousis, O., Deleuil, M., & Marcq, E. (2022). On the Stability of Low-mass Planets with Supercritical Hydrospheres. *931*(2), 143. https://doi.org/10.3847/1538-4357/ac66e2

DRAFT