The Journal of Open Source Software

KPMru.jl: A Julia Package for Kernel Polynomial Method in Condensed Matter Physics

Ang-Kun Wu [1,3], **Yixing Fu** [1], **Justin H. Wilson** [4,5], **and J. H. Pixley** [1,2¶]

1 Department of Physics and Astronomy, Center for Materials Theory, Rutgers University, Piscataway, New Jersey 08854, USA **2** Center for Computational Quantum Physics, Flatiron Institute, 162 5th Avenue, New York, New York 10010, USA **3** Department of Physics and Astronomy, University of Tennessee, Knoxville, Knoxville, Tennessee, 37996, USA **4** Department of Physics and Astronomy, Louisiana State University, Baton Rouge, LA 70803, USA **5** Center for Computation and Technology, Louisiana State University, Baton Rouge, LA 70803, USA ¶ Corresponding author

DOI: 10.xxxxxx/draft

Software

- Review
- Repository
- Archive

Editor:

Submitted: 02 March 2026
Published: unpublished

License
Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License (CC BY 4.0).

Summary

The Kernel Polynomial Method (KPM) is a numerical technique for approximating spectral densities and response functions of large Hermitian operators without performing full diagonalization (Weisse et al., 2006). KPM expands spectral functions in Chebyshev polynomials and evaluates them via recursive matrix-vector multiplications. Reconstructing the moments of the Chebyshev expansion depends on the observable of interest and can be computed through either a single inner product or through a trace, which is computed using a stochastic trace (Skilling, 1988). Carefully chosen kernels (e.g., the Jackson) suppress Gibbs oscillations and improve convergence of the truncated Chebyshev series. Because the dominant cost is sparse matrix–vector products, KPM scales (nearly) linearly with the size of the Hilbert space, which for non-interacting systems scales like the volume, and can be applied to extremely large sparse Hamiltonians to compute quantities such as the density of states (DOS) and DC conductivity (João & Lopes, 2020).

`KPMru.jl` is a Julia implementation (Bezanson et al., 2017) of KPM for tight-binding models in condensed matter physics, providing a high-performance, user-friendly toolbox. `KPMru.jl` targets large sparse Hamiltonians, integrates with `CUDA.jl` for automatic GPU acceleration when available, and is designed for scalability and ease of integration into Julia-based workflows for large-scale spectral and transport calculations.

Statement of need

`KPMru.jl` provides a high-performance implementation of the Kernel Polynomial Method tailored to tight-binding models. The method avoids ever doing a full diagonalization and instead uses Chebyshev expansions, the package enables spectral and transport calculations on very large sparse Hamiltonians that are infeasible with exact diagonalization (ED). It fills the gap between low-level C/Fortran libraries and interactive, reproducible Julia workflows by offering:

1. **Scalability:** Capable of handling large sparse matrices representing realistic tight-binding models.
2. **Versatility:** Support for DOS and LDOS, DC conductivity (both longitudinal and Hall), and linear and nonlinear optical response functions.
3. **Performance:** Automatic GPU acceleration via `CUDA.jl` when available.

DRAFT

4. **Integration:** As a Julia package, it easily interfaces with other tools in the Julia ecosystem for Hamiltonian generation and data analysis.

In condensed matter physics, understanding the effects of disorder, interactions, and complex lattice geometries often requires numerical simulations of very large Hamiltonians. Traditional ED methods for computing the full spectrum are limited to relatively small system sizes (typically $N \sim 10^4$ states), which can be insufficient to resolve the spectral features of disordered systems or incommensurate structures like twisted bilayer graphene. `KPMru.jl` is especially useful for studies of disorder, moiré systems, and topological materials where large system sizes ($N \sim 10^7$ states) are essential to capture realistic spectral and transport behavior.

State of the field

The Kernel Polynomial Method is well established for computing spectral and response functions of large sparse Hamiltonians (Weisse et al., 2006), and several open-source packages implement it. Kwant provides a general tight-binding transport framework with KPM routines for spectral densities and Kubo conductivity (Groth et al., 2014); pybinding couples Python model construction to C++ KPM routines for densities of states, Green's functions, and conductivity (Moldovan et al., 2017); and KITE targets large-scale real-space calculations of spectral densities, DC and optical conductivities, and nonlinear optical response (João et al., 2020). In Julia, Quantica.jl provides a KPM solver within a broader tight-binding framework (San-Jose, 2024).

`KPMru.jl` instead acts directly on user-supplied sparse Hamiltonians and current operators, separates Chebyshev-moment generation from observable reconstruction so moments can be reused, and runs on GPUs via `CUDA.jl`. We developed a standalone Julia package rather than extending the Python/C++ codes so that model construction, moment generation, and analysis can be modified and combined in a single language, without cross-language interfaces.

Software design

`KPMru.jl` is designed with modularity and performance in mind. The package operates on sparse Hamiltonians and computes Chebyshev moments using stochastic trace estimation with random vectors. The core functionality is divided into three main tiers corresponding to the complexity of the response function:

- **Density of States (1D):** The `KPMru.kpm_1d` function computes the Chebyshev moments for the DOS. It supports stochastic estimation using multiple random vectors (`NR`) and allows users to supply custom input vectors to compute the LDOS. The moments are converted to the spectral density $\rho(E)$ using `KPMru.dos`.
- **Linear Response (2D):** For transport properties like DC conductivity, `KPMru.kpm_2d` calculates the moments required for the Kubo-Greenwood formula involving two current operators (J_x, J_y). The function `KPMru.d_dc_cond` processes these moments to obtain the energy-dependent conductivity $\sigma_{\alpha\beta}(E)$ with $\alpha, \beta = x, y, z$.
- **Nonlinear Response (3D):** The package includes specialized routines (`KPMru.kpm_3d`) for frequency-dependent linear and nonlinear responses, such as the Circular Photogalvanic Effect (CPGE). This involves computing moments for three operators (J_x, J_y, J_z) and post-processing them (`KPMru.d_cpge`) to extract the second-order conductivity $\chi_{xyz}(\omega_1, \omega_2)$.

The software automatically detects available hardware and offloads matrix-vector multiplications to a GPU if a compatible CUDA device is present (`KPMru.whichcore()`), ensuring efficient performance on modern clusters. Further memory can be saved by computing the matrix elements of H and $\mathbf{J}$ on the fly, which is not implemented here.

DRAFT

The Journal of Open Source Software

Example Usage

The following minimal example computes the DOS of a 1-D tight-binding chain using `KPMru.jl`.

```julia
using KPMru
using LinearAlgebra
using SparseArrays
using Plots

# Simple dense 1D tight-binding Hamiltonian (periodic)
function tb1dchain(N::Integer; t::Real=1.0)
    H = zeros(Float64, N, N)
    for i in 1:(N-1)
        H[i, i+1] = -t
        H[i+1, i] = -t
    end
    H[1, N] = -t
    H[N, 1] = -t
    return H
end

# Parameters
N = 500               # system size
NC = 256              # Chebyshev order
NR = 32               # number of random vectors for stochastic trace
nE = 1000             # output energy grid points

H = tb1dchain(N)
# Rescale H -> (-1, 1)
# make the Hamiltonian sparse under complex number
Hsparse = sparse(H.*(1+0*1im))
b, H_norm = KPMru.normalizeH(Hsparse)

# Compute Chebyshev moments (DOS)
mu = KPMru.kpm_1d(H_norm, NC, NR)    # returns moments (array-like)

# Reconstruct DOS on a grid and map energies back to physical scale
E, rho = KPMru.dos(mu, b;kernel = KPMru.JacksonKernel, N_tilde=nE)

# Analytical DOS
rho_exact = zeros(length(E))
mask = abs.(E) .< 2
rho_exact[mask] = 1.0 ./ (π * sqrt.(4 .- E[mask].^2))

# plot the DOS
plot(E, rho, xlabel="E", ylabel="DOS", label="KPM")
plot!(E, rho_exact, lw=1, ls=:dash, label="Analytic")
```

Figure 1 shows the DOS from the same workflow for a larger chain and several expansion orders. Discrepancies between the KPM and analytic results arise from the finite system size, the truncated KPM expansion order, and stochastic trace estimation error.

DRAFT




Figure 1: Density of states for a simple 1D tight-binding chain $L = 10^4$: KPM (solid) with various expansion orders $N_C = 32, 64, 1024$ vs. analytic (dashed). The number of random vectors for the stochastic trace is $N_R = 10$.

Research impact statement

`KPMru.jl` has been utilized in numerous studies to investigate electronic structure and response in complex materials:

1. **Moiré systems and disorder:** Enabled large-scale simulations of twisted bilayer graphene and demonstrated how twist-angle disorder broadens flat bands and modifies spectral features (Chang et al., 2024; Chou et al., 2020; Fu et al., 2020; Wilson et al., 2020; Yi et al., 2022).
2. **Experimentally relevant transport calculations:** Applied to transport modeling based on tight-binding models derived from DFT+U to directly compare theory with experiment (Liu et al., 2021; L. Wu et al., 2020; T.-C. Wu et al., 2025).
3. **Effects of Quasiperiodic potentials:** Used to compute conductivity and disorder-averaged spectral properties and Green's function in disordered materials (Fu et al., 2021; Yi et al., 2022).
4. **Nonlinear optical responses and disordered topological material:** Employed to calculate nonlinear responses such as the CPGE in disordered topological semimetals and to identify spectral signatures of surface and bulk phase transitions in disordered axion insulators (Grindall et al., 2025; A.-K. Wu et al., 2024).
5. **Method benchmarking and numerical validation:** Served as a reference implementation for comparing and validating new Chebyshev-regularization and spectral-density algorithms (Yi et al., 2026).

These applications demonstrate the package's capability to tackle cutting-edge problems in condensed matter theory.

AI usage disclosure

GitHub Copilot was used to assist with updating CUDA-related code, improving tests and documentation, and polishing the manuscript for clarity and readability. The authors reviewed,

DRAFT

edited, and validated all AI-assisted contributions and take full responsibility for the content of this publication.

Acknowledgements

We acknowledge the support of NSF Career Grant No. DMR-1941569, NSF Grant No. DMR-2515945, US Department of Energy Office of Science, Basic Energy Sciences, under Award No. DESC0026023 and the Alfred P. Sloan Foundation through a Sloan Research Fellowship. J.H.W. acknowledges support from the National Science Foundation under Grant Number DMR-2238895.

References

Bezanson, J., Edelman, A., Karpinski, S., & Shah, V. B. (2017). Julia: A fresh approach to numerical computing. *SIAM Review*, *59*(1), 65–98. https://doi.org/10.1137/141000671

Chang, Y., Yi, J., Wu, A.-K., Kugler, F. B., Andrei, E. Y., Vanderbilt, D., Kotliar, G., & Pixley, J. H. (2024). Vacancy-induced tunable Kondo effect in twisted bilayer graphene. *Phys. Rev. Lett.*, *133*, 126503. https://doi.org/10.1103/PhysRevLett.133.126503

Chou, Y.-Z., Fu, Y., Wilson, J. H., König, E. J., & Pixley, J. H. (2020). Magic-angle semimetals with chiral symmetry. *Phys. Rev. B*, *101*, 235121. https://doi.org/10.1103/PhysRevB.101.235121

Fu, Y., König, E. J., Wilson, J. H., Chou, Y.-Z., & Pixley, J. H. (2020). Magic-angle semimetals. *Npj Quantum Materials*, *5*(1), 71. https://doi.org/10.1038/s41535-020-00271-9

Fu, Y., Wilson, J. H., & Pixley, J. H. (2021). Flat topological bands and eigenstate criticality in a quasiperiodic insulator. *Phys. Rev. B*, *104*, L041106. https://doi.org/10.1103/PhysRevB.104.L041106

Grindall, C., Tyner, A. C., Wu, A.-K., Hughes, T. L., & Pixley, J. H. (2025). Separate surface and bulk topological Anderson localization transitions in disordered axion insulators. *Phys. Rev. Lett.*, *135*, 226601. https://doi.org/10.1103/v7x8-ghfy

Groth, C. W., Wimmer, M., Akhmerov, A. R., & Waintal, X. (2014). Kwant: A software package for quantum transport. *New Journal of Physics*, *16*(6), 063065. https://doi.org/10.1088/1367-2630/16/6/063065

João, S. M., Anđelković, M., Covaci, L., Rappoport, T. G., Lopes, J. M. V. P., & Ferreira, A. (2020). KITE: High-performance accurate modelling of electronic structure and response functions of large molecules, disordered crystals and heterostructures. *Royal Society Open Science*, *7*(2), 191809. https://doi.org/10.1098/rsos.191809

João, S. M., & Lopes, J. M. V. P. (2020). Basis-independent spectral methods for non-linear optical response in arbitrary tight-binding models. *J. Phys. Condens. Matter*, *32*(12), 125901. https://doi.org/10.1088/1361-648X/ab59ec

Liu, X., Fang, S., Fu, Y., Ge, W., Kareev, M., Kim, J.-W., Choi, Y., Karapetrova, E., Zhang, Q., Gu, L., Choi, E.-S., Wen, F., Wilson, J. H., Fabbris, G., Ryan, P. J., Freeland, J. W., Haskel, D., Wu, W., Pixley, J. H., & Chakhalian, J. (2021). Magnetic Weyl semimetallic phase in thin films of $\mathrm{Eu_2Ir_2O_7}$. *Phys. Rev. Lett.*, *127*, 277204. https://doi.org/10.1103/PhysRevLett.127.277204

Moldovan, D., Anđelković, M., & Peeters, F. (2017). *Pybinding v0.9.4: A Python package for tight-binding calculations*. Zenodo. https://doi.org/10.5281/zenodo.826942

San-Jose, P. (2024). *Quantica.jl*. Zenodo. https://doi.org/10.5281/zenodo.4762963

DRAFT

Skilling, J. (1988). The axioms of maximum entropy. In *Maximum-entropy and Bayesian methods in science and engineering: foundations* (pp. 173–187). Springer. https://doi.org/10.1007/978-94-009-3049-0_8

Weisse, A., Wellein, G., Alvermann, A., & Fehske, H. (2006). The kernel polynomial method. *Rev. Mod. Phys.*, *78*, 275–306. https://doi.org/10.1103/RevModPhys.78.275

Wilson, J. H., Fu, Y., Das Sarma, S., & Pixley, J. H. (2020). Disorder in twisted bilayer graphene. *Phys. Rev. Res.*, *2*, 023325. https://doi.org/10.1103/PhysRevResearch.2.023325

Wu, A.-K., Guerci, D., Fu, Y., Wilson, J. H., & Pixley, J. H. (2024). Absence of quantization in the circular photogalvanic effect in disordered chiral Weyl semimetals. *Phys. Rev. B*, *110*, 014201. https://doi.org/10.1103/PhysRevB.110.014201

Wu, L., Wen, F., Fu, Y., Wilson, J. H., Liu, X., Zhang, Y., Vasiukov, D. M., Kareev, M. S., Pixley, J. H., & Chakhalian, J. (2020). Berry phase manipulation in ultrathin $\mathrm{SrRuO_3}$ films. *Phys. Rev. B*, *102*, 220406. https://doi.org/10.1103/PhysRevB.102.220406

Wu, T.-C., Chang, Y., Wu, A.-K., Terilli, M., Wen, F., Kareev, M., Choi, E. S., Graf, D., Zhang, Q., Gu, L., & others. (2025). Electronic anisotropy and rotational symmetry breaking at a Weyl semimetal/spin ice interface. *Science Advances*, *11*(24), eadr6202. https://doi.org/10.1126/sciadv.adr6202

Yi, J., König, E. J., & Pixley, J. H. (2022). Low energy excitation spectrum of magic-angle semimetals. *Phys. Rev. B*, *106*, 195123. https://doi.org/10.1103/PhysRevB.106.195123

Yi, J., Massatt, D., Horning, A., Luskin, M., Pixley, J. H., & Kaye, J. (2026). A high-order regularized delta-Chebyshev method for computing spectral densities. *Journal of Computational Physics*, *567*, 115299. https://doi.org/10.1016/j.jcp.2026.115299

DRAFT