The Journal of Open Source Software

PyIRD: A Python-Based Data Reduction Pipeline for Subaru/IRD and REACH

Yui Kasagi[1], **Hajime Kawahara**[1,2], **Ziying Gu**[2], **Teruyuki Hirano**[3,4,5], **Takayuki Kotani**[3,4,5], **Masayuki Kuzuhara**[3,5], and **Kento Masuda**[6]

1 Institute of Space and Astronomical Science, Japan Aerospace Exploration Agency, 3-1-1 Yoshinodai, Chuo-ku, Sagamihara, Kanagawa, 252-5210, Japan **2** Department of Astronomy, Graduate School of Science, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan **3** Astrobiology Center, 2-21-1 Osawa, Mitaka, Tokyo 181-8588, Japan **4** Astronomical Science Program, The Graduate University for Advanced Studies, SOKENDAI, 2-21-1 Osawa, Mitaka, Tokyo 181-8588, Japan **5** National Astronomical Observatory of Japan, 2-21-1 Osawa, Mitaka, Tokyo 181-8588, Japan **6** Department of Earth and Space Science, Osaka University, Osaka 560-0043, Japan

DOI: 10.xxxxxx/draft

Software

- Review
- Repository
- Archive

Editor: Arfon Smith

Submitted: 07 February 2025
Published: unpublished

License
Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License (CC BY 4.0).

DRAFT

Summary

PyIRD is a Python-based pipeline for reducing spectroscopic data obtained with IRD (InfraRed Doppler; Takayuki Kotani et al. (2018)) and REACH (Rigorous Exoplanetary Atmosphere Characterization with High dispersion coronagraphy; T. Kotani et al. (2020)) on the Subaru Telescope. It is designed to process raw images into one-dimensional spectra in a semi-automatic manner. Unlike traditional methods, it does not rely on IRAF (Tody, 1986, 1993), a software used for astronomical data reduction. This approach simplifies the workflow while maintaining efficiency and accuracy. Additionally, the pipeline includes an updated method for removing readout noise patterns from raw images, enabling efficient extraction of spectra even for faint targets such as brown dwarfs.

Statement of need

The reduction of high-dispersion spectroscopic data has traditionally been performed using IRAF, one of the most widely used software tools for astronomical data reduction and analysis. However, the National Optical Astronomy Observatories (NOAO) officially ceased its development and maintenance in 2013. As a result, there is a growing demand for a modern, flexible solution.

PyIRD addresses this need and has already been utilized in several papers (Kawahara et al., 2024; Kawashima et al., 2024; Tomoyoshi et al., 2024).

Key Features

PyIRD is designed to perform data reduction semi-automatically by following a general workflow for high-dispersion spectroscopic data reduction, as illustrated in Figure 1.

The Journal of Open Source Software



Figure 1: Flowchart of the reduction process for IRD and REACH data. The reduction process follows from top to bottom of this figure. Texts in the grey boxes represent instance names of each reduction step used in `PyIRD`.

Users can define a set of FITS-format files using a Python class named `FitsSet`, and functions in the `Stream2D` class are applied to generate the one-dimentional spectrum. Since all functions in `PyIRD` are written in Python rather than IRAF's subset preprocessor language (SPP), the package is easy to develop and maintain. This also significantly reduces the time required for the reduction process: users only need to execute a single Python script without complex IRAF configuration. For example, reducing data with `PyIRD` typically takes a few tens of minutes to produce one-dimensional spectra from raw data obtained during a single observing night, compared to approximately half a day with traditional IRAF methods.

Moreover, `PyIRD` achieves a higher level of readout noise pattern removal on final results. This feature is particularly important for processing data from faint objects such as brown dwarfs, where the astronomical signal is often comparable in strength to systematic noise. The dominant noise source is the readout pattern from the H2RG detector used in IRD. To address this, `PyIRD` models the noise by calculating a median profile for each readout channel and applying a 2D Gaussian Process using `gpkron` (Kawahara, 2022). This innovative method effectively mitigates the readout pattern, as shown in Figure 2, and improves data quality for faint targets.



Figure 2: (Left) Raw image; (Middle) Readout pattern model created by `PyIRD`; (Right) Pattern-corrected image

DRAFT

Acknowledgements

Y.K. acknowledges support from JST SPRING, Grant Number JPMJSP2104 and JSPS KAKENHI grant No. 24K22912. Z.G. acknowledges support from Forefront Physics and Mathematics Program to Drive Transformation (FoPM), a World-leading Innovative Graduate Study (WINGS) Program, the University of Tokyo.

References

Kawahara, H. (2022). Gpkron. In *GitHub repository*. GitHub. https://github.com/HajimeKawahara/gpkron

Kawahara, H., Kawashima, Y., Tada, S., Ishikawa, H. T., Hosokawa, K., Kasagi, Y., Kotani, T., Masuda, K., Nuguroho, S., Tamura, M., Yama, H., Kitzmann, D., Minesi, N., & Morris, B. M. (2024). Differentiable Modeling of Planet and Substellar Atmosphere: High-Resolution Emission, Transmission, and Reflection Spectroscopy with ExoJAX2. *arXiv e-Prints*, arXiv:2410.06900. https://doi.org/10.48550/arXiv.2410.06900

Kawashima, Y., Kawahara, H., Kasagi, Y., Ishikawa, H. T., Masuda, K., Kotani, T., Kudo, T., Hirano, T., Kuzuhara, M., Nugroho, S. K., Livingston, J., Harakawa, H., Nishikawa, J., Omiya, M., Takarada, T., Tamura, M., & Ueda, A. (2024). Atmospheric retrieval of Subaru/IRD high-resolution spectrum of the archetype T-type brown dwarf Gl 229 B. *arXiv e-Prints*, arXiv:2410.11561. https://doi.org/10.48550/arXiv.2410.11561

Kotani, T., Kawahara, H., Ishizuka, M., Jovanovic, N., Vievard, S., Lozi, J., Sahoo, A., Guyon, O., Yoneta, K., & Tamura, M. (2020). Extremely high-contrast, high spectral resolution spectrometer REACH for the Subaru Telescope. In L. Schreiber, D. Schmidt, & E. Vernet (Eds.), *Adaptive optics systems VII* (Vol. 11448, p. 1144878). https://doi.org/10.1117/12.2561755

Kotani, Takayuki, Tamura, M., Nishikawa, J., Ueda, A., Kuzuhara, M., Omiya, M., Hashimoto, J., Ishizuka, M., Hirano, T., Suto, H., Kurokawa, T., Kokubo, T., Mori, T., Tanaka, Y., Kashiwagi, K., Konishi, M., Kudo, T., Sato, B., Jacobson, S., … Oh, D. (2018). The infrared Doppler (IRD) instrument for the Subaru telescope: instrument description and commissioning results. In C. J. Evans, L. Simard, & H. Takami (Eds.), *Ground-based and airborne instrumentation for astronomy VII* (Vol. 10702, p. 1070211). https://doi.org/10.1117/12.2311836

Tody, D. (1993). IRAF in the Nineties. In R. J. Hanisch, R. J. V. Brissenden, & J. Barnes (Eds.), *Astronomical data analysis software and systems II* (Vol. 52, p. 173).

Tody, D. (1986). The IRAF Data Reduction and Analysis System. In D. L. Crawford (Ed.), *Instrumentation in astronomy VI* (Vol. 627, p. 733). https://doi.org/10.1117/12.968154

Tomoyoshi, M., Masuda, K., Hirano, T., Kasagi, Y., Kawahara, H., Kotani, T., Kudo, T., Tamura, M., & Vievard, S. (2024). Weighing Single-lined Spectroscopic Binaries Using Tidal Effects on Radial Velocities: The Case of V723 Monocerotis. *977*(2), 151. https://doi.org/10.3847/1538-4357/ad8cdd

DRAFT