The Journal of Open Source Software

MZsearch: A Python-Based Compound Identification Tool for GC-MS and LC-MS-Based Metabolomics

Hunter Dlugas [1,2*], **Xiang Zhang** [3], **and Seongho Kim** [1,2*]

1 Department of Oncology, School of Medicine, Wayne State University, Detroit, MI, USA **2** Biostatistics and Bioinformatics Core, Karmanos Cancer Institute, Wayne State University, Detroit, MI, USA **3** Department of Chemistry, University of Louisville, Louisville, KY, USA * These authors contributed equally.

DOI: 10.xxxxxx/draft

Software

- Review
- Repository
- Archive

Editor: Julia Romanowska

Reviewers:

- @aboffin
- @lionel42

Submitted: 03 September 2024
Published: unpublished

License
Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License (CC BY 4.0).

DRAFT

Summary

A primary goal in metabolomics, the qualitative and quantitative study of metabolites, is to identify chemical compounds in a given sample. This often involves identifying and quantifying metabolic biomarkers for use in diagnosing, treating, and stratifying the risk of various diseases. A central tool for compound identification in metabolomics is mass spectrometry, which produces the mass spectrum for a given chemical compound. Each peak in the mass spectrum represents an ion fragment, with one axis representing mass to charge ratio (m/z) and the other axis representing relative intensity. A popular method used to identify an unknown chemical compound based on its mass spectrum is spectral library matching. This process involves computing a measure of similarity between the mass spectrum of the unknown chemical compound and each mass spectrum in a reference library. The unknown chemical compound is then identified as the one from the reference library whose mass spectrum is most similar to that of the unknown compound. We present MZsearch, a command-line tool written in Python that implements spectral library matching. The MZsearch tool offers a range of spectrum preprocessing transformations and similarity measures, and it is capable of analyzing data generated from gas chromatography-mass spectrometry (GC-MS) and liquid chromatography-mass spectrometry (LC-MS).

Statement of need

There exist several open-source software tools for working with mass spectrometry data. The Python package matchms allows one to perform spectral library matching for both GC-MS and LC-MS and provides an application programming interface (API) for users to define their own spectrum preprocessing transformations and similarity measures (Huber et al., 2020). The pyOpenMS tool, which is a Python wrapper of the OpenMS tool, is for analyzing LC-MS data and provides common spectrum preprocessing transformations such as filtering based on m/z and intensity values (Röst et al., 2014, 2016). The Python package spectrum_utils provides functionality for preprocessing and visualizing mass spectrometry data (Bittremieux, 2020). The R package metID is a tool for LC-MS-based compound identification with a focus on allowing users to combine their in-house database(s) with public databases (Shen et al., 2022). The R Shiny package ShinyMetID provides users six similarity measures (Cosine, Weighted Cosine, Stein & Scott, Discrete Fourier Transformation, Discrete Wavelet Transformation, and Semi-Partial Correlation) and a graphical user interface (GUI) to perform spectral library matching to identify chemical compounds from GC-MS data (Oh et al., 2023).

In a GC-MS experiment, the resulting mass spectra typically have m/z values as integers. In contrast, LC-MS experiments produce mass spectra with high resolution m/z values, often with many decimal places. This necessitates several spectrum preprocessing steps unique to

LC-MS data analysis, namely centroiding (i.e., merging peaks that are 'close' with respect to their m/z values) and matching (i.e., aligning the m/z values so that the query spectrum and reference spectrum have the same length). In addition to these canonical spectrum preprocessing transformations for LC-MS data, weight factor transformations and low-entropy transformations have been proposed to improve the performance of compound identification for both GC-MS and LC-MS data (Dlugas et al., 2024; Kim et al., 2012; Li et al., 2021). The Shannon Entropy Similarity Measure has been shown to outperform the Cosine Similarity Measure with respect to LC-MS data (Li et al., 2021). A generalization of the Shannon Entropy Similarity Measure, the Tsallis Entropy Similarity Measure, slightly outperforms the Shannon Entropy Similarity Measure in both GC-MS and LC-MS data (Dlugas et al., 2024). This recent study has further demonstrated that the order of preprocessing transformations is also critical to achieving higher accuracy in compound identification. However, to our knowledge, there is no tool that includes the recently-developed, high-performance entropy-based similarity measures or considers the order of the spectrum preprocessing transformations, which are important for effective metabolomics studies.

To address the lack of spectral library matching software that considers both the order of spectrum preprocessing steps and novel entropy-based similarity measures, MZsearch was developed. The developed MZsearch is a command-line tool for performing spectral library matching on either GC-MS or LC-MS data. It allows users to construct their own spectrum preprocessing order using spectrum preprocessing transformations such as weight factor and low-entropy transformations. Additionally, the MZsearch tool includes a novel entropy-based similarity measure, the Rényi Entropy Similarity Measure, enabling users to choose among four similarity measures: the commonly-used Cosine Similarity Measure (Stein & Scott, 1994), the Shannon Entropy Similarity Measure (Li et al., 2021), the recently-developed Entropy Similarity Measure (Dlugas et al., 2024), and the novel Rényi Entropy Similarity Measure.

Functionality

Spectrum Preprocessing Transformations

Functionality implementing the following spectrum preprocessing transformations is offered in MZsearch (more details found in documentation:):

- Filtering based on mz-values and intensities
- Weight Factor Transformation
- Low-Entropy Transformation
- Noise Removal
- Centroiding (only applicable to LC-MS data)
- Matching (only applicable to LC-MS data)
- Normalization (only applicable to entropy-based similarity measures)

The flowchart in Figure 1 depicts the overall workflow of MZsearch.

The Journal of Open Source Software



Figure 1: Workflow of MZsearch.

Similarity Measures

Given a pair of processed spectra intensities $I = (a_1, a_2, ..., a_n), J = (b_1, b_2, ..., b_n) \in \mathbb{R}^n$ with $0 \leq a_i, b_i \leq 1$ for all $i \in \{1, 2, ..., n\}$ and $\sum_{i=1}^{n} a_i = \sum_{i=1}^{n} b_i = 1$, MZsearch provides functionality for computing the following similarity measures:

- Cosine Similarity Measure:

$$S_{Cosine}(I, J) = \frac{I \circ J}{|I|_2 \cdot |J|_2}$$

  where multiplication in the numerator refers to the dot product $I \circ J = a_1 b_1 + a_2 b_2 + ... + a_n b_n$ and multiplication in the denominator refers to multiplication of the L^2-norm of I and J, $|I|_2 = \sqrt{a_1^2 + a_2^2 + ... + a_n^2}, |J|_2 = \sqrt{b_1^2 + b_2^2 + ... + b_n^2}$.

- Shannon Entropy Similarity Measure:

$$S_{Shannon}(I, J) = 1 - \frac{2 \cdot H_{Shannon}\left(\frac{I+J}{2}\right) - H_{Shannon}(I) - H_{Shannon}(J)}{ln(4)},$$

$$H_{Shannon}(I) = -\sum_{i=1}^{n} a_i \cdot ln(a_i)$$

- Tsallis Entropy Similarity Measure (Havrda & Charvát, 1967; Tsallis, 1988):

$$S_{Tsallis}(I, J, q) = 1 - \frac{2 \times H_{Tsallis}(I/2 + J/2, q) - H_{Tsallis}(I, q) - H_{Tsallis}(J, q)}{N_{Tsallis}(I, J, q)},$$

$$N_{Tsallis}(I, J, q) := \frac{\sum_{i=1}^{n} \left(2\left(\frac{a_i}{2}\right)^q + 2\left(\frac{b_i}{2}\right)^q - a_i^q - b_i^q\right)}{1 - q},$$

$$H_{Tsallis}(I, q) = \frac{\left(\sum_{i=1}^{n} a_i^q\right) - 1}{1 - q},$$

$$q \neq 1, \ q > 0$$

- Rényi Entropy Similarity Measure:

$$S_{Renyi}(I, J, q) = 1 - \frac{2 \times H_{Renyi}(I/2 + J/2, q) - H_{Renyi}(I, q) - H_{Renyi}(J, q)}{N_{Renyi}(I, J, q)},$$

$$N_{Renyi}(I, J, q) := \left(\frac{1}{1 - q}\right)\left(2 \times ln\left(\sum_i (a_i/2)^q + \sum_j (b_j/2)^q\right) - ln(\sum_i a_i^q) - ln(\sum_i b_i^q)\right)$$

$$H_{Renyi}(I, q) = \frac{1}{1 - q} ln(\sum_{i=1}^{n} a_i^q),$$

$$q \neq 1, \ q > 0$$

DRAFT

Usage

MZsearch has three main capabilities: (i) converting the raw data to the necessary format for spectral library matching, (ii) running spectral library matching to identify compounds based on their mass spectrometry data, and (iii) plotting a query spectrum vs. a reference spectrum before and after preprocessing transformations.

These tasks are implemented separately for (i) GC-MS and (ii) LC-MS data due to the different spectrum preprocessing transformations. To see all parameters for any of the three main scripts (build_library.py, spec_lib_matching.py, plot_spectra.py), see the documentation at .

Acknowledgements

This research was partially funded by NIH grants (National Institutes of Health R21GM140352), (National Cancer Institute P30CA022453), (National Institutes of Health P20GM113226) and (National Institutes of Health P50AA024337). Computations were performed on Wayne State University's High-Performance Computing Grid.

References

Bittremieux, W. (2020). Spectrum_utils: A Python Package for Mass Spectrometry Data Processing and Visualization. *Analytical Chemistry*, *92*(1), 659–661. https://doi.org/10.1021/acs.analchem.9b04884

Dlugas, H., Zhang, X., & Kim, S. (2024). Comparative analysis of continuous similarity measures for compound identification in mass spectrometry-based metabolomics. *ChemRxiv*. https://doi.org/10.26434/chemrxiv-2024-5fm7t

Havrda, J., & Charvát, F. (1967). Quantification method of classification processes. Concept of structural a-entropy. *Kybernetika*, *3*, 30–35. http://dml.cz/dmlcz/125526

Huber, F., Verhoeven, S., Meijer, C., Spreeuw, H., Castilla, E. M. V., Geng, C., Hooft, J. J. j. van der, Rogers, S., Belloum, A., Diblen, F., & Spaaks, J. H. (2020). Matchms - processing and similarity evaluation of mass spectrometry data. *Journal of Open Source Software*, *5*(52), 2411. https://doi.org/10.21105/joss.02411

Kim, S., Koo, I., Wei, X., & Zhang, X. (2012). A method of finding optimal weight factors for compound identification in gas chromatography-mass spectrometry. *Bioinformatics*, *28*(8), 1158–1163. https://doi.org/10.1093/bioinformatics/bts083

Li, Y., Kind, T., Folz, J., Vaniya, A., Mehta, S. S., & Fiehn, O. (2021). Spectral entropy outperforms MS/MS dot product similarity for small-molecule compound identification. *Nature Methods*, *18*(12), 1524–1531. https://doi.org/10.1038/s41592-021-01331-z

Oh, Y., Kim, S., Kim, S., & Jeong, J. (2023). ShinyMetID: An r shiny package for metabolite identification by mass spectral matching. *Chemometrics and Intelligent Laboratory Systems*, *240*, 104861. https://doi.org/10.1016/j.chemolab.2023.104861

Röst, H. L., Sachsenberg, T., Aiche, S., Bielow, C., Weisser, H., Aicheler, F., Andreotti, S., Ehrlich, H.-C., Gutenbrunner, P., Kenar, E., Liang, X., Nahnsen, S., Nilse, L., Pfeuffer, J., Rosenberger, G., Rurik, M., Schmitt, U., Veit, J., Walzer, M., … Kohlbacher, O. (2016). OpenMS: A flexible open-source software platform for mass spectrometry data analysis. *Nature Methods*, *13*(9), 741–748. https://doi.org/10.1038/nmeth.3959

Röst, H. L., Schmitt, U., Aebersold, R., & Malmström, L. (2014). pyOpenMS: A Python-based interface to the OpenMS mass-spectrometry algorithm library. *PROTEOMICS*, *14*(1), 74–77. https://doi.org/10.1002/pmic.201300246

DRAFT

Shen, X., Wu, S., Liang, L., Chen, S., Contrepois, K., Zhu, Z.-J., & Snyder, M. (2022). metID: An r package for automatable compound annotation for LC-MS-based data. *Bioinformatics*, *38*(2), 568–569. https://doi.org/10.1093/bioinformatics/btab583

Stein, S. E., & Scott, D. R. (1994). Optimization and testing of mass spectral library search algorithms for compound identification. *Journal of the American Society for Mass Spectrometry*, *5*(9), 859–866. https://doi.org/10.1016/1044-0305(94)87009-8

Tsallis, C. (1988). Possible generalization of boltzmann-gibbs statistics. *Journal of Statistical Physics*, *52*, 479–487. https://doi.org/10.1007/bf01016429

DRAFT