The Journal of Open Source Software

OrbDot: A Python package for studying the secular evolution of exoplanet orbits

Simone R. Hagey [1] **and Aaron Boley** [1]

1 Department of Physics and Astronomy, The University of British Columbia, 6224 Agricultural Road Vancouver, BC, V6T 1Z1, Canada

DOI: 10.xxxxxx/draft

Software

- Review
- Repository
- Archive

Editor:

Submitted: 06 May 2025
Published: unpublished

License
Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License (CC BY 4.0).

In partnership with

AMERICAN ASTRONOMICAL SOCIETY

This article and software are linked with research article DOI 10.3847/xxxxx, published in the Astronomical Journal.

DRAFT

Summary

Gradual changes in exoplanet orbits, known as secular variations, can be detected through observations of transits, eclipses, and radial velocities that span multiple decades in time. Their detection and characterization enable the study of a wide range of dynamical phenomena, such as orbital decay and precession, which operate on timescales of millions of years. Under certain conditions, measurements of secular variations can even probe the interior structure of exoplanets, providing a unique tool for understanding exoplanet formation and evolution.

The necessity to search over many orbital epochs coupled with an ever-growing archive of exoplanet observations creates a need for fast and flexible open-source software that can reliably detect gradual changes in exoplanet orbits. `OrbDot` addresses this need by offering robust tools for fitting secular evolution models to exoplanet transit and eclipse mid-times, transit durations, and radial velocity data.

A key advantage of `OrbDot` is its ability to fit multiple data types simultaneously, which can help to break parameter degeneracies. It also excels at assisting in result interpretation by generating reports on model comparisons and assessments of various physical effects in the context of the models and their corresponding theory. For example, analysis reports could determine key parameters for assessing tidal energy dissipation, apsidal precession mechanisms, variations due to systemic proper motion, and the dynamical effects of non-resonant companion objects, depending on the applied models.

`OrbDot` remains highly efficient with multiple data types and a high number of free parameters, as it utilizes the powerful nested sampling algorithms of the `Nestle` (Barbary, 2021; Skilling, 2006) and `PyMultiNest` (Buchner et al., 2014; Feroz et al., 2009) packages. The intricacies of the implementation are abstracted such that the `OrbDot` input files are simple and the method calls require only a list of free parameters, along with the desired model for fitting.

Extensive documentation, including examples, is hosted on ReadTheDocs.

The examples demonstrate that `OrbDot` can quickly reproduce literature results using only a few lines of code. Readers may be especially interested in the `OrbDot` example analysis of the transit and eclipse mid-times of Hot Jupiter WASP-12 b, which is well-known for showing strong evidence for orbital decay.

A complementary case study of TrES-1 b (CITE) illustrates the full capabilities of `OrbDot`, placing it in a broader scientific context. Moreover, an early version of this code was used for the orbital analysis of the Hot Neptune LTT-9779 b, published in Edwards et al. (2023).

Statement of need

Many exoplanet systems now have transit and radial velocity data spanning over a decade, enabling studies of secular variations. While tools for analyzing short-term transit variations exist, there is a lack of open-source software dedicated to long-term orbital evolution. This does not, however, reflect a lack of interest, as the number of such studies is growing rapidly.

`OrbDot` lowers the barrier to entry for researchers at all levels, including undergraduates, by making advanced statistical methods accessible without requiring extensive computational experience. Despite its ease of use, `OrbDot` is not intended to be a black box. Rather, with extensive documentation, examples, and accessible source code, it is presented to the community with transparency that lends itself to community contributions and independent verification of results. It is designed to be easily extendable, as the nested sampling framework supports custom log-likelihood models with free parameters that are part of the `OrbDot` ecosystem. This ensures that the software may evolve to meet the needs of the research community.

Acknowledgements

This work was supported, in part, by an NSERC Discovery Grant (DG-2020-04635) and the University of British Columbia. SH's contribution was further supported, in part, by an NSERC PGS-D and a Li Tze Fong Fellowship.

References

Barbary, K. (2021). *nestle: Nested sampling algorithms for evaluating Bayesian evidence*.

Buchner, J., Georgakakis, A., Nandra, K., Hsu, L., Rangel, C., Brightman, M., Merloni, A., Salvato, M., Donley, J., & Kocevski, D. (2014). X-ray spectral modelling of the AGN obscuring region in the CDFS: Bayesian model selection and catalogue. *Astronomy & Astrophysics*, *564*, A125. https://doi.org/10.1051/0004-6361/201322971

Edwards, B., Changeat, Q., Tsiaras, A., Allan, A., Behr, P., Hagey, S. R., Himes, M. D., Ma, S., Stassun, K. G., Thomas, L., Thompson, A., Boley, A., Booth, L., Bouwman, J., France, K., Lowson, N., Meech, A., Phillips, C. L., Vidotto, A. A., … Ward-Thompson, D. (2023). Characterizing a World Within the Hot-Neptune Desert: Transit Observations of LTT 9779 b with the Hubble Space Telescope/WFC3. *The Astronomical Journal*, *166*(4), 158. https://doi.org/10.3847/1538-3881/acea77

Feroz, F., Hobson, M. P., & Bridges, M. (2009). MULTINEST: an efficient and robust Bayesian inference tool for cosmology and particle physics. *Monthly Notices of the Royal Astronomical Society*, *398*(4), 1601–1614. https://doi.org/10.1111/j.1365-2966.2009.14548.x

Skilling, J. (2006). Nested sampling for general Bayesian computation. *Bayesian Analysis*, *1*(4), 833–859. https://doi.org/10.1214/06-BA127

DRAFT