The Journal of Open Source Software

FRAGMENT-MNP: A model of micro- and nanoplastic fragmentation in the environment

Sam Harrison [1], **Cansu Uluseker** [1], **Patrizia Pfohl** [2], **Katherine Santizo**[2,3], **Joana Sipe** [4,7], **Brandon Lopez**[4], **Antonia Praetorius** [5], **Richard K. Cross** [6], **Gbotemi A. Adediran** [6,8], **Claus Svendsen** [6], **Wendel Wohlleben** [2], **and Mark Wiesner** [4]

1 UK Centre for Ecology & Hydrology, Library Avenue, Bailrigg, Lancaster, LA1 4AP, UK **2** BASF SE, Carl-Bosch-Str. 38, 67056 Ludwigshafen, Germany **3** Cefic aisbl, Rue Belliard 40 (box 15) – 1040, Brussels, Belgium **4** Duke University, Durham, NC 27708, United States **5** University of Amsterdam, 1090 GE Amsterdam, Netherlands **6** UK Centre for Ecology & Hydrology, Benson Lane, Crowmarsh Gifford, Wallingford, OX10 8BB, UK **7** Ira A. Fulton Schools of Engineering, Arizona State University, 1151 S Forest Ave, Tempe, AZ, United States **8** School of Earth and Environment, University of Leeds, Leeds, LS2 9JT, UK

DOI: 10.xxxxxx/draft

Software

- Review
- Repository
- Archive

Editor:

Submitted: 18 February 2025
Published: unpublished

License
Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License (CC BY 4.0).

DRAFT

Summary

The degradation and fragmentation of plastics in the environment is an important but relatively poorly characterised process. Fragmentation leads to the formation of micro- and nanoplastics, and research has shown that particle size and shape, and thereby fragmentation, impacts a variety of processes, such as the ability of organisms to uptake plastics, and the movement of plastics around the environment (Thompson et al., 2024; Thornton Hampton et al., 2022). In this paper, we present the FRAGMENT-MNP model as an open-source mechanistic model of micro- and nanoplastic degradation and fragmentation. FRAGMENT-MNP predicts the time evolution of particle size distributions, providing invaluable insights into fragmentation dynamics to help develop our understanding of plastic degradation and fragmentation, and offering predictive capabilities to enable better prediction of the fate and exposure of plastics in the environment.

Statement of need

By modifying particle sizes and shapes, fragmentation influences the potential risk caused by plastics in the environment. Therefore, to assess the risk caused by plastic pollution, understanding fragmentation is crucial. Predictive models are essential in helping this assessment, enabling us to fill gaps in observation data, better understand the results of experiments, and predict theoretical scenarios, such as in prospective risk assessments. Despite this, existing models that predict plastic transport, fate and exposure to organisms either do not consider fragmentation, include it only as a loss process (Quik et al., 2023), treat fragmentation as independent of the properties and residence time in the environment (Koelmans et al., 2017), or consider fragmentation as cascading (fragmenting mass can only be partitioned to the next biggest size class, rather than allowing the formation of nano-scale fragments) (Domercq et al., 2022; Kaandorp et al., 2021). In reality, fragmentation depends on the environmental stresses encountered in the environment, such as photolysis by sunlight, hydrolysis by water, enzymatic action and mechanical disruption causing the break-apart of particles (e.g. the action of waves or bioturbation by soil invertebrates). Fragmentation has also been shown to often occur via surface erosion into nanoscale fragments (Meides et al., 2021; Menzel et al., 2022), rather than in a cascading manner. There is a clear need for flexible and accessible model that can

account for these factors, and FRAGMENT-MNP fills this gap.

Overview of the model

The model is fully documented at https://fragmentmnp.ceh.ac.uk. Here, we provide a brief overview of its main conceptualisation and functionality.

Particle concentrations c_k are represented in binned size classes k, with the model allowing for the fragmentation of particles from larger to smaller size classes, and the dissolution of particles into a dissolved size class with concentration c_{diss}. Conceptually, the dissolved fraction consists of oligomers, monomers and volatile organic compounds. The solutions are obtained by numerically solving the following set of differential equations:

$$\frac{dc_k}{dt} = -k_{\text{frag},k}c_k + \sum_i f_{i,k}k_{\text{frag},i}c_i - k_{\text{diss},k}c_k$$

$$\frac{dc_{\text{diss}}}{dt} = \sum_k k_{\text{diss},k}c_k$$

Here, $k_{\text{frag},k}$ is the fragmentation rate constant of size class k, $f_{i,k}$ is the fraction of daughter fragments produced from a fragmenting particle of size i that are of size k, and $k_{\text{diss},k}$ is the dissolution rate from size class k. The rate constants k_{frag} and k_{diss} can be a function of time and particle surface area, and thus are represented internally as 2D arrays. The shape of these dependencies can be controlled by model input parameters, which allow for constant, linear, polynomial, power law, exponential, logarithmic or logistic dependencies. This flexible parameterisation allows for rate constants to model a variety of physical phenomena, such as the modulation of fragmentation as polymer particles undergo weathering in the environment.

FRAGMENT-MNP is released as a `pip` package (`pip install fragmentmnp`) and comes with example configuration and data,to enable users to get started using the model in as little time as possible. A bare minimum model run is shown below, which uses these examples to run an arbitrary fragmentation scenario (with no dissolution):

```
from fragmentmnp import FragmentMNP
from fragmentmnp.examples import minimal_config, minimal_data

# Create the model and pass it the example config and data
fmnp = FragmentMNP(minimal_config, minimal_data)
# Run the model
output = fmnp.run()
# Plot the results
output.plot()
```

DRAFT



Figure 1: Example model output showing the time evolution of particle size distributions undergoing fragmentation

Related work

The development of FRAGMENT-MNP is part of broader efforts to further our understanding of microplastics in the environment. This includes the development of other models covering emissions, additive release, long-range transport and exposure. Ongoing work is pursuing closer integration of these models and associated experimental data in order to provide holistic predictions of microplastic fate and exposure in the environment.

Acknowledgements

Thank you to the European Chemical Industry Council Long-Range Research Initiative (Cefic-LRI) for providing funding for this work, under project number ECO59.

References

Domercq, P., Praetorius, A., & MacLeod, M. (2022). The Full Multi: An open-source framework for modelling the transport and fate of nano- and microplastics in aquatic systems. *Environmental Modelling & Software*, *148*, 105291. https://doi.org/10.1016/j.envsoft.2021.105291

Kaandorp, M. L. A., Dijkstra, H. A., & Sebille, E. van. (2021). Modelling size distributions of marine plastics under the influence of continuous cascading fragmentation. *Environmental Research Letters*, *16*(5), 054075. https://doi.org/10.1088/1748-9326/abe9ea

Koelmans, A. A., Kooi, M., Law, K. L., & Sebille, E. van. (2017). All is not lost: Deriving a top-down mass budget of plastic at sea. *Environmental Research Letters*, *12*(11), 114028. https://doi.org/10.1088/1748-9326/aa9500

Meides, N., Menzel, T., Poetzschner, B., Löder, M. G. J., Mansfeld, U., Strohriegl, P., Altstaedt, V., & Senker, J. (2021). Reconstructing the Environmental Degradation of Polystyrene by Accelerated Weathering. *Environmental Science & Technology*, *55*(12), 7930–7938. https://doi.org/10.1021/acs.est.0c07718

DRAFT

Menzel, T., Meides, N., Mauel, A., Mansfeld, U., Kretschmer, W., Kuhn, M., Herzig, E. M., Altstädt, V., Strohriegl, P., Senker, J., & Ruckdäschel, H. (2022). Degradation of low-density polyethylene to nanoplastic particles by accelerated weathering. *Science of The Total Environment*, *826*, 154035. https://doi.org/10.1016/j.scitotenv.2022.154035

Quik, J. T. K., Meesters, J. A. J., & Koelmans, A. A. (2023). A multimedia model to estimate the environmental fate of microplastic particles. *Science of The Total Environment*, *882*, 163437. https://doi.org/10.1016/j.scitotenv.2023.163437

Thompson, R. C., Courtene-Jones, W., Boucher, J., Pahl, S., Raubenheimer, K., & Koelmans, A. A. (2024). Twenty years of microplastic pollution research—what have we learned? *Science*, *386*(6720), eadl2746. https://doi.org/10.1126/science.adl2746

Thornton Hampton, L. M., Brander, S. M., Coffin, S., Cole, M., Hermabessiere, L., Koelmans, A. A., & Rochman, C. M. (2022). Characterizing microplastic hazards: Which concentration metrics and particle characteristics are most informative for understanding toxicity in aquatic organisms? *Microplastics and Nanoplastics*, *2*(1), 20. https://doi.org/10.1186/s43591-022-00040-4

DRAFT