The Journal of Open Source Software

ParticleTracking: A GUI and library for particle tracking on stereo camera images

Adrian Niemann [1], **Dmitry Puzyrev** [1,3,4], **and Ralf Stannarius** [2,3,4,5]

1 Department of Nonlinear Phenomena, Institute of Physics, Otto von Guericke University Magdeburg, Germany **2** Institute of Physics, Otto von Guericke University Magdeburg, Germany **3** Department of Microgravity and Translational Regenerative Medicine, Medical Faculty, Otto von Guericke University Magdeburg, Germany **4** Research Group 'Magdeburger Arbeitsgemeinschaft für Forschungunter Raumfahrt-und Schwerelosigkeitsbedingungen' (MARS), Otto von Guericke University Magdeburg, Germany **5** Department of Engineering, Brandenburg University of Applied Sciences, Germany

DOI: 10.xxxxxx/draft

Software

- Review
- Repository
- Archive

Editor: Jeff Gostick

Reviewers:

- @mhubii
- @aquilesC
- @merrygoat
- @jgostick

Submitted: 15 August 2023
Published: unpublished

License
Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License (CC BY 4.0).

DRAFT

Summary

The `ParticleTracking` software is intended to facilitate semi-automatic detection, 3D position and orientation reconstruction and tracking of arbitrarily shaped particles from 2-view stereo camera footage. The software consists of two packages, `RodTracker` and `ParticleDetection`. The `ParticleDetection` package provides functionality for training and application of neural networks (e.g. Mask R-CNN) for particle detection in camera images, as well as automatic 3D matching and multi-object tracking of these particles. The `RodTracker` package is a graphical user interface (GUI) for the particle tracking task, encapsulating the functionality of `ParticleDetection` and providing means to manually correct the automatically generated particle coordinates and tracking data.

The main features of this software are given below with a more extensive feature description available in the documentation under https://particletracking.readthedocs.io/en/latest/:

- training and application of (Detectron2) Mask R-CNN models for detecting particles on images
- automated particle endpoint localization from segmentation masks
- automated assignment of particle correspondences (3D matching) between two camera views
- reconstruction of 3D coordinates and orientations of particles identified on camera images
- automated tracking of particles over multiple stereo camera frames, i.e. the course of an experiment
- providing a GUI for applying manual corrections to the automatically generated data with a typical workflow shown in Figure 1

The main focus of this software is currently on elongated (rod-shaped) particles, but it is extensible with new particle geometries. The software can also be modified for inclusion of additional camera views for more accurate 3D tracking, or for 1-view 2D particle tracking. The `RodTracker` software is currently employed for data extraction in the German Aerospace Center (DLR) projects EVA II (50WK2348), VICKI (50WM2252), and CORDYGA (50WM2242). Recently, it was used in research article on cooling of granular gas mixture in microgravity (Puzyrev et al., 2024). Other publications that use this library for data extraction are currently in preparation. The prototype software for particle detection and tracking was described in (Puzyrev et al., 2020).



Figure 1: Typical workflow with the `RodTracker` for data extraction.

Statement of need

Many natural and industrial processes deal with granular gases, i.e. dilute ensembles of macroscopic particles floating and colliding in space. One of the defining features of such systems is inelasticity of the collision, i.e. dissipation of particle kinetic energy. This leads to fascinating phenomena such as spontaneous clustering, absence of energy equipartition and non-Gaussian velocity distributions. While most of 2D experiments can be performed in normal gravity, 3D experiments with granular gases require microgravity conditions. Starting from the pioneering results on cluster formation (É. Falcon et al., 1999), 3D experiments have been reported for spherical grains (E. Falcon et al., 2006; Yu et al., 2020), ellipsoids (Pitikaris et al., 2022) and rods (K. Harth et al., 2013; Kirsten Harth et al., 2018).

In typical microgravity experiments, ensembles of particles are placed in the container, excited mechanically or magnetically and observed with a stereo-camera setup. Many experiments were performed in the VIP-Gran instrument by the Space Grains ESA Topical team (spacegrains.org) during parabolic flight campaigns. In the majority of VIP-Gran experiments, particle density does not allow for tracking individual grains.

Another possibility is to perform the experiment with dilute ensembles, where most particles can be directly observed on video footage (Kirsten Harth et al., 2018; Puzyrev et al., 2020). In this case, the focus has been on experiments with elongated particles, due to the fact that collision rates for such particles are much higher than for spheres for the same packing fraction. Thus, even if particles overlap on the camera views, usually their endpoints still can be observed and their 3D positions and orientations can be reconstructed. In addition, study of elongated particles allows to observe the evolution of their orientations and to find the kinetic energy associated with the rotational degrees of freedom. Experiments with other particle types are planned as well.

For the study of such systems, it is beneficial to know the 3D positions and orientations over time for as many particles as possible. To achieve statistically meaningful results, the tracking

DRAFT

of many tens to hundreds of particles is usually required. With that information, a reliable statistical analysis of the ensemble properties and their evolution over time can be achieved. Due to the large number of simultaneously tracked objects and their relatively high velocity, accurate experimental data analysis requires high frame rates. In one drop tower experimental run, around 9 seconds of 100 to 240 fps video footage must be analyzed. This makes manual data analysis exceptionally time-consuming. Due to the large number of overlapping particles, conventional particle detection methods based on color separation, morphological operations, and Hough transform have proven to be unstable. For this reason, an AI-assisted approach based on Matterport Mask R-CNN implementation (Abdulla, 2017; He et al., 2017) has been successfully employed (Puzyrev et al., 2020) in extraction and processing of data from the raw stereo camera images. This approach still suffered from long manual data processing times, due to the necessity to correct remaining errors after automatic particle detection, matching and tracking, as well as a suboptimal user interface to perform the correction tasks.

The `ParticleTracking` software is an evolution of the AI-assisted framework for the analysis of dilute granular ensembles, improved by the transition to the Detectron2 platform, inclusion of a GUI, and a documented and extensible codebase.

Dependencies

Among others, the software depends on the following open source libraries: For the particle detection the Detectron2 (Wu et al., 2019) framework is used. For tracking the software relies heavily on functions provided by numpy (Harris et al., 2020), scipy (Virtanen et al., 2020) and PuLP. The GUI was constructed with PyQt5 and is using pandas (team, 2022) for its data management.

Acknowledgements

We want to acknowledge the valuable feedback and bug reports given by the users of our software, specifically Mahdieh Mohammadi and Kirsten Harth. We also want to acknowledge the work of Meera Subramanian and Adithya Viswanathan that provided a first prototype of the `RodTracker` GUI.

The development of this software has been financially supported by the German Aerospace Center (DLR) within grants 50WM2048/50WK2348 (Projects EVA/EVA-II) and 50WM2252 (Project VICKI).

References

Abdulla, W. (2017). Mask R-CNN for object detection and instance segmentation on keras and TensorFlow. In *GitHub repository*. https://github.com/matterport/Mask_RCNN; Github.

Falcon, E., Aumaitre, S., Evesque, P., Palencia, F., Lecoutre-Chabot, C., Fauve, S., Beysens, D., & Garrabos, Y. (2006). Collision statistics in a dilute granular gas fluidized by vibrations in low gravity. *Eur. Phys. Lett.*, *74*, 830–836. https://doi.org/10.1209/epl/i2005-10589-8

Falcon, É., Wunenburger, R., Évesque, P., Fauve, S., Chabot, C., Garrabos, Y., & Beysens, D. (1999). Cluster formation in a granular medium fluidized by vibrations in low gravity. *Phys. Rev. Lett.*, *83*, 440–443. https://doi.org/10.1103/physrevlett.83.440

Harris, C. R., Millman, K. J., Walt, S. J. van der, Gommers, R., Virtanen, P., Cournapeau, D., Wieser, E., Taylor, J., Berg, S., Smith, N. J., Kern, R., Picus, M., Hoyer, S., Kerkwijk, M. H. van, Brett, M., Haldane, A., Río, J. F. del, Wiebe, M., Peterson, P., … Oliphant, T. E. (2020). Array programming with NumPy. *Nature*, *585*(7825), 357–362. https://doi.org/10.1038/s41586-020-2649-2

Harth, K., Kornek, U., Trittel, T., Strachauer, U., Höme, S., Will, K., & Stannarius, R. (2013). Granular gases of rod-shaped grains in microgravity. *Phys. Rev. Lett.*, *110*, 144102. https://doi.org/10.1103/PhysRevLett.110.144102

Harth, Kirsten, Trittel, T., Wegner, S., & Stannarius, R. (2018). Free cooling of a granular gas of rodlike particles in microgravity. *Phys. Rev. Lett.*, *120*, 214301. https://doi.org/10.1103/PhysRevLett.120.214301

He, K., Gkioxari, G., Dollár, P., & Girshick, R. B. (2017). Mask R-CNN. *CoRR*, *abs/1703.06870*. https://arxiv.org/abs/1703.06870

Pitikaris, S., Bartz, P., Yu, P., Cristoforetti, S., & Sperl, M. (2022). Granular cooling of ellipsoidal particles in microgravity. *Npj Microgravity*, *8*(1), 11. https://doi.org/10.1038/s41526-022-00196-6

Puzyrev, D., Harth, K., Trittel, T., & Stannarius, R. (2020). Machine learning for 3D particle tracking in granular gases. *Microgravity Science and Technology*, *32*(5), 897–906. https://doi.org/10.1007/s12217-020-09800-4

Puzyrev, D., Trittel, T., Harth, K., & Stannarius, R. (2024). Cooling of a granular gas mixture in microgravity. *Npj Microgravity*, *10*(1), 36. https://doi.org/10.1038/s41526-024-00369-5

team, T. pandas development. (2022). *Pandas-dev/pandas: pandas* (Version v1.5.2). Zenodo. https://doi.org/10.5281/zenodo.7344967

Virtanen, P., Gommers, R., Oliphant, T. E., Haberland, M., Reddy, T., Cournapeau, D., Burovski, E., Peterson, P., Weckesser, W., Bright, J., van der Walt, S. J., Brett, M., Wilson, J., Millman, K. J., Mayorov, N., Nelson, A. R. J., Jones, E., Kern, R., Larson, E., … SciPy 1.0 Contributors. (2020). SciPy 1.0: Fundamental Algorithms for Scientific Computing in Python. *Nature Methods*, *17*, 261–272. https://doi.org/10.1038/s41592-019-0686-2

Wu, Y., Kirillov, A., Massa, F., Lo, W.-Y., & Girshick, R. (2019). *Detectron2*. https://github.com/facebookresearch/detectron2.

Yu, P., Schröter, M., & Sperl, M. (2020). Velocity distribution of a homogeneously cooling granular gas. *Phys. Rev. Lett.*, *124*, 208007. https://doi.org/10.1103/PhysRevLett.124.208007

DRAFT