

strongcoca: A Python package for strong coupling in nanoscale systems

Jakub Fojt^{1*}, Tuomas P. Rossi^{2,3*}, Viktor Martvall¹, Tomasz J. Antosiewicz⁴, Mikael Kuisma^{5,6¶}, and Paul Erhart^{1*¶}

¹ Department of Physics and Astronomy, Chalmers University of Technology, 412 96 Gothenburg, Sweden ² Department of Applied Physics, Aalto University, 00076 Aalto, Finland ³ CSC – IT Center for Science Ltd., P.O. Box 405, 02101 Espoo, Finland ⁴ Faculty of Physics, University of Warsaw, Pasteura 5, PL-02-093 Warsaw, Poland ⁵ Department of Chemistry, University of Jyväskylä, 40014 Jyväskylä, Finland ⁶ CAMD, Department of Physics, Technical University of Denmark, 2800 Kongens Lyngby, Denmark ¶ Corresponding author * These authors contributed equally.

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#)
- [Repository](#)
- [Archive](#)

Editor: [Evan Spotte-Smith](#)

Reviewers:

- [@victorchanglee](#)

Submitted: 25 May 2026

Published: unpublished

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

Summary

Electromagnetic (EM) simulations are ubiquitous for the modeling of spectroscopy and near-field enhancement in plasmonics, connecting microscopic phenomena to experimentally observed quantities. Here, we present strongcoca, a Python package that calculates the EM response using a subsystem approach. In this approach, the response of a system is calculated from the interactions between subsystems and the individual response of each subsystem, e.g., a molecule or a nanostructure. The user supplies tabulated response functions for the subsystems as well as their positions and orientations. In the current implementation of strongcoca, the EM interactions are approximated up to the order of dipole–dipole interactions, as discussed further below.

Response functions for subsystems can be taken from various sources. For example, strongcoca provides parsers for the output of time-dependent density-functional theory (TDDFT) calculations carried out using the GPAW ([Mortensen et al., 2024](#)) and NWChem ([Aprà et al., 2020](#)) codes. It is also possible to use analytical Mie–Gans models with tabulated dielectric functions, which allows one to use experimental data. Additionally, strongcoca interfaces to the Atomic Simulation Environment (ASE) ([Larsen et al., 2017](#)) for visualization of the coupled systems and relies on NumPy ([Harris et al., 2020](#)) and SciPy ([Virtanen et al., 2020](#)) for linear algebra and vectorized operations.

The full documentation for strongcoca in addition to examples and tutorials can be found at <https://strongcoca.materialsmodeling.org/>.

Statement of need

In the common approach to EM simulations, matter is treated as a continuum, with material-specific response functions (e.g., dielectric functions) obtained from experiment or simulations of bulk materials. This approach fails to accurately capture the response of nanoscale materials with irregularities on the atomic scale, or of individual organic molecules, due to the inherent quantum nature of electrons on the atomic scale. On the other hand, fully quantum mechanical calculations, while accurate, exhibit steep polynomial scaling with system size and remain intractable even for moderately sized systems. In the approach to EM simulations taken in strongcoca, we instead rely on explicitly separating the system of interest into subsystems for which the response is known, thereby circumventing these issues. The response function (polarizability) of each subsystem can capture the correct quantum behavior if it is obtained

at the appropriate level of theory or taken from experiment. `strongcoca` is therefore aimed at researchers in nanophotonics and quantum optics who need to model interactions between distinct quantum and nanoscale subsystems.

In the current implementation, `strongcoca` neglects (1) higher-order multipolar interactions between subsystems, (2) charge transfer between subsystems, and (3) quantum mechanical hybridization between the subsystems. These approximations are less significant for well-separated subsystems, but lead to inaccurate results at small separations. It is possible to extend `strongcoca` to properly take into account higher-order multipoles (1) and charge transfer (2), but the lack of quantum mechanical hybridization (3) is fundamentally a limitation of our subsystem approach. The dipole-dipole interactions currently implemented in `strongcoca` are sufficient to study, e.g., strong light-matter coupling (Ebbesen, 2016) between nanoscale materials and organic molecules. `strongcoca` is thus an excellent tool for **strong coupling** calculations, from which the package takes its name.

State of the field

Two EM codes whose implementation most closely resembles that of `strongcoca` are PyGDM2 (Wiecha, 2018; Wiecha et al., 2022), implementing the Green dyadic method, and DDSCAT (Draine & Flatau, 1994), implementing the discrete-dipole approximation. Both of these codes calculate the response of a system made up of many interacting dipoles that together represent a single continuum material. In `strongcoca`, by contrast, each dipole represents an explicit subsystem, such as an organic molecule or a nanostructure, with its own individually known response function. `strongcoca` therefore has an inherently different scope than these packages: rather than modeling the EM response of a single material, it is designed to study interactions between distinct subsystems with individually known response functions. However, in principle a single larger polarizable nanoparticle could also be described in `strongcoca` as a collection of closely packed dipolar elements, just as in other discrete-dipole approximation implementations, thereby enabling the description of higher-order multipolar interactions between this subsystem and others.

The code PyCharge (Filipovich & Hughes, 2022) can also calculate dipole-dipole interactions, although it supports only single-Lorentzian response functions, i.e., subsystems with a single resonance. While it would be possible to describe a general material using a sum of Lorentzians, there is no user interface to conveniently do so. In addition, PyCharge performs simulations in the time domain (allowing for explicitly time-dependent positions), while `strongcoca` works in the frequency domain, which is orders of magnitude faster for the applications targeted here.

Software design

`strongcoca` follows an object-oriented structure, where physical subsystems are defined independently and then assembled into an interacting system. The response of each subsystem is encapsulated in a `PolarizableUnit` object, which carries a response function, a position, and an orientation in space. The response function can be provided through any of the available parsers, for example from a TDDFT calculation in GPAW or NWChem, or from an analytical Mie-Gans model with a tabulated dielectric function. This makes it straightforward to mix subsystems described at different levels of theory or obtained from different software packages. Once the individual subsystems are defined, they can be combined into a `CoupledSystem` object, which manages the dipole-dipole interaction tensor between all pairs of subsystems. Physical observables of the interacting system, such as absorption spectra and near-field enhancements, are then computed by passing the `CoupledSystem` to the appropriate calculator object.

The object-oriented design also makes it straightforward to extend `strongcoca` with new parsers, new response function types, and new calculators, each of which can be added independently without modifying the core framework.

Research impact statement

strongcoca has been under active development for approximately five years, during which it has been used internally by the developers to study strong light–matter coupling in plasmonic systems. In (Fojt et al., 2021) it was used to study strong coupling in assemblies of molecules and nanoparticles, and in (Kuisma et al., 2022) to study strong coupling between molecules and plasmonic nanocavities. Following requests from other researchers, we have decided to release strongcoca as an open-source package.

AI usage disclosure

No AI tools were used in developing the software included in this release. In ongoing development work, Claude Code has been used for code assistance, and all AI-assisted code is carefully reviewed by the authors.

Acknowledgements

This work has received funding from the Knut and Alice Wallenberg Foundation (Grant No. 2019.0140, J.F., P.E.), the Swedish Research Council (Grant No. 2015-04153, J.F., P.E.), the Research Council of Finland (Grant Nos. 332429, T.P.R; 295602, M.K.), the Polish National Science Center (Grant No. 2019/34/E/ST3/00359, T.J.A.), and been enabled by computational resources provided by the National Academic Infrastructure for Supercomputing in Sweden (NAISS) and the Swedish National Infrastructure for Computing (SNIC) at C3SE, NSC, and PDC partially funded by the Swedish Research Council (Grant Nos. 2018-05973 and 2022-06725), as well as by the CSC – IT Center for Science, Finland, by the Aalto Science-IT project, Aalto University School of Science, and by the Interdisciplinary Center for Mathematical and Computational Modeling, University of Warsaw (Grant No. G55-6).

References

- Aprà, E., Bylaska, E. J., de Jong, W. A., Govind, N., Kowalski, K., Straatsma, T. P., Valiev, M., van Dam, H. J. J., Alexeev, Y., Anchell, J., Anisimov, V., Aquino, F. W., Atta-Fynn, R., Autschbach, J., Bauman, N. P., Becca, J. C., Bernholdt, D. E., Bhaskaran-Nair, K., Bogatko, S., ... Harrison, R. J. (2020). NWChem: Past, present, and future. *The Journal of Chemical Physics*, 152(18), 184102. <https://doi.org/10.1063/5.0004997>
- Draine, B. T., & Flatau, P. J. (1994). Discrete-Dipole Approximation For Scattering Calculations. *Journal of the Optical Society of America A*, 11(4), 1491–1499. <https://doi.org/10.1364/JOSAA.11.001491>
- Ebbesen, T. W. (2016). Hybrid Light–Matter States in a Molecular and Material Science Perspective. *Accounts of Chemical Research*, 49(11), 2403–2412. <https://doi.org/10.1021/acs.accounts.6b00295>
- Filipovich, M. J., & Hughes, S. (2022). PyCharge: An open-source Python package for self-consistent electrodynamics simulations of Lorentz oscillators and moving point charges. *Computer Physics Communications*, 274, 108291. <https://doi.org/10.1016/j.cpc.2022.108291>
- Fojt, J., Rossi, T. P., Antosiewicz, T. J., Kuisma, M., & Erhart, P. (2021). Dipolar coupling of nanoparticle-molecule assemblies: An efficient approach for studying strong coupling. *The Journal of Chemical Physics*, 154(9), 094109. <https://doi.org/10.1063/5.0037853>
- Harris, C. R., Millman, K. J., van der Walt, S. J., Gommers, R., Virtanen, P., Cournapeau, D., Wieser, E., Taylor, J., Berg, S., Smith, N. J., Kern, R., Picus, M., Hoyer, S., van

- 133 Kerkwijk, M. H., Brett, M., Haldane, A., del Río, J. F., Wiebe, M., Peterson, P., ...
134 Oliphant, T. E. (2020). Array programming with NumPy. *Nature*, 585(7825), 357–362.
135 <https://doi.org/10.1038/s41586-020-2649-2>
- 136 Kuisma, M., Rousseaux, B., Czajkowski, K. M., Rossi, T. P., Shegai, T., Erhart, P., &
137 Antosiewicz, T. J. (2022). Ultrastrong coupling of a single molecule to a plasmonic
138 nanocavity: A first-principles study. *ACS Photonics*, 9(3), 1065–1077. [https://doi.org/10.](https://doi.org/10.1021/acsp Photonics.2c00066)
139 [1021/acsp Photonics.2c00066](https://doi.org/10.1021/acsp Photonics.2c00066)
- 140 Larsen, A. H., Mortensen, J. J., Blomqvist, J., Castelli, I. E., Christensen, R., Duřak, M.,
141 Friis, J., Groves, M. N., Hammer, B., Hargus, C., Hermes, E. D., Jennings, P. C.,
142 Jensen, P. B., Kermode, J., Kitchin, J. R., Kolsbjerg, E. L., Kubal, J., Kaasbjerg, K.,
143 Lysgaard, S., ... Jacobsen, K. W. (2017). The atomic simulation environment—a Python
144 library for working with atoms. *Journal of Physics: Condensed Matter*, 29(27), 273002.
145 <https://doi.org/10.1088/1361-648X/aa680e>
- 146 Mortensen, J. J., Larsen, A. H., Kuisma, M., Ivanov, A. V., Taghizadeh, A., Peterson, A., Haldar,
147 A., Dohn, A. O., Schäfer, C., Jónsson, E. Ö., Hermes, E. D., Nilsson, F. A., Kastlunger, G.,
148 Levi, G., Jónsson, H., Häkkinen, H., Fojt, J., Kangsabanik, J., Sødequist, J., ... Thygesen,
149 K. S. (2024). GPAW: An open Python package for electronic structure calculations. *The*
150 *Journal of Chemical Physics*, 160(9), 092503. <https://doi.org/10.1063/5.0182685>
- 151 Virtanen, P., Gommers, R., Oliphant, T. E., Haberland, M., Reddy, T., Cournapeau, D.,
152 Burovski, E., Peterson, P., Weckesser, W., Bright, J., van der Walt, S. J., Brett, M.,
153 Wilson, J., Millman, K. J., Mayorov, N., Nelson, A. R. J., Jones, E., Kern, R., Larson, E.,
154 ... van Mulbregt, P. (2020). SciPy 1.0: Fundamental algorithms for scientific computing in
155 Python. *Nature Methods*, 17(3), 261–272. <https://doi.org/10.1038/s41592-019-0686-2>
- 156 Wiecha, P. R. (2018). pyGDM—A python toolkit for full-field electro-dynamical simulations
157 and evolutionary optimization of nanostructures. *Computer Physics Communications*, 233,
158 167–192. <https://doi.org/10.1016/j.cpc.2018.06.017>
- 159 Wiecha, P. R., Majorel, C., Arbouet, A., Patoux, A., Brûlé, Y., Colas des Francs, G., & Girard,
160 C. (2022). “pyGDM” - new functionalities and major improvements to the python toolkit
161 for nano-optics full-field simulations. *Computer Physics Communications*, 270, 108142.
162 <https://doi.org/10.1016/j.cpc.2021.108142>