

DAA Taxa

0.1 Differential abundance analysis

- Group variable: **diet**
- Formula used: $\sim \text{diet} * \text{time} + (1 | \text{id})$

Table 1: Table of significant features for species - Total features analyzed: 215

feature	metadata	value	coef	qval_individual	qval_joint	model
Clostridiales_bacterium_KLE1615	diet	rice	-3.024	0.092	0.093	abundance
Clostridiales_bacterium_KLE1615	diet	rice:time	1.756	0.092	0.093	abundance
GGB34797_SGB14322	diet	rice	-3.154	0.092	0.093	abundance
Lawsonibacter_asaccharolyticus	time	time	-3.621	0.092	0.093	prevalence
GGB34797_SGB14322	diet	rice	15.979	0.184	0.093	prevalence
Lawsonibacter_asaccharolyticus	time	time	-0.520	0.559	0.093	abundance
Clostridiales_bacterium_KLE1615	diet	rice	1.358	1.000	0.093	prevalence
Clostridiales_bacterium_KLE1615	diet	rice:time	0.000	1.000	0.093	prevalence

Table 2: Table of significant features for genus - Total features analyzed: 139

feature	metadata	value	coef	qval_individual	qval_joint	model
Clostridiaceae_unclassified	diet	rice:time	1.078	0.066	0.06	abundance
Eubacteriales_unclassified	diet	rice:time	1.164	0.066	0.06	abundance
GGB34797	diet	rice	-3.154	0.066	0.06	abundance
Evtepia	diet	rice	15.971	0.066	0.06	prevalence
Lacrimispora	diet	rice:time	-7.649	0.066	0.06	prevalence
Lawsonibacter	diet	rice	-14.254	0.066	0.06	prevalence
Lawsonibacter	time	time	-3.621	0.066	0.06	prevalence
GGB34797	diet	rice:time	1.705	0.092	0.06	abundance
GGB34797	diet	rice	15.979	0.161	0.06	prevalence
Lawsonibacter	time	time	-0.520	0.399	0.06	abundance
Lacrimispora	diet	rice:time	0.901	0.424	0.06	abundance
Lawsonibacter	diet	rice	-1.156	0.596	0.06	abundance
Evtepia	diet	rice	-0.371	1.000	0.06	abundance
Clostridiaceae_unclassified	diet	rice:time	-1.318	1.000	0.06	prevalence
Eubacteriales_unclassified	diet	rice:time	0.039	1.000	0.06	prevalence
GGB34797	diet	rice:time	0.000	1.000	0.06	prevalence

Table 3: Table of significant features for family - Total features analyzed: 68

feature	metadata	value	coef	qval_individual	qval_joint	model
Eubacteriaceae	diet	rice:time	-9.428	0.000	0.000	prevalence
Clostridiales_unclassified	time	time	-3.621	0.029	0.031	prevalence

Table 3: Table of significant features for family - Total features analyzed: 68 (continued)

feature	metadata	value	coef	qval_individual	qval_joint	model
FGB12107	diet	rice	-3.154	0.037	0.039	abundance
Clostridiales_unclassified	diet	rice	-14.254	0.051	0.044	prevalence
FGB12107	diet	rice:time	1.705	0.070	0.044	abundance
FGB12107	diet	rice	15.979	0.129	0.039	prevalence
Clostridiales_unclassified	time	time	-0.520	0.481	0.031	abundance
Clostridiales_unclassified	diet	rice	-1.156	0.689	0.044	abundance
Eubacteriaceae	diet	rice:time	0.517	0.742	0.000	abundance
FGB12107	diet	rice:time	0.000	1.000	0.044	prevalence

Table 4: Table of significant features for phylum - Total features analyzed: 10

feature	metadata	value	coef	qval_individual	qval_joint	model
*						

1 species






Figure 1: taxa_image18


2 family





3 genus




4 phylum







5 Associations-species





6 Associations-family





7 Associations-genus

