

Differential abundance analysis

Analysis information:

- Taxonomic level: genus_prevalent

Taxa that are less prevalent than this (in relabundance) have been agglomerated:

- Detection threshold: 0.1%
- Prevalence threshold: 10%
- Number of taxonomic groups: 139

Number of significant findings within time points (Kruskal test, Benjamini-Hochberg FDR adjustment).

before	after
0	5

Number of significant findings within treatments (Paired Wilcoxon test, Benjamini-Hochberg FDR adjustment).



rice	oat
5	2

Let us look more closely to the significant hits. The logFC corresponds to logFC in CLR (tp2-tp1).



For a complete table of results, see [complete results](#)

Group	Taxon	logFC.clr	FDR	mean.rel1	mean.rel2
rice	Candidatus_Cibionibacter	-1.1231	0.0000	2.336	0.822
oat	GGB27106	1.6815	0.1020	0.019	0.115
oat	GGB4569	0.7232	0.1020	0.029	0.090
rice	Bifidobacterium	-0.7623	0.1128	6.265	4.463
rice	GGB34797	0.4275	0.1176	0.355	0.851
rice	Akkermansia	-1.1755	0.1176	1.147	1.116
rice	GGB3612	1.5879	0.1570	0.017	0.043
oat	Candidatus_Cibionibacter	-0.2537	0.4231	1.548	1.014
rice	GGB27106	-0.8557	0.5326	0.094	0.050
oat	Akkermansia	-0.2480	0.7113	0.704	0.904
oat	GGB3612	-0.3678	0.8582	0.046	0.061
oat	Bifidobacterium	0.2431	0.9049	3.874	3.871
oat	GGB34797	-0.0360	0.9351	0.484	0.410
rice	GGB4569	0.2229	0.9905	0.239	0.479



Akkermansia / p = NA





Bifidobacterium / p = NA





Candidatus_Cibionibacter / p = NA





GGB27106 / p = NA



GGB34797 / p = NA



GGB3612 / p = NA



GGB4569 / p = NA

oat/ p = 0.933

rice/ p = NA

