

# Differential abundance analysis

Analysis information:

- Taxonomic level: genus\_prevalent

Taxa that are less prevalent than this (in relabundance) have been agglomerated:

- Detection threshold: 0.001
- Prevalence threshold: 0.1
- Number of taxonomic groups: 139


Number of significant findings within time points (Kruskal test, Benjamini-Hochberg FDR adjustment).

before after  
0 5

Number of significant findings within treatments (Paired Wilcoxon test, Benjamini-Hochberg FDR adjustment).

rice oat  
5 2

Let us look more closely to the significant hits. The logFC corresponds to logFC in CLR (tp2-tp1).



For a complete table of results, see [complete results](#)

| Group | Taxon                    | logFC.clr | FDR    | mean.rel1 | mean.rel2 |
|-------|--------------------------|-----------|--------|-----------|-----------|
| rice  | Candidatus_Cibionibacter | -1.1231   | 0.0000 | 2.336     | 0.822     |
| oat   | GGB27106                 | 1.6815    | 0.1020 | 0.019     | 0.115     |
| oat   | GGB4569                  | 0.7232    | 0.1020 | 0.029     | 0.090     |
| rice  | Bifidobacterium          | -0.7623   | 0.1128 | 6.265     | 4.463     |
| rice  | GGB34797                 | 0.4275    | 0.1176 | 0.355     | 0.851     |
| rice  | Akkermansia              | -1.1755   | 0.1176 | 1.147     | 1.116     |
| rice  | GGB3612                  | 1.5879    | 0.1570 | 0.017     | 0.043     |

