

Differential abundance analysis

Analysis information:

- Taxonomic level: species_prevalent

Taxa that are less prevalent than this (in relabundance) have been agglomerated:

- Detection threshold: 0.001
- Prevalence threshold: 0.1
- Number of taxonomic groups: 215

Number of significant findings within time points (Kruskal test, Benjamini-Hochberg FDR adjustment).

```
before  after
      0      5
```

Number of significant findings within treatments (Paired Wilcoxon test, Benjamini-Hochberg FDR adjustment).

```
rice  oat
    16   2
```

Let us look more closely to the significant hits. The logFC corresponds to logFC in CLR (tp2-tp1).

For a complete table of results, see [complete results](#)

Group	Taxon	logFC.clr	FDR	mean.rel1	mean.rel2
rice	Candidatus_Cibionibacter_quicibialis	-1.0893	0.0002	2.336	0.822
rice	Bifidobacterium_longum	-0.8433	0.0034	1.794	1.311
rice	Blautia_sp_Marseille_P3087	0.7361	0.0295	0.016	0.054
rice	Eubacterium_siraeum	-1.7692	0.0516	0.262	0.205
rice	Bilophila_wadsworthia	0.7231	0.0796	0.066	0.096
rice	GGB34797_SGB14322	0.4614	0.0796	0.355	0.851
rice	GGB79734_SGB15291	0.4818	0.1168	0.116	0.178
rice	Firmicutes_bacterium_AF16_15	0.4403	0.1866	0.317	0.627
rice	Blautia_SGB4815	0.2757	0.1866	0.151	0.166
oat	GGB27106_SGB6188	1.6089	0.1929	0.019	0.115
oat	Blautia_massiliensis	0.6438	0.1929	1.169	2.183
rice	Streptococcus_salivarius	1.2865	0.2221	0.070	0.176
rice	GGB14025_SGB14906	0.5171	0.2221	0.040	0.064
rice	Clostridiaceae_bacterium	0.3150	0.2221	0.648	1.422
rice	Bacteroides_cellulosilyticus	0.3472	0.2316	0.224	0.446
rice	Alistipes_communis	-0.4789	0.2316	0.295	0.135
rice	GGB9760_SGB15374	0.7116	0.2339	0.049	0.109
rice	Akkermansia_muciniphila	-0.8478	0.2339	1.110	1.113



