

Differential abundance analysis

0.1 Differential abundance analysis

Analysis information:

- Taxonomic level: genus_prevalent

Taxa that are less prevalent than this (in relabundance) have been agglomerated:

- Detection threshold: 0.1%
- Prevalence threshold: 0.1
- Number of taxonomic groups: 139

Number of significant findings within time points (Kruskal test, Benjamini-Hochberg FDR adjustment).


before	after
0	5

Number of significant findings within treatments (Paired Wilcoxon test, Benjamini-Hochberg FDR adjustment).

rice	oat
5	2

Let us look more closely to the significant hits. The logFC corresponds to logFC in CLR (tp2-tp1).

Group	Taxon	logFC.clr	FDR	mean.relundance1	mean.relundance2
rice	Candidatus_Cibionibacter	-1.1231	0.0000	2.336	0.822
oat	GGB27106	1.6815	0.1020	0.019	0.115
oat	GGB4569	0.7232	0.1020	0.029	0.090
rice	Bifidobacterium	-0.7623	0.1128	6.265	4.463
rice	GGB34797	0.4275	0.1176	0.355	0.851
rice	Akkermansia	-1.1755	0.1176	1.147	1.116
rice	GGB3612	1.5879	0.1570	0.017	0.043



Volcano Plot: Effect Size vs Significance

