

Describing the population experiencing COVID-19 vaccine breakthrough following second vaccination in England: A cohort study from OpenSAFELY

The OpenSAFELY Collaborative: Amelia Green¹, Helen Curtis¹, William Hulme¹, Elizabeth Williamson², Helen McDonald², Krishnan Bhaskaran², Christopher Rentsch², Anna Schultze², Brian MacKenna¹, Viyaasan Mahalingasivam², Laurie Tomlinson², Alex Walker¹, Louis Fisher¹, Jon Massey¹, Colm Andrews¹, Lisa Hopcroft¹, Caroline Morton¹, Richard Croker¹, Jessica Morley¹, Amir Mehrkar¹, Seb Bacon¹, David Evans¹, Peter Inglesby¹, George Hickman¹, Tom Ward¹, Simon Davy¹, Rohini Mathur², John Tazare², Rosalind M Eggo², Kevin Wing², Angel Wong², Harriet Forbes², Chris Bates³, Jonathan Cockburn³, John Parry³, Frank Hester³, Sam Harper³, Ian Douglas², Stephen Evans², Liam Smeeth², Ben Goldacre¹

¹ The DataLab, Nuffield Department of Primary Care Health Sciences, University of Oxford, OX26GG

² London School of Hygiene and Tropical Medicine, Keppel Street, London WC1E 7HT

³ TPP, TPP House, 129 Low Lane, Horsforth, Leeds, LS18 5PX

Abstract

Background

While the vaccines against COVID-19 are highly effective, COVID-19 vaccine breakthrough is possible despite being fully vaccinated. With SARS-CoV-2 variants still circulating, describing the characteristics of individuals who have experienced COVID-19 vaccine breakthroughs could be hugely important in helping to determine who may be at greatest risk.

Method

With the approval of NHS England we conducted a retrospective cohort study using routine clinical data from the OpenSAFELY-TPP database of fully vaccinated individuals, linked to secondary care and death registry data, and described the characteristics of those experiencing COVID-19 vaccine breakthroughs.

Results

As of 01st November 2021, a total of 15,501,550 individuals were identified as being fully vaccinated against COVID-19, with a median follow-up time of 149 days (IQR: 107-179). From within this population, a total of 579,780 (<4%) individuals reported a positive SARS-CoV-2 test. For every 1,000 years of patient follow-up time, the corresponding incidence rate (IR) was 98.06 (95% CI 97.93-98.19). There were 28,580 COVID-19-related hospital admissions, 1,980 COVID-19-related critical care admissions and 6,435 COVID-19-related deaths; corresponding IRs 4.77 (95% CI 4.74-4.80), 0.33 (95% CI 0.32-0.34) and 1.07 (95% CI 1.06-1.09), respectively. The highest rates of breakthrough COVID-19 were seen in those in care homes and in patients with chronic kidney disease, dialysis, transplant, haematological malignancy or who were immunocompromised.

Conclusion

While the majority of COVID-19 vaccine breakthrough cases in England were mild, some differences in rates of breakthrough cases have been identified in several clinical groups. While it is important to note that these findings are simply descriptive and cannot be used to answer why certain groups have higher rates of COVID-19 breakthrough than others, the emergence of the Omicron variant of COVID-19 coupled with the number of positive SARS-CoV-2 tests still occurring is concerning and as numbers of fully vaccinated (and boosted) individuals increases and as follow-up time lengthens, so too will the number of COVID-19 breakthrough cases. Additional analyses, to assess vaccine waning and rates of breakthrough COVID-19 between different variants, aimed at identifying individuals at higher risk, are needed.

Keywords

COVID-19; Vaccine breakthrough; EHR data

Background

The vaccination programme for COVID-19 in the United Kingdom (UK) was started on 8 December 2020. Vaccination was in order of priority groups determined by the Joint Committee on Vaccination and Immunisation (JCVI) expert advisory group [1]: initially to people aged 80+ years, health and care workers and care home residents, people aged 70-79 and those extremely clinically vulnerable, followed by remaining adults in order of decreasing age or at increased risk. The Medicines and Healthcare products Regulatory Agency (MHRA) has so far approved four COVID-19 vaccines for use in the United Kingdom (UK), with each of the three vaccines currently in use requiring two doses to produce maximum protection [2]. To allow a higher percentage of the population to receive one vaccine dose quicker, second doses initially followed a 12-week interval from the first. This interval has since been shortened. These vaccines and their date of first administration and current second dose schedule are as follows: the Pfizer-BioNTech BNT162b2 mRNA COVID-19 vaccine (BNT162b2; first administered 8 Dec 2020, second dose at least 21 days later); the Oxford-AstraZeneca ChAdOx1 nCoV-19 vaccine (ChAdOx1; first administered 4 Jan 2021, second dose between 4 and 12 weeks later); the Moderna mRNA-1273 vaccine (mRNA-1273; first administered 7 April 2021, second dose 28 days later); and the Janssen Ad26.COV2.S vaccine (Ad26.COV2.S; not yet administered). As of November 1st, 2021 78.7% of individuals aged over 16 years in England had been fully vaccinated (i.e., ≥14 days have passed since the receipt of their second dose of a COVID-19 vaccine) [3].

The vaccines against COVID-19 are considered to be highly effective and there is strong evidence that the UK's COVID-19 vaccination programme has reduced infection and severe outcomes in vaccinated individuals [4, 5][4, 6], [4, 5]. However, breakthrough infections after vaccination against SARS-CoV-2 are increasingly reported [7] and there are concerns that the effectiveness of the vaccines may fade over time, due to the vaccines being less effective against the delta variant, and/or the waning of the immune response [8]. In addition, as no vaccine is 100% effective, COVID-19 vaccine breakthrough is likely (i.e., a small number of people will still get sick, be hospitalised, or die from COVID-19, despite being fully vaccinated). Describing individuals who have experienced a COVID-19 vaccine breakthrough could provide the first indication that the COVID-19 vaccine is less effective in certain groups and could be hugely important in helping to determine who may be at greatest risk and therefore might benefit most from booster doses.

To that end we used the new secure data analytics platform, OpenSAFELY [9] (built by our group on behalf of NHS England to support analysis of important questions related to COVID-19) to: describe breakthrough COVID-19 among fully vaccinated individuals in England; and to describe how breakthrough COVID-19 varied between priority groups and by clinical and demographic characteristics.

Methods

Data sources

This OpenSAFELY study was conducted using the OpenSAFELY-TPP database which contains records for approximately 24 million people currently registered with GP surgeries using TPP SystmOne software (approximately 40% of the English population).

Study population

The base population consisted of all individuals aged 16 years or over with records indicating that they had received two COVID-19 vaccination doses since 8 December 2020 (the earliest vaccination date in England) and who were still alive and registered 2 weeks after their second dose. Individuals were excluded if they tested positive for SARS-CoV-2 or had been hospitalised or died due to COVID-19 within the 2 weeks after their second dose. Follow up started 14 days after an individual's second dose of the COVID-19 vaccination (the point by which individuals were classified as being fully vaccinated) and individuals were followed up until 01st November 2021 or until the first occurrence of the outcome of interest; death; practice de-registration.

Outcomes

Four outcomes were assessed: positive SARS-CoV-2 swab test, via SGSS and based on swab date (we did not distinguish between symptomatic and asymptomatic infection for this outcome, nor PCR and lateral flow testing); COVID-19 related hospital admissions via Hospital Episode Statistics (HES) in-patient records (using both primary and non-primary diagnosis codes); COVID-19 related critical care admissions via HES and COVID-19-related death, via linked death registry records, which included individuals who died within 28-days of positive SARS-CoV-2 test or who had COVID-19 mentioned on the death certificate as one of the causes. COVID-19 hospitalisation variables were derived using two ICD-10 COVID-19 diagnosis codes; U071, U072 [10]. Outcomes were only included if they occurred 14 days or more after an individual's second dose of a COVID-19 vaccination.

Priority groups for vaccination

Individuals were classified into seven priority groups (box 1), based on the Joint Committee on Vaccination and Immunisation (JCVI) priority groups [11] using SNOMED-CT codelists and logic defined in the national COVID-19 Vaccination Uptake Reporting Specification developed by PRIMIS [12]. Individuals were assigned only to their highest priority group and not included again as part of any other priority group. In line with the national reporting specification, most criteria were ascertained using the latest available data at the time of analysis, with the exception of age which was calculated as at 31 March 2021 as recommended by Public Health England.

Box 1 – Priority groups for vaccination.

JCVI Risk group	Priority group	
	Description	Name
1	Residents in a care home for older adults, aged 65+	Care home
2	All those 80 years of age and over	80+
	Health and social care workers	Health/Care workers
3	All those 70 years of age and over	70-79
4	Clinically extremely vulnerable individuals	CEV
5,7,8,9	All those 50 years of age and over	50-69
6	“At Risk” individuals, some misclassified health/care workers and care home residents (except those age 70+ or shielding who will be included in another group), carers, household contacts of those at increased risk, and those not in priority groups who had the vaccine opportunistically, e.g. by responding to calls to use up excess doses.	Others not in the above groups

Key demographic and clinical characteristics

Key clinical and demographic groups which were considered to have a higher possibility of experiencing a COVID-19 vaccine breakthrough were identified from previous studies [13–17]. This included age, sex, body mass index (BMI; kg/m²), smoking status, deprivation (measured by the Index of Multiple Deprivation (IMD) as quintiles, ethnicity, NHS region of patient's general practice, asplenia, asthma, blood pressure, cancer, chronic kidney disease, diabetes mellitus, dialysis, heart disease, haematological malignancy, immunocompromised, learning disability, liver disease, neurological diseases, respiratory disease, severe mental illness and transplant. Other variables considered were time since being fully vaccinated, time between vaccinations and any evidence of a prior SARS-CoV-2 infection.

Codelists and implementation

Information on all **variables** were obtained from primary care records by searching TPP SystmOne records for specific coded data. Detailed information on compilation and sources for every individual codelist is available at <https://codelists.opensafely.org/> and the lists are available for inspection and re-use by the broader research community.

Missing data

Individuals with missing ethnicity, IMD and region were included as “Unknown”.

Statistical methods

Simple descriptive statistics were used to summarise the counts and rates of COVID-19 vaccine breakthrough. Rates for each outcome were estimated by dividing the count by

person-years, with 95% confidence intervals. Counts and rates were stratified by initial priority groups for all outcomes and by selected clinical and demographic groups.

Software and Reproducibility

Data management and analysis was performed using the OpenSAFELY software libraries, Python 3 and R version 4.0.2. All code for the OpenSAFELY platform for data management, analysis and secure code execution is shared for review and re-use under open licenses at [GitHub.com/OpenSAFELY](https://github.com/OpenSAFELY). All code for data management and analysis for this paper is shared for scientific review and re-use under open licenses on GitHub (<https://github.com/opensafely/covid-19-vaccine-breakthrough>).

Patient and Public Involvement

Any patient or member of the public is invited to contact us at <https://opensafely.org/> regarding this study or the broader OpenSAFELY project.

Results

Study population

Out of approximately 24 million patients, 15,501,550 individuals were identified as being fully vaccinated against COVID-19 and included in this study (Figure 1). The median follow-up time was 149 days (interquartile range: 107 - 179 days). Since being fully vaccinated, 8,370,837 (54%) of the base population had at least one record for a (positive/negative) SARS-CoV-2 test recorded, with a positivity rate of 3.17%. Testing behaviours varied between priority groups with individuals in care homes testing more regularly than other groups; 91% of care home residents had 3+ SARS-CoV-2 tests since being fully vaccinated vs 41%-73% in other groups. The total number of COVID-19 vaccine breakthrough cases was 590,279 (<4%). Table 1 shows the number and rate of individuals fully vaccinated broken down by initial JCVI priority groups, along with the number and rate (per 1,000 patient years) of each outcome.

Positive SARS-CoV-2 test

In fully vaccinated individuals, the median number of days to a positive test for SARS-CoV-2 was 99 (IQR 63 - 134 days) with a total of 579,780 individuals testing positive for SARS-CoV-2 (about 1 in 25 or <4%). For every 1,000 years of patient follow-up time, the corresponding incidence rate was 98.06 (95% CI 97.93-98.19). Within the initial JCVI priority groups, positive SARS-CoV-2 test rates were highest in the CEV group (107.84, 95% CI 107.27-108.42) and lowest in those over 80 years of age (26.69, 95% CI 26.48-26.90). The overall cumulative incidence of positive SARS-CoV-2 tests at 275 days from the study start date was <0.1% (Figure 2).

When broken down into clinical/demographic groups and comorbidities (Table 2), rates of individuals testing positive for SARS-CoV-2 were highest in 40-49 year olds (179.96, 95% CI 179.50-180.42) and lowest in individuals over 80 years of age (28.90, 95% CI 28.69-29.11). Rates were higher in females than in males: 102.60 (95% CI 102.42-102.78) vs 92.76 (95% CI 92.58-92.95), respectively. Comorbidities with the highest rates of positive SARS-CoV-2

tests included kidney transplant, dialysis and immunocompromised; 143.51 (95% CI 136.46-150.93), 134.75 (95% CI 124.32-146.06), 91.53 (95% CI 90.91-92.15), respectively. Rates of positive SARS-CoV-2 tests were lowest 4-8 weeks after being fully vaccinated and highest at 12+ weeks; 48.65 (95% CI 47.97-49.33) vs 100.3 (95% CI 100.16-100.43), respectively.

COVID-19 related hospital admission

From within the fully vaccinated population, 28,580 had a COVID-19 related hospital admission (about 1 in 550 or 0.18%), with a median time to hospitalisation of 143 days (IQR 102-174). Of those who had a COVID-19 related hospital admission, 18,800 (66%) had a positive SARS-CoV-2 test in their records; 9,170 (49%) occurred prior to an individual being hospitalised, 7030 (37%) occurred within the first 2 days of their hospital admission, 2365 (13%) occurred between 3 and 29 days after being hospitalised and 235 (1%) occurred 30 days or more after an individual had been hospitalised.

Rates of COVID-19 related hospital admissions increased with age; 1.09 (1.84-1.96) for 16-29 year olds vs 12.75 (12.61-12.89) for those over 80 years, respectively. Rates were higher in those who were more deprived; most deprived IMD quintile versus least deprived IMD quintile was 7.32 (7.23-7.41) vs 3.55 (3.5-3.61), respectively. Comorbidities with the highest rates of COVID-19 related hospital admissions included kidney transplant, dialysis, and chronic kidney disease; 76.08 (95% CI 71.03-81.49), 70.73 (95% CI 63.34-78.99), 49.49 (95% CI 45.33-54.02), respectively.

COVID-19 related critical care admission

Of the 28,580 COVID-19 related hospital admission, 1,980 needed to be admitted to critical care (about 1 in 1000 of fully vaccinated people or 1 in 14 of hospitalised patients), with a median time to critical care admission of 136 days (IQR 94-166). For every 1,000 years of patient follow-up time, the corresponding incidence rate was 0.33 (95% CI 0.32-0.34). Within the initial JCVI priority groups, COVID-19 related critical care admissions rates were highest in the CEV group (1.73, 95% CI 1.66-1.80) and lowest in health care workers 0.10 (95% CI 0.08-0.12).

Rates of COVID-19 related critical care admissions were higher in males than females and in; 0.45 (95% CI 0.44-0.46) vs 0.23 (95% CI 0.22-0.24), respectively. Rates were higher in those who were obese compared to not obese; BMI >30 (kg/m²) versus BMI <30 (kg/m²) was 0.68 (0.66-0.71) vs 0.22 (0.21-0.23), respectively. Comorbidities with the highest rates of COVID-19 related critical care admissions included dialysis, kidney transplant and chronic kidney disease; 19.74 (95% CI 16.03-24.32), 17.86 (95% CI 15.5-20.57) and 5.31 (95% CI 4.06-6.93), respectively.

COVID-19 related death

Of those who were fully vaccinated, 6,435 had a COVID-19 related death (about 1 in 2500 or 0.04%) with a median number of days to death of 165 (IQR 129-188). While the majority (43%) of deaths occurred in the 80+ JCVI priority group, COVID-19 related death rates were three times as high among individuals living in care homes compared to those over 80 years living in private residences; 13.37 (95% CI 12.86-13.91) vs 4.64 (95% CI 4.56-4.73), respectively.

There were very few deaths in those under 30 years of age with rates of COVID-19-related death increasing with increased age; 0.02 (95% CI 0.02-0.03) for 16-29 year olds vs 5.26 (95% CI 5.17-5.35) for those over 80 years, respectively. Comorbidities with the highest rates of COVID-19-related death included dialysis, transplant and chronic kidney disease; 25.71 (95% CI 21.42-30.86), 18.9 (95% CI 16.47-21.68), and 17.81 (95% CI 15.40-20.61), respectively.

Discussion

Summary

This descriptive analysis in over 15 million people living in England shows that COVID-19 vaccine breakthrough is occurring, however, events are currently rare and mostly mild in nature. Nevertheless, some individuals are experiencing higher rates of serious illness and death due to COVID-19, such as those in care homes, with chronic kidney disease, on dialysis, with a transplant or who are immunosuppressed. It is important to note that this analysis is a simple descriptive piece of analysis. While it is possible to adjust rates (as demonstrated for patients with chronic kidney disease in Supplementary Table S1) to help inform decisions around rollout of vaccine/booster programme for patients at high risk of adverse outcomes, it is important to stress that this study was not designed to estimate risk factors for COVID-19 vaccine breakthrough and thus cannot be used to answer why certain groups have higher rates of breakthrough cases than others or to estimate vaccine effectiveness.

Strengths and weaknesses

This study used large-scale, routinely-collected primary care records, linked to coronavirus testing surveillance, hospital, and death registry data. This allowed us to describe a substantial proportion of the English population in rich longitudinal detail and to detect variations in COVID-19 vaccine breakthrough cases, as early as possible.

We acknowledge several important limitations of these findings. First, even though the base population consisted of over 15 million fully vaccinated individuals the numbers of COVID-19 vaccine breakthrough cases were relatively small, especially for hospitalisations and deaths. This made comparisons between outcomes, specifically at selected clinical and demographic levels difficult, meaning rates could be imprecisely estimated. Second, due to the targeted roll out of the COVID-19 vaccination programme in England, this cohort represents mostly older and vulnerable populations. In addition, follow-up time is systematically different amongst individuals included in this study and no adjustment for this has been made. Third, asymptomatic testing patterns vary between individuals. Apart from health and care workers, asymptomatic individuals without any underlying health issues or comorbidities are less likely to get tested than those with underlying health issues or comorbidities (i.e., haemodialysis patients) who will undergo asymptomatic testing regularly. Most lateral flow tests conducted at home are not included in our data. The number of reported positive SARS-CoV-2 tests is therefore likely to be an undercount of SARS-CoV-2 among fully vaccinated individuals without any underlying health issues or comorbidities,

which may have led to underestimation of the corresponding rates. Fourth, individuals were excluded if they tested positive for SARS-CoV-2 or had been hospitalised or died due to COVID-19 within the two weeks after their second dose, we acknowledge that there is still the potential that some patients will have been included whose outcome was related to a previous infection pre second vaccination, in particular in some of the highest risk groups such as dialysis patients and those in care homes where there a high amount of repeated screening in, resulting in a potential increase in rates. Fifth, characteristics linked to COVID-19 vaccine breakthrough in fully vaccinated individuals may be reflective of higher infection rates regardless of vaccination in some groups, not because of vaccination (i.e. due to higher exposure due to behavioural differences) [18]. Finally, this study includes data from December 2020 until November 2021, during which time two different variants of SARS-COV-2 were dominating in England at different times; the Alpha variant (B.1.1.7) between January 2021 and May 2021, and the Delta variant (B.1.617.2) after June 2021[19]. These have since been supplanted by other variants, in particular the Omicron variant. This is a descriptive study and does not aim to quantify the contribution of vaccine breakthrough due to the change in dominant variant from Alpha to Delta or the extent of waning in the months immediately following vaccination. Further studies are needed to assess vaccine waning and rates of breakthrough COVID-19 between different variants and in newer, more prevalent variants, such as Omicron (B.1.1.529).

Findings in Context

There is strong evidence that the UK's COVID-19 vaccination programme has reduced infection and severe outcomes in vaccinated individuals[4, 5]. However, breakthrough infections after vaccination against SARS-CoV-2 are increasingly reported[7] and there are concerns that the effectiveness of the vaccines may reduce over time, due to the vaccines being less effective against the delta variant, and/or the waning of the immune response[8]. While we did find that breakthrough infections were more frequent in the earlier-vaccinated individuals, potentially due to waning vaccine efficacy or impaired immune response, further studies are needed to gain a clearer picture of the long-term effectiveness of COVID-19 vaccines. In addition, with only a handful of studies investigating risk factors for post-vaccination infection [13, 16, 17], very little is known about how breakthrough COVID-19 varies between key clinical and demographic groups. Our findings on COVID-19 vaccine breakthrough are consistent with patterns observed worldwide; COVID-19 vaccine breakthrough cases are rare and mild. However, there are potentially several groups who are at higher risk of COVID-19 vaccine breakthrough including those in care-homes, with chronic kidney disease, on dialysis, with transplants or who are immunocompromised.

Conclusion

As of 1st November 2021, the majority of COVID-19 vaccine breakthrough cases in England were mild with relatively smaller numbers of fully vaccinated individuals being hospitalised or dying as a result of COVID-19. While these numbers are in line with expectations, and while follow-up time is systematically different amongst fully vaccinated individuals and variation in vaccination coverage between groups and regions will have many complex drivers, some differences in rates of breakthrough cases were identified in several clinical groups. the emergence of the Omicron variant of COVID-19 coupled with the number of positive SARS-CoV-2 tests still occurring is concerning and as numbers of fully vaccinated (and

boosted) individuals increases and as follow-up time lengthens, so too will the number of COVID-19 breakthrough cases. Additional analyses are therefore, to assess vaccine waning and rates of breakthrough COVID-19 between different variants, and to enable identification of individuals at higher risk, who would require continued strict precautions, and additional vaccination.

Abbreviations

Body Mass index (BMI); Joint Committee Vaccination and Immunisation (JCVI); Index of Multiple Deprivation (IMD); United Kingdom (UK).

Declarations

Information governance and ethical approval

NHS England is the data controller; TPP is the data processor; and the key researchers on OpenSAFELY are acting with the approval of NHS England. This implementation of OpenSAFELY is hosted within the TPP environment which is accredited to the ISO 27001 information security standard and is NHS IG Toolkit compliant^[17,18]; Patient data has been pseudonymised for analysis and linkage using industry standard cryptographic hashing techniques; all pseudonymised datasets transmitted for linkage onto OpenSAFELY are encrypted; access to the platform is via a virtual private network (VPN) connection, restricted to a small group of researchers; the researchers hold contracts with NHS England and only access the platform to initiate database queries and statistical models; all database activity is logged; only aggregate statistical outputs leave the platform environment following best practice for anonymisation of results such as statistical disclosure control for low cell counts¹⁶. The OpenSAFELY research platform adheres to the obligations of the UK General Data Protection Regulation (GDPR) and the Data Protection Act 2018. In March 2020, the Secretary of State for Health and Social Care used powers under the UK Health Service (Control of Patient Information) Regulations 2002 (COPI) to require organisations to process confidential patient information for the purposes of protecting public health, providing healthcare services to the public and monitoring and managing the COVID-19 outbreak and incidents of exposure; this sets aside the requirement for patient consent¹⁷. Taken together, these provide the legal bases to link patient datasets on the OpenSAFELY platform. GP practices, from which the primary care data are obtained, are required to share relevant health information to support the public health response to the pandemic, and have been informed of the OpenSAFELY analytics platform.

This study was approved by the Health Research Authority (REC reference 20/LO/0651) and by the LSHTM Ethics Board (reference 21863).

Availability of data and materials

Access to the underlying identifiable and potentially re-identifiable pseudonymised electronic health record data is tightly governed by various legislative and regulatory frameworks, and restricted by best practice. The data in OpenSAFELY is drawn from General Practice data across England where TPP is the Data Processor. TPP developers (CB, RC, JP, FH, and SH) initiate an automated process to create pseudonymised records in the core OpenSAFELY database, which are copies of key structured data tables in the identifiable records. These are linked onto key external data resources that have also been pseudonymised via SHA-512 one-way hashing of NHS numbers using a shared salt. DataLab developers and PIs (BG, LS, CEM, SB, AJW, WH, DE, PI, and CTR) holding contracts with NHS England have access to the OpenSAFELY pseudonymised data tables as needed to develop the OpenSAFELY tools. These tools in turn enable researchers with OpenSAFELY Data Access Agreements to write and execute code for data management and data analysis without direct access to the underlying raw pseudonymised patient data, and to review the outputs of this code. All code for the full data management pipeline—from raw data to completed results for this analysis—and for the OpenSAFELY platform as a whole is available for review at github.com/OpenSAFELY.

Competing interests

All authors have completed the ICMJE uniform disclosure form and declare the following: BG has received research funding from Health Data Research UK (HDRUK), the Laura and John Arnold Foundation, the Wellcome Trust, the NIHR Oxford Biomedical Research Centre, the NHS National Institute for Health Research School of Primary Care Research, the Mohn-Westlake Foundation, the Good Thinking Foundation, the Health Foundation, and the World Health Organisation; he also receives personal income from speaking and writing for lay audiences on the misuse of science. IJD holds shares in GlaxoSmithKline (GSK).

Funding

This work was supported by the Medical Research Council MR/V015737/1 and the Longitudinal Health and wellbeing strand of the National Core Studies programme. The OpenSAFELY platform is funded by the Wellcome Trust. TPP provided technical expertise and infrastructure within their data centre pro bono in the context of a national emergency. BGs work on clinical informatics is supported by the NIHR Oxford Biomedical Research Centre and the NIHR Applied Research Collaboration Oxford and Thames Valley.

BG's work on better use of data in healthcare more broadly is currently funded in part by: NIHR Oxford Biomedical Research Centre, NIHR Applied Research Collaboration Oxford and Thames Valley, the Mohn-Westlake Foundation, NHS England, and the Health Foundation; all DataLab staff are supported by BG's grants on this work. LS reports grants from Wellcome, MRC, NIHR, UKRI, British Council, GSK, British Heart Foundation, and Diabetes UK outside this work. KB holds a Wellcome Senior Research Fellowship (220283/Z/20/Z). HIM is funded by the NIHR Health Protection Research Unit in Immunisation, a partnership between Public Health England and London School of Hygiene & Tropical Medicine. AYSW holds a fellowship from the British Heart Foundation. EJW holds grants from MRC. RM holds a Sir Henry Wellcome Fellowship funded by the Wellcome Trust (201375/Z/16/Z). HF holds a UKRI fellowship. IJD holds grants from NIHR and GSK.

Funders had no role in the study design, collection, analysis, and interpretation of data; in the writing of the report and in the decision to submit the article for publication.

The views expressed are those of the authors and not necessarily those of the NIHR, NHS England, Public Health England or the Department of Health and Social Care.

For the purpose of Open Access, the author has applied a CC BY public copyright licence to any Author Accepted Manuscript (AAM) version arising from this submission.

Authors' contributions

Contributions are as follows:

Conceptualisation: RM, CTR, KB, RME, LS, BG, BM, HJC, SJWE, KK, DH, KR

Funding acquisition: LS, BG, RME

Methodology: AG, HC, WH, EW, HMD, AW, LF, CA, LH, CM, BMK, RC, JM, AM, KB, AS, CR, AR, LF, LT, RM, AW, HF, SE, LS, BG

Formal analysis: AG, HC

Codelists: RM, LT, AS, AJW, CM, BG, WJH, SB, AM

Software: AW, CB, JC, DE, PI, CM, WH, BN, SB, HC, ND, RC, JP, FH, SH

Visualisation: AG, HC, WH, EW

Writing - original draft: AG

Writing- review & editing: ALL

Information governance: CB LS BG AM

Acknowledgements

We are very grateful for all the support received from the TPP Technical Operations team throughout this work, and for generous assistance from the information governance and database teams at NHS England / NHSX.

Guarantor

BG is guarantor.

References

1. COVID-19: the green book, chapter 14a. GOV.UK. 2020.
<https://www.gov.uk/government/publications/covid-19-the-green-book-chapter-14a>. Accessed 20 Oct 2021.
2. Dosing information COVID-19 primary vaccination. SPS - Specialist Pharmacy Service. 2021. <https://www.sps.nhs.uk/articles/dosing-information-covid-19-primary-vaccination/>. Accessed 1 Mar 2022.
3. Coronavirus (COVID-19) in the UK. GOV.UK.
<https://coronavirus.data.gov.uk/details/vaccinations>. Accessed 9 Sep 2021.
4. Lopez Bernal J, Andrews N, Gower C, Robertson C, Stowe J, Tessier E, et al. Effectiveness of the Pfizer-BioNTech and Oxford-AstraZeneca vaccines on covid-19 related symptoms, hospital admissions, and mortality in older adults in England: test negative case-control study. *BMJ*. 2021;373:n1088.
5. Lopez Bernal J, Andrews N, Gower C, Gallagher E, Simmons R, Thelwall S, et al. Effectiveness of Covid-19 Vaccines against the B.1.617.2 (Delta) Variant. *N Engl J Med*. 2021;385:585–94.
6. Bernal JL, Andrews N, Gower C, Gallagher E, Simmons R, Thelwall S, et al. Effectiveness of Covid-19 Vaccines against the B.1.617.2 (Delta) Variant. *New England Journal of Medicine*. 2021;385:585–94.
7. Bates TA, McBride SK, Winders B, Schoen D, Trautmann L, Curlin ME, et al. Antibody Response and Variant Cross-Neutralization After SARS-CoV-2 Breakthrough Infection. *JAMA*. 2022;327:179–81.
8. Krause PR, Fleming TR, Peto R, Longini IM, Figueroa JP, Sterne JAC, et al. Considerations in boosting COVID-19 vaccine immune responses. *Lancet*. 2021;398:1377–80.
9. OpenSAFELY. <https://www.opensafely.org/>. Accessed 20 Oct 2021.
10. Emergency use ICD codes for COVID-19 disease outbreak.
<https://www.who.int/classifications/classification-of-diseases/emergency-use-icd-codes-for-covid-19-disease-outbreak>. Accessed 20 Oct 2021.
11. UK Health Security Agency. COVID-19: the green book, chapter 14a. 2020.
<https://www.gov.uk/government/publications/covid-19-the-green-book-chapter-14a>. Accessed 20 Oct 2021.
12. COVID-19 Vaccination Uptake Reporting Specification. PRIMIS.
<https://www.nottingham.ac.uk/primis/covid-19/covid-19.aspx>. Accessed 20 Jun 2021.
13. Cook C, Patel NJ, D'Silva KM, Hsu TY-T, Dilorio M, Prisco L, et al. Clinical characteristics and outcomes of COVID-19 breakthrough infections among vaccinated patients with systemic autoimmune rheumatic diseases. *Ann Rheum Dis*. 2021.
<https://doi.org/10.1136/annrheumdis-2021-221326>.
14. Keehner J, Horton LE, Binkin NJ, Laurent LC, Pride D, Longhurst CA, et al. Resurgence of SARS-CoV-2 Infection in a Highly Vaccinated Health System Workforce. *N Engl J Med*. 2021;385:1330–2.

15. Oster Y, Benenson S, Yochi Harpaz L, Buda I, Nir-Paz R, Strahilevitz J, et al. Association Between Exposure Characteristics and the Risk for COVID-19 Infection Among Health Care Workers With and Without BNT162b2 Vaccination. *JAMA Netw Open*. 2021;4:e2125394.
16. Antonelli M, Penfold RS, Merino J, Sudre CH, Molteni E, Berry S, et al. Risk factors and disease profile of post-vaccination SARS-CoV-2 infection in UK users of the COVID Symptom Study app: a prospective, community-based, nested, case-control study. *Lancet Infect Dis*. 2021. [https://doi.org/10.1016/S1473-3099\(21\)00460-6](https://doi.org/10.1016/S1473-3099(21)00460-6).
17. Hippisley-Cox J, Coupland CA, Mehta N, Keogh RH, Diaz-Ordaz K, Khunti K, et al. Risk prediction of covid-19 related death and hospital admission in adults after covid-19 vaccination: national prospective cohort study. *BMJ*. 2021;374:n2244.
18. Day M. Covid-19: Stronger warnings are needed to curb socialising after vaccination, say doctors and behavioural scientists. *BMJ*. 2021;372.
19. British Heart Foundation. Covid variant. British Heart Foundation. 2022. <https://www.bhf.org.uk/information-support/heart-matters-magazine/news/coronavirus-and-your-health/covid-variant>. Accessed 1 Mar 2022.

Figures, tables and additional files

Table 1. Number of individuals fully vaccinated (2 doses + 2 weeks) in initial priority groups in OpenSAFELY-TTP, and number and rate with each outcome.

Priority Group	Fully vaccinated						Positive SARS-CoV-2 test †			COVID-19-related hospital admission †		COVID-19-related critical care admission †		COVID-19 related death †	
	Count*	Follow-up time, medium days (IQR)	Tests conducted (%)				Positivity (%)	Events* (PYs)	Incidence rate per 1000 person-years (95% CI)	Events* (PYs)	Incidence rate per 1000 person-years (95% CI)	Events* (PYs)	Incidence rate per 1000 person-years (95% CI)	Events* (PYs)	Incidence rate per 1000 person-years (95% CI)
			0	1	2	3+									
All	15501550	149 (107-179)	46	19	10	19	579780 (5912498)	3.17	98.06 (97.93-98.19)	28580 (5993535)	4.77 (4.74-4.8)	1980 (5996007)	0.33 (0.32-0.34)	6435 (5996178)	1.07 (1.06-1.09)
Care home (priority group 1)	98400	200 (180-207)	9	5	4	82	3110 (48207)	0.90	64.51 (63.37-65.68)	755 (48600)	15.58 (15.02-16.15)	REDACTED	REDACTED	650 (48674)	13.37 (12.86-13.91)
80+ (priority group 2)	1073145	200 (191-213)	59	14	7	20	15945 (597368)	1.96	26.69 (26.48-26.9)	7545 (598868)	12.60 (12.45-12.74)	180 (599613)	0.30 (0.28-0.32)	2785 (599626)	4.64 (4.56-4.73)
Health/care workers (priority groups 1-2)	555820	200 (182-211)	27	21	14	39	30045 (282813)	3.31	106.23 (105.62-106.84)	440 (287822)	1.53 (1.46-1.60)	30 (287862)	0.10 (0.08-0.12)	20 (287866)	0.07 (0.05-0.08)
70-79 (priority groups 3-4)	1990755	184 (176-191)	56	17	8	19	41325 (985544)	2.16	41.93 (41.73-42.14)	6875 (990497)	6.94 (6.86-7.03)	525 (991111)	0.53 (0.51-0.56)	1640 (991147)	1.66 (1.62-1.70)
CEV (age 16-69) (priority group 4)	725045	171 (157-181)	42	18	10	30	34820 (322885)	2.92	107.84 (107.27-108.42)	4560 (327932)	13.90 (13.7-14.11)	565 (328308)	1.73 (1.66-1.80)	785 (328361)	2.39 (2.31-2.48)
50 - 69 (priority groups 5-9)	4730155	151 (138-166)	47	18	9	26	184890 (1946451)	2.27	94.99 (94.77-95.21)	5075 (1973346)	2.57 (2.54-2.61)	515 (1973743)	0.26 (0.25-0.27)	495 (1973791)	0.25 (0.24-0.26)
Others not in the above groups***	15501550	149 (107-179)	46	19	10	19	579780 (5912498)	3.17	98.06 (97.93-98.19)	28580 (5993535)	4.77 (4.74-4.8)	1980 (5996007)	0.33 (0.32-0.34)	6435 (5996178)	1.07 (1.06-1.09)

[†] Calculated as of 01st November 2021

*All counts (of people and events) have been rounded to the nearest 5 and counts < 8 have been redacted

**This is a count of (positive and negative) test results and may include multiple tests for an individual person

***Others includes: others in priority groups ("At Risk"), some misclassified health/care workers and care home residents (except those age 70+ or shielding who will be included in another group), carers, household contacts of those at increased risk, and those not in priority groups who had the vaccine opportunistically, e.g. by responding to calls to use up excess doses.

Table 2 Count and rates of breakthrough positive SARS-CoV-2 tests and hospitalisation and death in fully vaccinated individuals in OpenSAFELY-TPP, broken down by selected clinical and demographic groups

Clinical/ demographic group	Category	Fully vaccinated						Positive SARS-CoV-2 test			Hospitalised with COVID-19		COVID-19-related critical care admission		COVID-19 related death	
		Count*	Follow-up time, medium days (IQR)	Tests conducted (%)				Positivity (%)	Events* (PYs)	Incidence rate per 1000 person-years (95% CI)	Events* (PYs)	Incidence rate per 1000 person-years (95% CI)	Events* (PYs)	Incidence rate per 1000 person-years (95% CI)	Events* (PYs)	Incidence rate per 1000 person-years (95% CI)
				0	1	2	3+									
Age	16 - 29	2227120	66 (47-129)	45	21	11	22	63620 (516289)	5.47	123.22 (122.74-123.71)	1000 (527073)	1.90 (1.84-1.96)	40 (527145)	0.07 (0.06-0.08)	10 (527150)	0.02 (0.01-0.02)
	30 - 39	2289165	94 (72-145)	42	21	12	25	93375 (660313)	4.50	141.41 (140.95-141.88)	1550 (673014)	2.30 (2.25-2.36)	80 (673118)	0.12 (0.10-0.13)	35 (673127)	0.05 (0.04-0.06)
	40 - 50	2459075	126 (103-155)	40	20	11	29	153210 (851368)	3.86	179.96 (179.5-180.42)	2320 (871625)	2.66 (2.61-2.72)	210 (871799)	0.24 (0.22-0.26)	120 (871821)	0.14 (0.12-0.15)
	50 - 59	2942440	146 (135-163)	43	19	10	28	136470 (1188958)	2.49	114.78 (114.47-115.09)	3850 (1209351)	3.18 (3.13-3.23)	420 (1209663)	0.35 (0.33-0.37)	370 (1209705)	0.31 (0.29-0.32)
	60 - 69	2421245	163 (150-174)	49	18	9	24	72735 (1064063)	1.94	68.36 (68.1-68.61)	4715 (1074117)	4.39 (4.33-4.45)	530 (1074498)	0.49 (0.47-0.52)	845 (1074542)	0.78 (0.76-0.81)
	70 - 79	2016365	184 (176-191)	56	17	8	19	42075 (998433)	2.11	42.14 (41.93-42.34)	7050 (1003480)	7.03 (6.94-7.11)	530 (1004114)	0.53 (0.51-0.55)	1720 (1004149)	1.71 (1.67-1.76)
	80 +	1146150	200 (191-213)	55	14	7	24	18295 (633074)	1.72	28.9 (28.69-29.11)	8095 (634874)	12.75 (12.61-12.89)	180 (635670)	0.29 (0.27-0.31)	3345 (635683)	5.26 (5.17-5.35)
Sex	Female	8072115	155 (116-184)	43	19	10	28	326565 (3182819)	2.69	102.6 (102.42-102.78)	13500 (3228984)	4.18 (4.14-4.22)	740 (3230164)	0.23 (0.22-0.24)	2655 (3230233)	0.82 (0.81-0.84)
	Male	7429435	144 (99-174)	50	19	10	21	253215 (2729679)	4.04	92.76 (92.58-92.95)	15080 (2764551)	5.46 (5.41-5.50)	1250 (2765843)	0.45 (0.44-0.46)	3780 (2765945)	1.37 (1.34-1.39)
BMI (kg/m2)	Not obese	12078955	145 (98-178)	46	19	10	25	429315 (4498201)	3.20	95.44 (95.3-95.59)	18330 (4558086)	4.02 (3.99-4.05)	1010 (4559694)	0.22 (0.21-0.23)	4475 (4559783)	0.98 (0.97-1.00)
	Obese (>30)	3422600	159 (135-181)	46	19	10	26	150465 (1414297)	3.07	106.39 (106.11-106.66)	10255 (1435449)	7.14 (7.07-7.21)	980 (1436313)	0.68 (0.66-0.71)	1965 (1436395)	1.37 (1.34-1.40)
Smoking	Non-smoker and unknown	7833690	143 (91-176)	45	19	10	26	291000 (2835776)	3.19	102.62 (102.43-102.81)	9960 (2876640)	3.46 (3.43-3.50)	675 (2877506)	0.24 (0.23-0.24)	1860 (2877571)	0.65 (0.63-0.66)
	Current and former	7667865	156 (125-182)	47	19	10	24	288780 (3076722)	3.14	93.86 (93.68-94.03)	18620 (3116894)	5.97 (5.93-6.02)	1310 (3118501)	0.42 (0.41-0.43)	4575 (3118607)	1.47 (1.45-1.49)

Clinical/ demographic group	Category	Fully vaccinated						Positive SARS-CoV-2 test			Hospitalised with COVID-19		COVID-19-related critical care admission		COVID-19 related death	
		Count*	Follow-up time, medium days (IQR)	Tests conducted (%)				Positivity (%)	Events* (PYs)	Incidence rate per 1000 person-years (95% CI)	Events* (PYs)	Incidence rate per 1000 person-years (95% CI)	Events* (PYs)	Incidence rate per 1000 person-years (95% CI)	Events* (PYs)	Incidence rate per 1000 person-years (95% CI)
				0	1	2	3+									
Ethnicity	White	11895195	154 (119-182)	44	19	10	26	466170 (4688366)	3.08	99.43 (99.29-99.58)	25500 (4753382)	5.36 (5.33-5.40)	1760 (4755592)	0.37 (0.36-0.38)	5905 (4755742)	1.24 (1.23-1.26)
	Mixed	151275	129 (76-164)	44	20	11	25	4980 (50030)	3.78	99.56 (98.16-100.98)	195 (50764)	3.82 (3.56-4.11)	20 (50779)	0.35 (0.28-0.45)	25 (50780)	0.51 (0.42-0.62)
	Asian or Asian British	901520	134 (81-170)	55	19	10	17	29830 (309199)	4.24	96.48 (95.92-97.04)	1735 (313404)	5.54 (5.40-5.67)	120 (313565)	0.39 (0.36-0.43)	300 (313577)	0.95 (0.90-1.01)
	Black or Black British	224740	138 (87-172)	49	18	9	23	5930 (78696)	2.98	75.37 (74.39-76.35)	415 (79581)	5.20 (4.95-5.46)	25 (79614)	0.34 (0.28-0.41)	50 (79616)	0.63 (0.55-0.72)
	Other ethnic groups	242335	122 (79-159)	50	19	10	20	5970 (78690)	3.66	75.85 (74.88-76.84)	295 (79509)	3.69 (3.48-3.91)	30 (79531)	0.38 (0.31-0.45)	55 (79534)	0.67 (0.58-0.76)
	Unknown	2086480	135 (82-164)	53	18	9	21	66900 (707517)	3.36	94.56 (94.19-94.92)	445 (716894)	0.62 (0.59-0.65)	30 (716926)	0.04 (0.03-0.05)	105 (716928)	0.14 (0.13-0.16)
IMD quintile	1 (most deprived)	2519265	146 (100-176)	54	18	9	20	96800 (927354)	3.56	104.39 (104.05-104.72)	6890 (941680)	7.32 (7.23-7.41)	460 (942314)	0.49 (0.46-0.51)	1535 (942355)	1.63 (1.59-1.67)
	2	2871570	148 (101-178)	49	19	9	23	105430 (1074696)	3.30	98.10 (97.80-98.4)	5860 (1089699)	5.38 (5.31-5.45)	410 (1090201)	0.38 (0.36-0.39)	1310 (1090236)	1.20 (1.17-1.23)
	3	3318680	150 (109-180)	46	19	10	25	119680 (1274137)	3.06	93.93 (93.66-94.2)	5595 (1290631)	4.33 (4.28-4.39)	390 (1291108)	0.30 (0.29-0.32)	1265 (1291143)	0.98 (0.95-1.01)
	4	3274460	151 (113-181)	44	19	10	27	121740 (1272731)	3.07	95.65 (95.38-95.93)	5210 (1289336)	4.04 (3.98-4.10)	380 (1289775)	0.29 (0.28-0.31)	1195 (1289806)	0.92 (0.90-0.95)
	5 (least deprived)	3115025	152 (115-181)	41	19	11	29	120245 (1220876)	2.98	98.49 (98.21-98.77)	4395 (1237218)	3.55 (3.50-3.61)	310 (1237580)	0.25 (0.24-0.26)	1020 (1237606)	0.82 (0.80-0.85)
	Unknown	402555	139 (87-173)	43	20	11	26	15885 (142704)	3.64	111.31 (110.43-112.19)	635 (144971)	4.37 (4.20-4.55)	45 (145029)	0.30 (0.25-0.35)	120 (145032)	0.82 (0.75-0.90)
Region	London	891660	129 (83-170)	40	20	12	28	21970 (304836)	4.19	72.07 (71.59-72.56)	1100 (308097)	3.57 (3.46-3.68)	70 (308184)	0.23 (0.21-0.26)	220 (308190)	0.72 (0.67-0.77)
	East of England	3629745	149 (104-179)	44	19	10	27	113725 (1380061)	2.82	82.41 (82.16-82.65)	5105 (1394975)	3.66 (3.61-3.71)	410 (1395387)	0.29 (0.28-0.31)	1215 (1395416)	0.87 (0.84-0.89)

Clinical/ demographic group	Category	Fully vaccinated						Positive SARS-CoV-2 test			Hospitalised with COVID-19		COVID-19-related critical care admission		COVID-19 related death	
		Count*	Follow-up time, medium days (IQR)	Tests conducted (%)				Positivity (%)	Events* (PYs)	Incidence rate per 1000 person-years (95% CI)	Events* (PYs)	Incidence rate per 1000 person-years (95% CI)	Events* (PYs)	Incidence rate per 1000 person-years (95% CI)	Events* (PYs)	Incidence rate per 1000 person-years (95% CI)
				0	1	2	3+									
Region (continued)	East Midlands	2702030	150 (113-177)	48	19	10	24	108255 (1029678)	3.47	105.14 (104.82-105.46)	5480 (1044633)	5.24 (5.17-5.32)	420 (1045072)	0.4 (0.38-0.42)	1215 (1045111)	1.16 (1.13-1.20)
	North East	745775	150 (115-181)	49	20	10	22	37590 (289494)	4.54	129.85 (129.18-130.52)	2170 (294874)	7.36 (7.21-7.52)	140 (295080)	0.47 (0.43-0.51)	495 (295093)	1.68 (1.61-1.76)
	North West	1402645	150 (115-180)	49	19	9	23	63570 (541871)	3.55	117.32 (116.85-117.78)	3445 (551601)	6.24 (6.14-6.35)	265 (551937)	0.48 (0.45-0.51)	780 (551962)	1.42 (1.37-1.47)
	South East	1078015	151 (108-180)	43	19	10	27	34655 (417431)	2.65	83.02 (82.58-83.47)	1715 (422133)	4.06 (3.96-4.16)	110 (422262)	0.27 (0.24-0.29)	370 (422271)	0.88 (0.84-0.93)
	South West	2314235	152 (111-180)	45	19	10	27	83030 (896729)	2.56	92.59 (92.27-92.91)	3460 (907535)	3.81 (3.75-3.88)	185 (907805)	0.20 (0.19-0.22)	760 (907822)	0.83 (0.81-0.87)
	West Midlands	548600	149 (109-179)	52	18	9	22	21340 (208833)	3.69	102.19 (101.49-102.89)	1425 (212055)	6.73 (6.55-6.91)	105 (212201)	0.50 (0.45-0.55)	300 (212211)	1.40 (1.33-1.49)
	Yorkshire and The Humber	2179520	151 (111-181)	50	19	9	22	95360 (840635)	3.44	113.44 (113.07-113.81)	4670 (854660)	5.46 (5.38-5.54)	280 (855109)	0.33 (0.31-0.35)	1080 (855132)	1.26 (1.22-1.30)
	Unknown	9325	116 (67-164)	44	20	11	25	280 (2929)	3.85	96.27 (90.7-102.17)	15 (2969)	4.71 (3.61-6.16)	REDACTED	REDACTED	REDACTED	REDACTED
Asplenia		108350	170 (154-186)	40	19	11	30	4890 (48994)	2.62	99.79 (98.37-101.22)	280 (49718)	5.59 (5.27-5.94)	15 (49744)	0.32 (0.25-0.41)	55 (49746)	1.13 (0.98-1.29)
Asthma		2336650	151 (110-178)	42	20	11	27	105015 (893242)	3.43	117.57 (117.21-117.93)	5900 (908166)	6.5 (6.41-6.58)	425 (908684)	0.47 (0.44-0.49)	1070 (908724)	1.18 (1.14-1.21)
Blood pressure	Normal	3190995	137 (87-171)	41	20	11	28	136085 (1113098)	3.44	122.26 (121.93-122.59)	4975 (1132373)	4.4 (4.33-4.46)	260 (1132792)	0.23 (0.22-0.24)	1185 (1132816)	1.05 (1.02-1.08)
	Elevated/high	3428615	151 (122-179)	46	19	10	25	136200 (1347493)	3.08	101.08 (100.8-101.35)	6565 (1366494)	4.8 (4.74-4.86)	480 (1367064)	0.35 (0.34-0.37)	1420 (1367106)	1.04 (1.01-1.07)
	Unknown	8881945	152 (113-181)	48	19	10	24	307500 (3451907)	3.09	89.08 (88.92-89.24)	17040 (3494668)	4.88 (4.84-4.91)	1250 (3496150)	0.36 (0.35-0.37)	3830 (3496255)	1.10 (1.08-1.11)
Cancer (non-haematological)		891935	180 (160-194)	45	17	9	28	25625 (426015)	2.08	60.15 (59.77-60.52)	4065 (429286)	9.47 (9.32-9.62)	235 (429659)	0.54 (0.51-0.58)	1305 (429678)	3.04 (2.96-3.13)

Clinical/ demographic group	Category	Fully vaccinated						Positive SARS-CoV-2 test			Hospitalised with COVID-19		COVID-19-related critical care admission		COVID-19 related death	
		Count*	Follow-up time, medium days (IQR)	Tests conducted (%)				Positivity (%)	Events* (PYs)	Incidence rate per 1000 person-years (95% CI)	Events* (PYs)	Incidence rate per 1000 person-years (95% CI)	Events* (PYs)	Incidence rate per 1000 person-years (95% CI)	Events* (PYs)	Incidence rate per 1000 person-years (95% CI)
				0	1	2	3+									
Chronic kidney disease	Stage 3a	529000	187 (173-201)	53	16	8	23	12170 (268808)	2.18	45.27 (44.87-45.69)	3370 (270184)	12.48 (12.27-12.7)	190 (270500)	0.70 (0.65-0.75)	1080 (270514)	4.00 (3.88-4.12)
	Stage 3b	206450	192 (178-205)	52	14	7	27	4450 (107113)	2.16	41.56 (40.95-42.19)	2105 (107527)	19.59 (19.16-20.02)	95 (107716)	0.89 (0.80-0.99)	795 (107722)	7.39 (7.13-7.66)
	Stage 4	51210	191 (174-204)	47	13	8	33	1280 (25907)	2.19	49.41 (48.05-50.81)	780 (26000)	29.92 (28.87-31.01)	45 (26064)	1.80 (1.56-2.09)	335 (26067)	12.85 (12.17-13.57)
	Stage 5	5505	184 (164-200)	29	12	7	52	215 (2607)	1.75	82.47 (77.04-88.29)	130 (2627)	49.49 (45.33-54.02)	15 (2638)	5.31 (4.06-6.93)	45 (2638)	17.81 (15.4-20.61)
Diabetes		1476880	173 (157-191)	50	17	9	24	52540 (683694)	2.80	76.85 (76.51-77.18)	8645 (690614)	12.52 (12.39-12.65)	740 (691375)	1.07 (1.03-1.11)	2255 (691436)	3.26 (3.19-3.33)
Dialysis	Previous kidney transplant	2345	183 (172-198)	25	15	9	51	155 (1143)	2.54	134.75 (124.32-146.06)	80 (1159)	70.73 (63.34-78.99)	25 (1165)	19.74 (16.03-24.32)	30 (1167)	25.71 (21.42-30.86)
	No previous kidney transplant	10505	180 (165-199)	22	9	7	63	535 (4982)	1.38	107.19 (102.65-111.93)	230 (5048)	45.36 (42.46-48.46)	25 (5068)	5.33 (4.40-6.46)	65 (5070)	13.22 (11.7-14.93)
Heart disease		1745845	180 (163-194)	48	17	9	26	51745 (844423)	2.38	61.28 (61.01-61.55)	10590 (850928)	12.45 (12.33-12.57)	650 (851904)	0.76 (0.73-0.79)	3405 (851951)	4.00 (3.93-4.07)
Haematological malignancy		103625	180 (166-193)	42	17	9	31	3725 (49945)	2.91	74.56 (73.35-75.79)	1225 (50386)	24.35 (23.67-25.06)	140 (50481)	2.79 (2.57-3.04)	435 (50492)	8.62 (8.21-9.04)
Immunocompromised		508930	177 (159-191)	42	18	10	30	21695 (237016)	2.90	91.53 (90.91-92.15)	3950 (240002)	16.45 (16.19-16.71)	450 (240303)	1.88 (1.79-1.97)	1220 (240339)	5.08 (4.94-5.23)
Learning disability		102730	160 (146-177)	47	13	6	33	3230 (43892)	1.78	73.57 (72.28-74.87)	315 (44339)	7.06 (6.67-7.47)	30 (44367)	0.65 (0.54-0.79)	50 (44371)	1.13 (0.98-1.30)
Liver disease		348855	164 (144-182)	45	19	10	26	14020 (151587)	2.96	92.5 (91.72-93.29)	1620 (153529)	10.55 (10.29-10.81)	130 (153667)	0.85 (0.78-0.93)	360 (153678)	2.36 (2.23-2.48)
Neurological disease		874635	176 (156-193)	45	16	8	30	28410 (407866)	2.29	69.66 (69.24-70.07)	5075 (411606)	12.33 (12.16-12.51)	245 (412073)	0.59 (0.55-0.63)	1685 (412095)	4.09 (3.99-4.19)
Respiratory disease		581930	180 (166-193)	50	17	9	25	16785 (282315)	2.53	59.45 (58.99-59.91)	4900 (284321)	17.24 (16.99-17.49)	305 (284751)	1.07 (1.01-1.14)	1515 (284773)	5.32 (5.19-5.46)

Clinical/ demographic group	Category	Fully vaccinated						Positive SARS-CoV-2 test			Hospitalised with COVID-19		COVID-19-related critical care admission		COVID-19 related death	
		Count*	Follow-up time, medium days (IQR)	Tests conducted (%)				Positivity (%)	Events* (PYs)	Incidence rate per 1000 person-years (95% CI)	Events* (PYs)	Incidence rate per 1000 person-years (95% CI)	Events* (PYs)	Incidence rate per 1000 person-years (95% CI)	Events* (PYs)	Incidence rate per 1000 person-years (95% CI)
				0	1	2	3+									
Severe mental illness		181865	159 (141-177)	51	15	8	26	5135 (76184)	2.44	67.42 (66.48-68.36)	650 (76905)	8.44 (8.11-8.78)	50 (76961)	0.64 (0.55-0.73)	155 (76967)	2.01 (1.86-2.18)
Transplant	Kidney (with previous dialysis)	5745	179 (172-191)	35	18	11	37	395 (2745)	4.83	143.51 (136.46-150.93)	210 (2787)	76.08 (71.03-81.49)	50 (2800)	17.86 (15.5-20.57)	55 (2804)	18.9 (16.47-21.68)
	Kidney (without previous dialysis)	4515	179 (171-191)	36	19	11	35	280 (2160)	4.65	130.07 (122.54-138.06)	130 (2192)	58.4 (53.46-63.8)	25 (2202)	10.45 (8.48-12.87)	30 (2203)	14.52 (12.17-17.33)
	Other organ	6780	178 (170-188)	35	18	11	36	405 (3202)	3.91	127.12 (120.97-133.58)	165 (3247)	50.2 (46.42-54.29)	40 (3259)	11.66 (9.91-13.71)	55 (3262)	16.86 (14.73-19.29)
Time since being fully vaccinated	0-4 weeks	331535	16 (8-23)	63	19	7	11	1100 (13569)	5.91	81.14 (78.73-83.62)	170 (13580)	12.37 (11.45-13.36)	REDACTED	REDACTED	80 (13582)	5.74 (5.13-6.43)
	4 - 8 weeks	886870	44 (37-51)	59	20	9	12	5135 (105598)	7.09	48.65 (47.97-49.33)	335 (105819)	3.15 (2.98-3.32)	20 (105825)	0.18 (0.14-0.23)	125 (105826)	1.17 (1.07-1.28)
	8 - 12 weeks	1430095	69 (62-76)	50	22	11	17	19585 (270294)	6.32	72.45 (71.94-72.97)	610 (271783)	2.25 (2.16-2.34)	55 (271801)	0.20 (0.18-0.23)	270 (271804)	0.99 (0.93-1.05)
	12+ weeks	12853055	159 (135-184)	44	19	10	27	553960 (5523038)	2.92	100.3 (100.16-100.43)	27470 (5602353)	4.90 (4.87-4.93)	1910 (5604799)	0.34 (0.33-0.35)	5965 (5604967)	1.06 (1.05-1.08)
Time between vaccinations	6 weeks or less	499185	185 (111-283)	39	18	11	31	17760 (253744)	2.43	69.99 (69.47-70.52)	1630 (256550)	6.35 (6.20-6.51)	85 (256777)	0.34 (0.03-0.38)	450 (256788)	1.75 (1.67-1.83)
	6-12 weeks	13565570	150 (108-179)	46	19	10	25	519470 (5161343)	3.20	100.65 (100.51-100.79)	23120 (5233675)	4.42 (4.39-4.45)	1635 (5235630)	0.31 (0.03-0.32)	5005 (5235771)	0.96 (0.94-0.97)
	12 weeks or more	1436800	143 (100-170)	53	18	8	21	42550 (497411)	3.19	85.54 (85.13-85.96)	3830 (503309)	7.61 (7.49-7.73)	270 (503599)	0.53 (0.05-0.57)	980 (503619)	1.95 (1.89-2.01)
Prior COVID-19	Between 1st and 2nd vaccination	191320	65 (42-166)	48	19	9	24	1720 (49491)	1.14	34.75 (33.93-35.6)	160 (49786)	3.25 (3.01-3.52)	REDACTED	REDACTED	55 (49815)	1.08 (0.95-1.24)
	Before 1st vaccination	926960	138 (88-170)	42	19	10	29	8895 (326947)	1.19	27.21 (26.92-27.5)	755 (328374)	2.31 (2.22-2.39)	30 (328454)	0.09 (0.08-0.11)	200 (328458)	0.61 (0.57-0.66)

Figure 1 - Inclusion/exclusion flowchart




Figure 2 Kaplan-Meier plot for positive SARS-CoV-2 test over time, by priority group

