


A

MAP2 p.Leu867fs | JLF V2, JLF V1 | 24 aa


disclosed minimal epitope RVVPFTKAL at 9-17

CYTOSOLIC CROSS-PRESENTATION - conditional after SLP uptake/export


four model-native tracks



MHC-I: 7 shown / 39 eligible (top-10 cap)



MHC-II: 5 shown / 11 eligible (top-10 cap)



ENDOLYSOSOMAL / CLASS-II PROCESSING - primary SLP uptake route

NetCleave-II is not a named-cathepsin assay




EXTRACELLULAR BEFORE/DURING SLP UPTAKE - exposure-dependent evidence



SCORES Separate native 0-1 tracks; joined assessed bonds are not smoothed. Dashed = 0.5 display threshold; gaps = unassessed.

RED All four cytosolic tracks assessed the bond and ≥ 3 reached 0.5. Four filled/open beads show model concurrence, never probability.


MHC Up to 10/class: intended overlap, then allele diversity, then native rank. Red ticks inside bars are pre-binding cut evidence; full results stay in CSV.



C

PIP5K1A p.Gly474fs | JLF V3 | 20 aa

disclosed minimal epitope AAPVATPAL at 7-15



SCORES Separate native 0-1 tracks; joined assessed bonds are not smoothed. Dashed = 0.5 display threshold; gaps = unassessed.

RED All four cytosolic tracks assessed the bond and ≥ 3 reached 0.5. Four filled/open beads show model concurrence, never probability.

MHC Up to 10/class: intended overlap, then allele diversity, then native rank. Red ticks inside bars are pre-binding cut evidence; full results stay in CSV.

