


NME1 p.Gly24Arg | CeGaT | 13 aa

minimal epitope/window not disclosed for this SLP

CYTOSOLIC CROSS-PRESENTATION - conditional after SLP uptake/export

four model-native tracks



C01:02 1.1%

P P I S S C D T R T M A N

MHC-II: none <= 5% rank

1

5

10

13

ENDOLYSOSOMAL / CLASS-II PROCESSING - primary SLP uptake route

NetCleave-II is not a named-cathepsin assay

NetCleave II native 0-1



EXTRACELLULAR ENZYME EVIDENCE - requires enzyme exposure

P P I S S C D T R T M A N

1

5


10

13


DPP4 / qPISA  
N-terminal 2-residue removal

NOT SCORED / OUTSIDE MODEL SCOPE  
Dataset EV2 has missing coefficients for P1=P: P2:P


MME / neprilysin  
Internal hydrophobic preference




FAP - internal  
Gly-Pro|non-Pro recognition




FAP - N-terminal  
Remove 2 residues: X-Pro|non-Pro



ANPEP / CD13  
exposed N-terminus only



ENPEP  
exposed N-terminus only



Triangle = motif match | hollow circle = no motif match | NA = not assessed. Enzyme presence and exposure are not measured.

DPP4: predicted peptide-signal loss in the source 4 h assay. Native scores remain in CSV. Motifs indicate sequence matches only.

SCORES Separate native 0-1 tracks; joined assessed bonds are not smoothed. Dashed = 0.5 display threshold; gaps = unassessed.

RED All four cytosolic tracks assessed the bond and >=3 reached 0.5. Four filled/open beads show model concurrence, never probability.

MHC Up to 10/class: intended overlap, then allele diversity, then native rank. Red ticks inside bars are pre-binding cut evidence; full results stay in CSV.