

EXOC4 p.Ser34Ile | JLF V3, JLF V2 | 23 aa

minimal epitope/window not disclosed for this SLP

CYTOSOLIC CROSS-PRESENTATION - conditional after SLP uptake/export

four model-native tracks



NetChop Cterm (ligand) native 0-1

Pepsickle human (in vivo) native 0-1

NetCleave I native 0-1

NetChop 20S (in vitro) native 0-1

MHC-I: 3 shown / 4 eligible (top-10 cap)



ENDOLYSOSOMAL / CLASS-II PROCESSING - primary SLP uptake route

NetCleave-II is not a named-cathepsin assay

NetCleave II native 0-1

EXTRACELLULAR ENZYME EVIDENCE - requires enzyme exposure



DPP4 / qPISA
N-terminal 2-residue removal


MME / neprilysin
Internal hydrophobic preference

FAP - internal
Gly-Pro|non-Pro recognition

FAP - N-terminal
Remove 2 residues: X-Pro|non-Pro

ANPEP / CD13
exposed N-terminus only

ENPEP
exposed N-terminus only



Triangle = motif match | hollow circle = no motif match | NA = not assessed. Enzyme presence and exposure are not measured.

DPP4: predicted peptide-signal loss in the source 4 h assay. Native scores remain in CSV. Motifs indicate sequence matches only.

SCORES Separate native 0-1 tracks; joined assessed bonds are not smoothed. Dashed = 0.5 display threshold; gaps = unassessed.

RED All four cytosolic tracks assessed the bond and >=3 reached 0.5. Four filled/open beads show model concurrence, never probability.

MHC Up to 10/class: intended overlap, then allele diversity, then native rank. Red ticks inside bars are pre-binding cut evidence; full results stay in CSV.