

MAP2 p.Leu867fs | JLF V3 | 23 aa

disclosed minimal epitope RVVPFTKAL at 9-17

CYTOSOLIC CROSS-PRESENTATION - conditional after SLP uptake/export

four model-native tracks

NetChop Cterm (ligand) native 0-1

Pepsickle human (in vivo) native 0-1

NetCleave I native 0-1

NetChop 20S (in vitro) native 0-1

0.5 display


0.5 display

0.5 display

0.5 display

MHC-I: 6 shown / 35 eligible (top-10 cap)

MHC-II: 5 shown / 11 eligible (top-10 cap)



K T V R I Y Q G R V V P F T K A L M I K F E E

ENDOLYSOSOMAL / CLASS-II PROCESSING - primary SLP uptake route

NetCleave-II is not a named-cathepsin assay

NetCleave II native 0-1

0.5 display

EXTRACELLULAR ENZYME EVIDENCE - requires enzyme exposure

K T V R I Y Q G R V V P F T K A L M I K F E E

1 5 10 15 20 23

DPP4 / qPISA

N-terminal 2-residue removal

29% predicted loss

MME / neprilysin

Internal hydrophobic preference

FAP - internal

Gly-Pro|non-Pro recognition

FAP - N-terminal

Remove 2 residues: X-Pro|non-Pro

ANPEP / CD13

exposed N-terminus only

ENPEP

exposed N-terminus only

Triangle = motif match | hollow circle = no motif match | NA = not assessed. Enzyme presence and exposure are not measured.

DPP4: predicted peptide-signal loss in the source 4 h assay. Native scores remain in CSV. Motifs indicate sequence matches only.

SCORES Separate native 0-1 tracks; joined assessed bonds are not smoothed. Dashed = 0.5 display threshold; gaps = unassessed.

RED All four cytosolic tracks assessed the bond and >=3 reached 0.5. Four filled/open beads show model concurrence, never probability.

MHC Up to 10/class: intended overlap, then allele diversity, then native rank. Red ticks inside bars are pre-binding cut evidence; full results stay in CSV.