

minimal epitope/window not disclosed for this SLP

NetCleave I 0-1

NetChop Cterm 0-1

Pepsickle all-mammal 0-1

Pepsickle human 0-1

NetChop 20S 0-1

MHC-I candidate ligands

MHC-II candidate ligands

NetCleave II 0-1

DPP4 native

ERAP1 / ERAMER

plasma / extracellular motifs

cytosol / ER motifs

S F M L R A V S F | F V K D A V L | Y | S G A K | K | K

1	5	10	DR03:01 2.4%	15	20	23
---	---	----	--------------	----	----	----

0.012

MME

NPEPPS

▼
MME

▼
TPP2

MME

MME

MME

▼
ACE

CPB2+CPN1

Reading the map. Each profile joins exact scores at adjacent assessed bonds; it is not smoothed, averaged, or calibrated, and gaps are unassessed. Dots at or beyond the dashed 0.5 line are display candidates. Red slashes between residues require ≥ 3 of the five class-I-processing models. Pale blue/purple sequence tint marks coverage by a displayed MHC window; outlined windows identify the individual candidate MHC ligands at $\leq 2\%/\leq 5\%$ rank (top-ranked non-overlapping spans shown; all rows are in the CSV); red ticks inside them are internal candidate cuts before binding. These counts are not probabilities, and ligand bars do not model binding occupancy, timing, or post-binding protection.