

ZNF436 p.Ser207Cys | CeGaT | 16 aa

disclosed minimal epitope KSFGRSCHL at 2-10

INTRACELLULAR / CLASS-I PROCESSING - four complementary score tracks

higher native score upward

NetChop Cterm (ligand) native 0-1

0.5 display

Pepsickle human (in vivo) native 0-1

0.5 display

NetCleave I native 0-1

0.5 display

NetChop 20S (in vitro) native 0-1

0.5 display

ERAP1 / ERAMER

cytosol / ER motifs

-0.061

TPP2

+7 more in CSV

MHC-I candidate ligands

C01:02 0.62%

B27:05 0.28%

B08:01 0.16%

G

K

S

F

G

R

S

C

H

L

I

Q

H

Q

T

I

MHC-II: none <= 5% rank

1

5

10

15

16

ENDOLYSOSOMAL / CLASS-II PROCESSING - C-terminal score track

higher native score downward

NetCleave II native 0-1

0.5 display

SERUM / EXTRACELLULAR - native scores and recognition motifs

DPP4 native (intact SLP)

serum / extracellular motifs

MME

MME

MME

ACE

MME

SCORES Native 0-1 bond scores; lines join adjacent assessed bonds only (no smoothing). Dashed line = 0.5 display threshold; gaps = unassessed.

RED BOND MARKS All four intracellular tracks assess the bond; >=3 reach 0.5. Four beads show support (filled = hit, open = miss), not probability.

MHC WINDOWS <=2%/<=5% rank; display favors intended overlap, then allele diversity, then rank. Internal red ticks are pre-binding cut evidence; all predictions remain in CSV.