

GLIS3 p.Ser775fs | JLF V3 | 80 aa - segment 1/3, bonds 1-27

minimal epitope/window not disclosed for this SLP

INTRACELLULAR / CLASS-I PROCESSING - four complementary score tracks

higher native score upward

NetChop Cterm (ligand) native 0-1

0.5 display

Pepsickle human (in vivo) native 0-1

0.5 display

NetCleave I native 0-1

0.5 display

NetChop 20S (in vitro) native 0-1

0.5 display

ERAP1 / ERAMER

cytosol / ER motifs

XPNPEP1

▲

MHC-I candidate ligands

C07:01 0.24%

A01:01 1%

B27:05 0.97%

C01:02 1.4%

C01:02 0.69%

B08:01 1.2%

+12 more in CSV

Q | P | P | E | S | S | S | S | F | F | N | T | A | K | N | T | A | S | L | Y | P | A | A | I | R | F | T | P

MHC-II candidate ligands

DR08:01 3.7%

DR08:01 3.5%

DR08:01 4.2%

ENDOLYSOSOMAL / CLASS-II PROCESSING - C-terminal score track

higher native score downward

NetCleave II native 0-1

0.5 display

SERUM / EXTRACELLULAR - native scores and recognition motifs

DPP4 native (intact SLP)

0.11

serum / extracellular motifs

XPNPEP2

▼

SCORES Native 0-1 bond scores; lines join adjacent assessed bonds only (no smoothing). Dashed line = 0.5 display threshold; gaps = unassessed.

RED BOND MARKS All four intracellular tracks assess the bond; >=3 reach 0.5. Four beads show support (filled = hit, open = miss), not probability.

MHC WINDOWS <=2%/<=5% rank; display favors intended overlap, then allele diversity, then rank. Internal red ticks are pre-binding cut evidence; all predictions remain in CSV.