

CUL9 p.Asp403Tyr | JLF V2 | 23 aa

minimal epitope/window not disclosed for this SLP

INTRACELLULAR / CLASS-I PROCESSING - four complementary score tracks

higher native score upward

NetChop Cterm (ligand) native 0-1

0.5 display

Pepsickle human (in vivo) native 0-1

0.5 display

NetCleave I native 0-1

0.5 display

NetChop 20S (in vitro) native 0-1

0.5 display

ERAP1 / ERAMER

cytosol / ER motifs

ERAP2+NPEPP5

TPP2

MHC-I candidate ligands

+3 more in CSV

C01:02 0.12%

B27:05 1%

B27:05 0.68%

R | M | L | D | Y | Y | E | E | I | S | A | G | D | E | G | E | F | R | Q | S | K | K | K

MHC-II candidate ligands

DQ04:01/04:02 4.3%

DR08:01 3.4%

DQ05:01/02:01 2.3%

ENDOLYSOSOMAL / CLASS-II PROCESSING - C-terminal score track

higher native score downward

NetCleave II native 0-1

0.5 display

SERUM / EXTRACELLULAR - native scores and recognition motifs

DPP4 native (intact SLP)

serum / extracellular motifs

MME

MME

MME

MME

MME

ACE

CPB2+CPN1

SCORES Native 0-1 bond scores; lines join adjacent assessed bonds only (no smoothing). Dashed line = 0.5 display threshold; gaps = unassessed.

RED BOND MARKS All four intracellular tracks assess the bond; >=3 reach 0.5. Four beads show support (filled = hit, open = miss), not probability.

MHC WINDOWS <=2%/<=5% rank; display favors intended overlap, then allele diversity, then rank. Internal red ticks are pre-binding cut evidence; all predictions remain in CSV.