

SLC25A12 p.Ile347Thr | JLF V2, JLF V1 | 26 aa

disclosed minimal epitope ATAVYPTDL at 9-17

INTRACELLULAR / CLASS-I PROCESSING - four complementary score tracks

higher native score upward

NetChop Cterm (ligand) native 0-1

0.5 display

Pepsickle human (in vivo) native 0-1

0.5 display

NetCleave I native 0-1

0.5 display

NetChop 20S (in vitro) native 0-1

0.5 display

ERAP1 / ERAMER

cytosol / ER motifs

TPP2

PREP

+14 more in CSV

MHC-I candidate ligands

C01:02 1.3%

C01:02 0.93%

B08:01 0.67%

G | S | V | A | G | A | V | G | A | T | A | V | Y | P | T | D | L | V | K | T | R | M | Q | N | Q | R

MHC-II candidate ligands

DQ05:01/02:01 4.3%

DQ05:01/02:01 3.9%

DQ04:01/04:02 4.7%

ENDOLYSOSOMAL / CLASS-II PROCESSING - C-terminal score track

higher native score downward

NetCleave II native 0-1

0.5 display

SERUM / EXTRACELLULAR - native scores and recognition motifs

DPP4 native (intact SLP)

-0.58

serum / extracellular motifs

MME

MME

ACE

CPB2+CPN1

SCORES Native 0-1 bond scores; lines join adjacent assessed bonds only (no smoothing). Dashed line = 0.5 display threshold; gaps = unassessed.

RED BOND MARKS All four intracellular tracks assess the bond; >=3 reach 0.5. Four beads show support (filled = hit, open = miss), not probability.

MHC WINDOWS <=2%/<=5% rank; display favors intended overlap, then allele diversity, then rank. Internal red ticks are pre-binding cut evidence; all predictions remain in CSV.