

NR2F2 p.Gly65Val | JLF V2 | 26 aa

minimal epitope/window not disclosed for this SLP

INTRACELLULAR / CLASS-I PROCESSING - four complementary score tracks

higher native score upward

NetChop Cterm (ligand) native 0-1

0.5 display

Pepsickle human (in vivo) native 0-1

0.5 display

NetCleave I native 0-1

0.5 display

NetChop 20S (in vitro) native 0-1

0.5 display

ERAP1 / ERAMER

cytosol / ER motifs

NPEPPS

TPP2

PREP

PREP

+6 more in CSV

MHC-I candidate ligands

B27:05 1.5%

B27:05 1.3%

B27:05 1.2%

B27:05 1.8%

A | Q | T | A | A | G | G | Q | G | G | P | G | V | P | G | S | D | K | Q | Q | Q | Q | Q | H | I | E

MHC-II: none <= 5% rank

1 5 10 15 20 25 26

ENDOLYSOSOMAL / CLASS-II PROCESSING - C-terminal score track

higher native score downward

NetCleave II native 0-1

0.5 display

SERUM / EXTRACELLULAR - native scores and recognition motifs

DPP4 native (intact SLP)

0.024

serum / extracellular motifs

ANPEP

FAP

MME

SCORES Native 0-1 bond scores; lines join adjacent assessed bonds only (no smoothing). Dashed line = 0.5 display threshold; gaps = unassessed.

RED BOND MARKS All four intracellular tracks assess the bond; >=3 reach 0.5. Four beads show support (filled = hit, open = miss), not probability.

MHC WINDOWS <=2%/<=5% rank; display favors intended overlap, then allele diversity, then rank. Internal red ticks are pre-binding cut evidence; all predictions remain in CSV.