



A

MAP2 p.Leu867fs | JLF V2, JLF V1 | 24 aa

disclosed minimal epitope RVVPFTKAL at 9-17

INTRACELLULAR / CLASS-I PROCESSING - four complementary score tracks

higher native score upward



ENDOLYSOSOMAL / CLASS-II PROCESSING - C-terminal score track

higher native score downward



SERUM / EXTRACELLULAR - native scores and recognition motifs



SCORES Native 0-1 bond scores; lines join adjacent assessed bonds only (no smoothing). Dashed line = 0.5 display threshold; gaps = unassessed.

RED BOND MARKS All four intracellular tracks assess the bond; ≥ 3 reach 0.5. Four beads show support (filled = hit, open = miss), not probability.

MHC WINDOWS $\leq 2\% / \leq 5\%$ rank; display favors intended overlap, then allele diversity, then rank. Internal red ticks are pre-binding cut evidence; all predictions remain in CSV.


B

SMC5 p.Arg371Lys | JLF V3 | 21 aa

disclosed minimal epitope RQKRIGNTR at 2-10

INTRACELLULAR / CLASS-I PROCESSING - four complementary score tracks

higher native score upward



ENDOLYSOSOMAL / CLASS-II PROCESSING - C-terminal score track

higher native score downward



SERUM / EXTRACELLULAR - native scores and recognition motifs



SCORES Native 0-1 bond scores; lines join adjacent assessed bonds only (no smoothing). Dashed line = 0.5 display threshold; gaps = unassessed.

RED BOND MARKS All four intracellular tracks assess the bond; ≥ 3 reach 0.5. Four beads show support (filled = hit, open = miss), not probability.

MHC WINDOWS $\leq 2\%$ / $\leq 5\%$ rank; display favors intended overlap, then allele diversity, then rank. Internal red ticks are pre-binding cut evidence; all predictions remain in CSV.

C

PIP5K1A p.Gly474fs | JLF V3 | 20 aa

disclosed minimal epitope AAPVATPAL at 7-15

INTRACELLULAR / CLASS-I PROCESSING - four complementary score tracks

higher native score upward

NetChop Cterm (ligand) native 0-1

0.5 display

Pepsickle human (in vivo) native 0-1

0.5 display

NetCleave I native 0-1

0.5 display

NetChop 20S (in vitro) native 0-1

0.5 display

ERAP1 / ERAMER
cytosol / ER motifs

NPEPP5

TPP2

PREP

PREP

+23 more in CSV

MHC-I candidate ligands

B08:01 0.36%

C01:02 0.036%

C07:01 0.2%

S | S | F | S | R | R | A | A | P | V | A | T | P | A | L | L | T | S | H | R

MHC-II: none <= 5% rank

1

5

10

15

20

ENDOLYSOSOMAL / CLASS-II PROCESSING - C-terminal score track

higher native score downward

NetCleave II native 0-1

0.5 display

SERUM / EXTRACELLULAR - native scores and recognition motifs

DPP4 native (intact SLP)

0.62

serum / extracellular motifs

MME

MME

MME

ACE

CPB2+CPN1

SCORES Native 0-1 bond scores; lines join adjacent assessed bonds only (no smoothing). Dashed line = 0.5 display threshold; gaps = unassessed.

RED BOND MARKS All four intracellular tracks assess the bond; >=3 reach 0.5. Four beads show support (filled = hit, open = miss), not probability.

MHC WINDOWS <=2%/<=5% rank; display favors intended overlap, then allele diversity, then rank. Internal red ticks are pre-binding cut evidence; all predictions remain in CSV.



D

DYNC1H1 p.Val314Ile | JLF V3, JLF V2 | 24 aa

disclosed minimal epitope KRFHATISF at 2-10

INTRACELLULAR / CLASS-I PROCESSING - four complementary score tracks

higher native score upward



ENDOLYSOSOMAL / CLASS-II PROCESSING - C-terminal score track

higher native score downward



SERUM / EXTRACELLULAR - native scores and recognition motifs



SCORES Native 0-1 bond scores; lines join adjacent assessed bonds only (no smoothing). Dashed line = 0.5 display threshold; gaps = unassessed.

RED BOND MARKS All four intracellular tracks assess the bond; ≥ 3 reach 0.5. Four beads show support (filled = hit, open = miss), not probability.

MHC WINDOWS $\leq 2\%/\leq 5\%$ rank; display favors intended overlap, then allele diversity, then rank. Internal red ticks are pre-binding cut evidence; all predictions remain in CSV.