


GLIS3 p.Ser775fs | JLF V3 | 80 aa- segment 2/3, bonds 28-54


minimal epitope/window not disclosed for this SLP

CYTOSOLIC CROSS-PRESENTATION - conditional after SLP uptake/export


four model-native tracks



MHC-I: 8 shown / 14 eligible (top-10 cap)




MHC-II: 5 shown / 19 eligible (top-10 cap)



ENDOLYSOSOMAL / CLASS-II PROCESSING - primary SLP uptake route

NetCleave-II is not a named-cathepsin assay



EXTRACELLULAR ENZYME EVIDENCE - requires enzyme exposure




DPP4 / qPISA
N-terminal 2-residue removal

SITE OUTSIDE THIS SEGMENT | full-SLP score +0.113 log2 depletion
Intact SLP bond 2 is outside this segment

MME / neprilysin
Internal hydrophobic preference

NOT ASSESSED / OUTSIDE RULE SCOPE
Rule is limited to peptides of at most 30 residues

FAP - internal
Gly-Pro|non-Pro recognition



FAP - N-terminal
Remove 2 residues: X-Pro|non-Pro

SITE OUTSIDE THIS SEGMENT
Intact N-terminal bond 2 is outside this segment

ANPEP / CD13
exposed N-terminus only

SITE OUTSIDE THIS SEGMENT
Intact N-terminal bond 1 is outside this segment

ENPEP
exposed N-terminus only

SITE OUTSIDE THIS SEGMENT
Intact N-terminal bond 1 is outside this segment

M = motif match | hollow circle = assessed, no motif match | NA = not assessed. Enzyme presence and exposure are not measured.
DPP4: predicted log2(buffer control / treated substrate), source assay 4 h. Motifs have no numeric score; neither view gives a cleavage rate.

SCORES Separate native 0-1 tracks; joined assessed bonds are not smoothed. Dashed = 0.5 display threshold; gaps = unassessed.

RED All four cytosolic tracks assessed the bond and >=3 reached 0.5. Four filled/open beads show model concurrence, never probability.

MHC Up to 10/class: intended overlap, then allele diversity, then native rank. Red ticks inside bars are pre-binding cut evidence; full results stay in CSV.