


CUL9 p.Asp403Tyr | JLF V2 | 23 aa


minimal epitope/window not disclosed for this SLP

CYTOSOLIC CROSS-PRESENTATION - conditional after SLP uptake/export


four model-native tracks



MHC-I: 4 shown / 7 eligible (top-10 cap)



MHC-II: 5 shown / 13 eligible (top-10 cap)



ENDOLYSOSOMAL / CLASS-II PROCESSING - primary SLP uptake route

NetCleave-II is not a named-cathepsin assay



EXTRACELLULAR ENZYME EVIDENCE - requires enzyme exposure



DPP4 / qPISA

N-terminal 2-residue removal

NOT SCORED / OUTSIDE MODEL SCOPE

Dataset EV2 has missing coefficients for P1=M: P2:R

5 motif matches / 22 assessed bonds - partial recognition rule



MME / neprilysin

Internal hydrophobic preference

0 motif matches / 22 assessed bonds - partial recognition rule



FAP - internal

Gly-Pro|non-Pro recognition

0 motif matches / 1 assessed bond - partial recognition rule

FAP - N-terminal

Remove 2 residues: X-Pro|non-Pro

0 motif matches / 1 assessed bond - partial recognition rule

ANPEP / CD13

exposed N-terminus only

0 motif matches / 1 assessed bond - partial recognition rule

ENPEP

exposed N-terminus only

M = motif match | hollow circle = assessed, no motif match | NA = not assessed. Enzyme presence and exposure are not measured.

DPP4: predicted log₂(buffer control / treated substrate), source assay 4 h. Motifs have no numeric score; neither view gives a cleavage rate

SCORES Separate native 0-1 tracks; joined assessed bonds are not smoothed. Dashed = 0.5 display threshold; gaps = unassessed

RED All four cytosolic tracks assessed the bond and ≥ 3 reached 0.5. Four filled/open beads show model concurrence, never probability.

MHC Up to 10/class: intended overlap, then allele diversity, then native rank. Red ticks inside bars are pre-binding cut evidence; full results stay in CSV.