

SLC25A12 p.Ile347Thr | CeGaT | 17 aa

disclosed minimal epitope ATAVYPTDL at 3-11

CYTOSOLIC CROSS-PRESENTATION - conditional after SLP uptake/export

four model-native tracks

NetChop Cterm (ligand) native 0-1

0.5 display

Pepsickle human (in vivo) native 0-1

0.5 display

NetCleave I native 0-1


0.5 display

NetChop 20S (in vitro) native 0-1

0.5 display

MHC-I: 5 shown / 18 eligible (top-10 cap)

MHC-II: none <= 5% rank




ENDOLYSOSOMAL / CLASS-II PROCESSING - primary SLP uptake route

NetCleave-II is not a named-cathepsin assay

NetCleave II native 0-1

0.5 display

EXTRACELLULAR ENZYME EVIDENCE - requires enzyme exposure



M = motif match | hollow circle = assessed, no motif match | NA = not assessed. Enzyme presence and exposure are not measured.

DPP4: predicted log2(buffer control / treated substrate), source assay 4 h. Motifs have no numeric score; neither view gives a cleavage rate.

SCORES Separate native 0-1 tracks; joined assessed bonds are not smoothed. Dashed = 0.5 display threshold; gaps = unassessed.

RED All four cytosolic tracks assessed the bond and >=3 reached 0.5. Four filled/open beads show model concurrence, never probability.

MHC Up to 10/class: intended overlap, then allele diversity, then native rank. Red ticks inside bars are pre-binding cut evidence; full results stay in CSV.