


CENPL p.Pro26His |JLF V3 | 23 aa


minimal epitope/window not disclosed for this SLP

CYTOSOLIC CROSS-PRESENTATION - conditional after SLP uptake/export


four model-native tracks



MHC-I: 9 shown / 20 eligible (top-10 cap)



MHC-II: 4 shown / 5 eligible (top-10 cap)



ENDOLYSOSOMAL / CLASS-II PROCESSING - primary SLP uptake route

NetCleave-II is not a named-cathepsin assay



EXTRACELLULAR ENZYME EVIDENCE - requires enzyme exposure



DPP4 / qPISA

N-terminal 2-residue removal


NOT SCORED / OUTSIDE MODEL SCOPE

Dataset EV2 has missing coefficients for P1=S: P1':R

5 motif matches / 22 assessed bonds - partial recognition rule

MME / neprilysin

Internal hydrophobic preference



0 motif matches / 22 assessed bonds - partial recognition rule

FAP - internal


Gly-Pro|non-Pro recognition



0 motif matches / 1 assessed bond - partial recognition rule

FAP - N-terminal

Remove 2 residues: X-Pro|non-Pro



0 motif matches / 1 assessed bond - partial recognition rule


ANPEP / CD13

exposed N-terminus only



ENPEP

exposed N-terminus only



M = motif match | hollow circle = assessed, no motif match | NA = not assessed. Enzyme presence and exposure are not measured.

DPP4: predicted log2(buffer control / treated substrate), source assay 4 h. Motifs have no numeric score; neither view gives a cleavage rate.

SCORES Separate native 0-1 tracks; joined assessed bonds are not smoothed. Dashed = 0.5 display threshold; gaps = unassessed.

RED All four cytosolic tracks assessed the bond and >=3 reached 0.5. Four filled/open beads show model concurrence, never probability.

MHC Up to 10/class: intended overlap, then allele diversity, then native rank. Red ticks inside bars are pre-binding cut evidence; full results stay in CSV.