


PTH1R p.Ile237Phe | JLF V3 | 23 aa


minimal epitope/window not disclosed for this SLP

CYTOSOLIC CROSS-PRESENTATION - conditional after SLP uptake/export

four model-native tracks




MHC-I: 7 shown / 23 eligible (top-10 cap)



1 5 15 20 23

MHC-II: 5 shown / 15 eligible (top-10 cap)




ENDOLYSOSOMAL / CLASS-II PROCESSING - primary SLP uptake route

NetCleave-II is not a named-cathepsin assay



EXTRACELLULAR ENZYME EVIDENCE - requires enzyme exposure



DPP4 / qPISA
N-terminal 2-residue removal




MME / neprilysin
Internal hydrophobic preference




FAP - internal
Gly-Pro|non-Pro recognition




FAP - N-terminal
Remove 2 residues: X-Pro|non-Pro



ANPEP / CD13
exposed N-terminus only



ENPEP
exposed N-terminus only



M = motif match | hollow circle = assessed, no motif match | NA = not assessed. Enzyme presence and exposure are not measured.

DPP4: predicted log2(buffer control / treated substrate), source assay 4 h. Motifs have no numeric score; neither view gives a cleavage rate.

SCORES Separate native 0-1 tracks; joined assessed bonds are not smoothed. Dashed = 0.5 display threshold; gaps = unassessed.

RED All four cytosolic tracks assessed the bond and >=3 reached 0.5. Four filled/open beads show model concurrence, never probability.

MHC Up to 10/class: intended overlap, then allele diversity, then native rank. Red ticks inside bars are pre-binding cut evidence; full results stay in CSV.