

ADGRF5 p.Asn1279Ser | JLF V2 | 22 aa

minimal epitope/window not disclosed for this SLP

INTRACELLULAR / CLASS-I PROCESSING - four complementary score tracks

higher native score upward

NetChop Cterm (ligand) native 0-1

0.5 display

Pepsickle human (in vivo) native 0-1

0.5 display

NetCleave I native 0-1

0.5 display

NetChop 20S (in vitro) native 0-1

0.5 display

ERAP1 / ERAMER

cytosol / ER motifs

NPEPP5

TPP2

+4 more in CSV

MHC-I candidate ligands

B08:01 0.36%

B08:01 0.35%

B27:05 1.4%

B27:05 1.9%

B27:05 0.38%

V | Q | E | A | L | L | S | K | F | S | L | S | R | W | S | S | Q | H | S | K | S | T

MHC-II candidate ligands

DQ04:01/04:02 1.1%

DQ04:01/04:02 0.79%

DQ04:01/04:02 0.98%

+11 more in CSV

SERUM / EXTRACELLULAR - native scores and recognition motifs

higher native score downward

NetCleave II native 0-1

0.5 display

DPP4 native (intact SLP)

0.019

serum / extracellular motifs

MME

MME

MME

MME

ACE

SCORES Native 0-1 bond scores; lines join adjacent assessed bonds only (no smoothing). Dashed line = 0.5 display threshold; gaps = unassessed.

RED BOND MARKS All four intracellular tracks assess the bond; >=3 reach 0.5. Four beads show support (filled = hit, open = miss), not probability.

MHC WINDOWS <=2%/<=5% rank; display favors intended overlap, then allele diversity, then rank. Internal red ticks are pre-binding cut evidence; all predictions remain in CSV.