

WWC1 p.Val179Gly | JLF V3 | 20 aa

disclosed minimal epitope GHLQHELQF at 8-16

INTRACELLULAR / CLASS-I PROCESSING - four complementary score tracks

higher native score upward

NetChop Cterm (ligand) native 0-1

0.5 display

Pepsickle human (in vivo) native 0-1

0.5 display

NetCleave I native 0-1

0.5 display

NetChop 20S (in vitro) native 0-1

0.5 display

ERAP1 / ERAMER

cytosol / ER motifs

NPEPP5

TPP2

+10 more in CSV

MHC-I candidate ligands

C07:01 0.082%

C01:02 0.67%

B27:05 0.56%

N

K

L

K

R

E

M

G

H

L

Q

H

E

L

Q

F

K

E

R

G

MHC-II: none <= 5% rank

1

5

10

15

20

ENDOLYSOSOMAL / CLASS-II PROCESSING - C-terminal score track

higher native score downward

NetCleave II native 0-1

0.5 display

SERUM / EXTRACELLULAR - native scores and recognition motifs

DPP4 native (intact SLP)

0.065

serum / extracellular motifs

MME

MME

MME

MME

ACE

SCORES Native 0-1 bond scores; lines join adjacent assessed bonds only (no smoothing). Dashed line = 0.5 display threshold; gaps = unassessed.

RED BOND MARKS All four intracellular tracks assess the bond; >=3 reach 0.5. Four beads show support (filled = hit, open = miss), not probability.

MHC WINDOWS <=2%/<=5% rank; display favors intended overlap, then allele diversity, then rank. Internal red ticks are pre-binding cut evidence; all predictions remain in CSV.